Figure 1 Implanting the spacer in the gap that is firmly secured by the traction.

method to provide traction at the knee joint while inserting a static spacer has involved manual traction. This is an impractical method. We hereby describe a handy and easy-to-apply alternative technique using an ankle sling known from ankle arthroscopy to create a steady traction at the knee joint.

TECHNIQUE

After removal of infected knee implants and debridement of infected tissue, an ankle sling is placed around the foot with the components positioned behind the tendo-Achilles and over the dorsum of the foot. The pulling string of the sling is attached ventrally to a belt around the waist of the assistant situated at the patient's feet. When the assistant leans backwards, the tibia and femur are efficiently pulled apart. The surgeon can now implant the spacer in the gap that is firmly secured by the traction (Fig. 1). The method is comfortable for both surgeon and assistant. Maintaining tension and alignment, while the cement sets, is easy.

DISCUSSION

This is an easy method to obtain proper limb alignment and force of traction for insertion of a static spacer in the knee joint. The advantages are: (i) the traction created is steady and easily applied; (ii) it is inexpensive; and (iii) it secures a suitably shaped and sized spacer. Preformed moulds for creating moveable spacers intra-operatively are commercially available. These have the advantage of allowing early ROM-exercises, but these moveable spacers are often impractical.

A method preventing circlage wire displacement in intra-operative proximal femoral fractures

S HOSKINSON, I GILL, P FADERO

Department of Orthopaedics and Trauma Surgery, Mayday University Hospital, Croydon, UK

CORRESPONDENCE TO

Simon Hoskinson, Department of Orthopaedics, Mayday University Hospital, 530 London Road, Croydon CR7 7YE, UK
E: simon.hoskinson@mayday.nhs.uk

BACKGROUND

Intra-operative fractures of the proximal femur can occur during hip hemi-arthroplasty. One method of management is with circlage wires which prevents fracture propagation and allows fixation.¹ However, due to the funnel shape of the proximal femur, wires can slip on the femoral shaft. We describe a method of circlage wire placement to prevent this.

TECHNIQUE

This technique requires a 2.5-mm drill-bit, and an appropriate diameter of wire (18-gauge). A hole is drilled within the cortex of the calcar in an anterior-posterior direction, the wire is then passed through the drill hole and looped around the proximal femoral shaft twice and tied (Fig. 1). Effort must be made to ensure that the drill hole is placed entirely within the cortex and does not breach the intramedullary canal. If this occurs, the wire will be passing through the canal and have the potential to affect stem fixation.

DISCUSSION

This method has the advantage of anchorage of the wire loop through bone preventing wire slippage and fracture displacement.

Figure 1 Sawbone model showing wire *in situ* around proximal femur. (A) Anterior-posterior view; (B) lateral view.

This method is ideally suited to standard hemi-arthroplasty implants. We recognise the potential for canal compromise with poorly placed wires which may affect implant fixation. However, in cemented implants a wire which breaches the canal will be surrounded by cement mantle and its effect will be negligible. With uncemented implants, which rely on metaphyseal fixation for primary and secondary stability, the presence of the wire within the canal may cause micromotion and compromise osseointegration.

Reference

1. Berend KR, Lombardi Jr AV, Mallory TH, Chonko DJ, Dodds KL, Adams JB. Cerclage wires or cables for the management of intraoperative fracture associated with a cementless, tapered femoral prosthesis: results at 2 to 16 years. *J Arthroplasty* 2004; **19** (Suppl 2): 17–21.

The use of over-penetrated radiographs for assessment of chronic osteomyelitis in children

H WYNN JONES¹, J BATES², WJ HARRISON¹

¹Department of Orthopaedics, Beit Cure International Hospital, Blantyre, Malawi

²Department of Orthopaedics, Queen Elizabeth Central Hospital, Blantyre, Malawi

CORRESPONDENCE TO:

Henry Wynn Jones, Senior Orthopaedic Fellow, Beit Cure International Hospital, PO Box 31236, Blantyre 3, Malawi
T: +265 (0)1875015; F: +265 (0)1874397;
E: hwj@doctors.org.uk

BACKGROUND

Childhood chronic osteomyelitis is a condition that remains prevalent in the developing world. There are several imaging techniques that are useful in the investigation and management of chronic osteomyelitis such as plain radiographs, scintigraphy (bone, leukocyte and gallium), computer tomography, magnetic resonance imaging and positron emission tomography. Comparison between these has shown that some of the more sophisticated imaging techniques have some advantages.¹ However, plain radiographs are the only means of imaging that are generally available in the developing world, where this condition is most common.

TECHNIQUE

We have found that deliberately over-penetrated radiographs often reveal sequestra and abscesses that are not seen on radiographs taken at a standard exposure. This is because they are masked by the thick involucrum, which is frequently sclerotic. The radiographs in Figures 1 and 2 show anterior-posterior radiographs of the right tibia and fibula of a child with chronic haematogenous osteomyelitis. Figure 1 shows a

Figure 1 Anterior-posterior radiograph of the tibia of a boy with chronic haematogenous osteomyelitis taken at a standard exposure.

Figure 2 The same tibia seen on an overexposed radiograph: the sequestra are visualised much more clearly.

radiograph taken at a standard exposure (12 mAs/84 kVp) and Figure 2 deliberately overexposed (12 mAs/88 kVp). The sequestrae are much more clearly seen in Figure 2.