Recognition of Depression by Non-psychiatric Physicians—A Systematic Literature Review and Meta-analysis

Monica Cepoiu, MD, MSc^{1,2}, Jane McCusker, MD, DrPH^{1,3}, Martin G. Cole, MD, FRCP(C)^{4,5}, Maida Sewitch, PhD^{1,6}, Eric Belzile, MSc¹, and Antonio Ciampi, PhD^{1,3}

¹Department of Clinical Epidemiology and Community Studies, St. Mary's Hospital, Montreal, QC, Canada; ²Department of Community Health Studies, University of Calgary, Calgary, AB, Canada; ³Department of Epidemiology and Biostatistics, McGill University, Montreal, QC, Canada; ⁴Department of Psychiatry, St. Mary's Hospital, Montreal, Canada; ⁵Department of Psychiatry, McGill University, Montreal, Canada; ⁶Department of Medicine, McGill University, Montreal, QC, Canada.

BACKGROUND: Depression, with up to 11.9% prevalence in the general population, is a common disorder strongly associated with increased morbidity. The accuracy of non-psychiatric physicians in recognizing depression may influence the outcome of the illness, as unrecognized patients are not offered treatment for depression.

OBJECTIVES: To describe and quantitatively summarize the existing data on recognition of depression by non-psychiatric physicians.

METHODS: We searched the following databases: MEDLINE (1966–2005), Psych INFO (1967–2005) and CINAHL (1982–2005). To summarize data presented in the papers reviewed, we calculated the Summary receiver operating characteristic (ROC) and the summary sensitivity, specificity and odds ratios (ORs) of recognition, and their 95% confidence intervals using the random effects model.

MEASUREMENTS AND MAIN RESULTS: The summary sensitivity, specificity, and OR of recognition using the random effects model were: 36.4% (95% CI: 27.9–44.8), 83.7% (95% CI: 77.5–90.0), and 4.0 (95% CI: 3.2–4.9), respectively. We also calculated the Summary ROC. We performed a metaregression analysis, which showed that the method of documentation of recognition, the age of the sample, and the date of study publication have significant effect on the summary sensitivity and the odds of recognition, in the univariate model. Only the method of documentation had a significant effect on summary sensitivity, when the age of the sample and the date of publication were added to the model.

CONCLUSION: The accuracy of depression recognition by non-psychiatrist physicians is low. Further research should focus on developing standardized methods of documenting non-psychiatric physicians' recognition of depression.

KEY WORDS: recognition; depression; meta-analysis.

J Gen Intern Med 23(1):25–36

DOI: 10.1007/s11606-007-0428-5

© Society of General Internal Medicine 2007

BACKGROUND

Depression, with up to 11.9% prevalence in the general population,^{1,2} is a common disorder strongly associated with increased morbidity.³ It has been estimated that in 2020 depression will become the second leading cause of disability,⁴ which emphasizes the importance of its early detection and treatment. The accuracy of non-psychiatric physicians in recognizing depression may influence the outcome of the illness, as unrecognized patients are not offered treatment for depression. Currently, less than half of patients with depression are recognized by their primary care physicians, even after 5 years of follow-up.⁵ Moreover, studies show that among recognized depressed patients only a few receive appropriate care, which may further lead to poor outcome of depression and increased health service use and mortality rates in these patients.^{6,7}

There are many potential reasons for the underrecognition and undertreatment of depression; patient, provider, and system barriers have been identified. Patients reduce the likelihood of being diagnosed by presenting with somatic rather than emotional complaints^{8–10} and may resist a diagnosis of depression or anxiety by attributing their symptoms to physical causes.^{11,12} Provider barriers include concerns about potential patient stigma,^{13,14} time pressures,¹⁴ a belief that such diagnoses are burdensome,¹⁵ inadequate knowledge about diagnostic criteria or treatment options,⁶ lack of a psychosocial orientation,^{16–18} and inadequate insight into different cultural presentations of mental disorders.¹⁹ System barriers include productivity pressures, limitations of third-party mental health coverage, restrictions on specialist, drug, and psychotherapeutic care,^{13,14} lack of a systematic method for detecting and managing such patients,²⁰ and inadequate continuity of care.¹³

The accuracy of recognition of depression by attending physicians can be assessed using measures such as sensitivity, specificity, and odds ratio, when the clinical diagnosis made by the physician is compared to a gold standard diagnosis of depression. The estimates of these measures reported in the literature vary according to the method used

Received April 3, 2007

Revised September 29, 2007

Accepted October 4, 2007

Published online October 26, 2007

to document the clinical diagnosis made by physicians and the definition of gold standard.

The main objectives of this literature review and meta-analysis were to describe and quantitatively summarize existing data on recognition of depression by non-psychiatric physicians in adult outpatients and inpatients. To our knowledge, this is the first systematic review of studies on recognition of depression by non-psychiatric physicians.

METHODS

Search Strategy

The search was conducted by MC (MSc in Epidemiology) using OVID search engine (2000–2005 version). MEDLINE (1966–2006), Psych INFO (1967–2006), CINAHL (1982–2006), and EMBASE (1986–2006) databases were searched for the following keywords: “depression” or “depression disorder” and “detection” or “recognition” or “identification” or “diagnosis”. Letters, editorials, reviews, and case-reports were excluded. The search was limited to studies with subjects of ages 18 and more and written in English or French. This search yielded 7,105 papers.

Inclusion and Exclusion Criteria

Titles and abstracts of these papers were reviewed to identify those of potential relevance, based on the following inclusion criteria:

1. The study samples were adult patients attending primary care facilities, hospital emergency departments or outpa-

tient clinics, or admitted to hospital in either medical or surgical wards.

2. A gold standard diagnosis of depression was made by a study psychiatrist or by research staff, using a structured clinical interview or a rating scale with a specified cut point.
3. A clinical diagnosis of depression (or other method of recognition, such as antidepressant prescription, referral to a mental health specialist, or identification of depressive symptoms) was made by a non-psychiatric physician.

Eighty-one studies of potential relevance were retained for review. The reference lists of the papers selected were searched and another 2 studies of potential relevance were retained for review. From the 85 reviewed papers, we excluded 49 for the following reasons:

1. In 13 studies, no gold standard diagnosis of depression was used to evaluate the validity of physicians' recognition of depression.^{21–33}
2. Twenty-three studies evaluated the recognition of depression by nurses or the recognition of depression was attributed to a health care team (including physicians, nurses, social workers, etc).^{34–56} The large body of literature looking at the recognition of depression by other health professionals than physicians justifies a separate review of this subject.
3. In 7 studies, recognition of depression by physicians was evaluated in training cases (vignettes of patients, actors, or videotaped patients).^{18,57–62}
4. Two studies were randomized controlled trials that evaluated educational programs for improving recognition of depression by physicians.^{63,64}

Figure 1. The systematic review flow

Table 1. Studies that Used the Chart Review Method (Sample Characteristics, Physician Specialties, and Methods)

Author/ Year	Sample size (% depressed)	Age (years)	Physician specialties	Methods			
				Method of sampling	Blinding of physicians	Criterion measure for depression	Method and criteria of recognition of depression by physician
Asch 2003 ⁸⁶	1,140 (39.3%)	Not specified	HIV specialists, general internists, gynecologists	Sampling rates	Yes	CIDI (DSM-IV)	<i>Criteria:</i> diagnosis of depression
Balestrieri 2002 ⁸⁷	309 (63%)	45–65	Medical and surgical physicians	Not specified	Yes	CIDI-PHC	<i>Criteria:</i> any of the following – prescription of psychotropic drugs – any statements about the presence of depression – referral to psychiatrist <i>Criteria:</i> diagnosis of depression
Bertakis 2001 ⁷³	508 (25.6%)	Not specified	Family practice physicians and general internal medicine residents	Consecutive	Not specified	BDI—cut point of 9	<i>Criteria:</i> diagnosis of depression
Callahan 1997 ⁸⁸	508 (25.6%)	19–92	Family practice and general internal medicine residents	Consecutive	Yes	BDI—cut point of 9	<i>Criteria:</i> diagnosis of depression
Crawford 1998 ⁷⁶	318 (19.5%)	>65	General practitioners	Consecutive	Not specified	Short CARE scale— cut point of 6	<i>Criteria:</i> active treatment = antidepressant or referral to psychiatrist, psychiatric nurse or social services. <i>Criteria:</i> – diagnosis of depression – one or more visits to a mental health specialist – prescription for one or more antidepressant medication
Garrard 1998 ⁸⁹	3410 (16.4%)	>65	All specialties	Consecutive	Yes	GDS—cut point of 11	<i>Criteria:</i> any of the following – diagnosis of depression – use of word 'depression' – use of words descriptive of depression ('blue', 'sad', 'dysphoric')
Lichtenberg 1993 ⁹⁰	150 (34%) 90 Female 60 Male	>60	Emergency physicians	Consecutive	Yes	GDS—cut point of 10	<i>Criteria:</i> any of the following – note of depression – prescription of antidepressant medication <i>Criteria:</i> referral to mental health evaluation
McCusker 1996 ¹⁰³	94 (52.1%)	>60	Primary care physicians	Random	Yes	GDS—cut point of 11	<i>Criteria:</i> any of the following – diagnosis of depression – term 'depression' or 'depressed' mentioned – prescription of antidepressant medication <i>Criteria:</i> any of the following – depressed mood mentioned in the discharge diagnoses – depressed mood mentioned in the discussion section – antidepressant or benzodiazepine prescribed at discharge
Meldon 1997 ¹⁰⁴	101 (29.7%)	>65	Emergency physicians	Consecutive	Yes	SRDS—cut point of 4	
Meldon 1997 ¹⁰⁵	259 (27%)	>65	Emergency physicians	Convenience sample	Yes	KS—cut point of 4	
Nuyen 2005 ⁷⁸	191 (28.8%)	>18	General practitioners	Random	Not specified	CIDI (DSM-IV)	<i>Criteria:</i> code of depression in the Dutch National Survey database
Perez- Stable 1990 ⁹¹	265 (26.4%)	18–69	Not specified	Not specified	Yes	DIS—DSM III	<i>Criteria:</i> any of the following – diagnosis of depression – term 'depression' or 'depressed' mentioned – prescription of antidepressant medication
Pouget 2000 ⁹²	401 (22.4%)	>75	Not specified	Alternate	Yes	15-item GDS— cut point of 6	<i>Criteria:</i> any of the following – depressed mood mentioned in the discharge diagnoses – depressed mood mentioned in the discussion section – antidepressant or benzodiazepine prescribed at discharge

(continued on next page)

Table 1. (continued)

Author/ Year	Sample size (% depressed)	Age (years)	Physician specialties	Methods			
				Method of sampling	Blinding of physicians	Criterion measure for depression	Method and criteria of recognition of depression by physician
Rapp 1988 ⁹³	150 (15.3%)	>65	Not specified	Random	Yes	SADS – evaluated using Research Diagnostic Criteria (RDC)	<i>Criteria:</i> any of the following – a formal diagnosis of depression – use of the word ‘depressed’ in the chart – use of words descriptive of depression, such as ‘dysphoric’, ‘sad’, or ‘despondent’ – any stated need for initiation or continuation of treatment for emotional or mental distress
Volkers 2004 ⁷⁴	237 (23.2%)	>55	General practitioners	Random	Not specified	CIDI (DSM-IV)	<i>Criteria:</i> diagnosis of depression, depressive symptoms (down/depressed feelings) in the electronic medical record database
Whooley 1997 ⁹⁴	429 (15.8%)	39–67	Attending physicians and resident physicians	Consecutive	Yes	DIS—Quick DIS-III-R	<i>Criteria:</i> any of the following – term ‘depression’ or ‘depressed’ was noted – referral of the patient to a psychiatrist for further evaluation of depressive symptoms
Zung 1983 ⁹⁵	1086 (13.2%)	>20	Family practitioners	Random	Yes	SRDS—cut point of 55	<i>Criteria:</i> notations regarding depression and its treatment

*These studies presented results for both methods.

BDI = Beck Depression Inventory, CARE = Comprehensive Assessment and Referral Evaluation, CIDI = Composite International Diagnostic Interview, CIDI-PHC = Composite International Diagnostic Interview—Primary Care, DIS = Diagnostic Interview Schedule, DSM = Diagnostic and Statistical Manual of Mental Disorders, GDS = Geriatric Depression Scale, KS = Koenig Scale, SADS = Schedule for Affective Disorders and Schizophrenia, SRDS = Zung Self Rating Depression Scale

- In 3 studies,^{65–67} data were not available to calculate at least the overall sensitivity.
- In one study,⁶⁸ recognition of depression was based on patients’ self-report on antidepressant medication, with no reference to the specialty of the physician who prescribed the medication (possibly prescribed by psychiatrists).

The flow chart of the systematic review is presented in Figure 1.

The papers selected based on our inclusion and exclusion criteria were independently reviewed by 2 reviewers with training in Epidemiology using a review form and guidelines developed by the authors. Data abstracted included: study design, study setting, method of selecting the sample, age and gender of the sample, sample size used in the analysis, specialties and number of physicians participating in the study, blinding of physicians, criterion measure for diagnosing depression, data source for diagnosing depression, data source and indicators of recognition, and results (sensitivity, specificity, positive predictive value, negative predictive value, crude odds ratio, other) either presented in the study or possibly calculated from the data available.

Quality Assessment

Differences in data abstraction between the 2 reviewers were discussed and resolved by the authors. We reviewed the following aspects of study quality: selection of a clinically

relevant cohort, the consistent use of a good gold standard, blinding of physicians to the gold standard results, and incomplete reporting.⁶⁹

In all the studies reviewed, the authors consistently used a gold standard tool for detecting depression symptoms or diagnosing depression. Moreover, details pertaining to the validity and reliability of the tool used were included.

The authors of all the studies reviewed indicated the method of sampling the patients (consecutive or randomized sampling) and the health care settings where the patients were recruited. The age or gender distribution, as well as the prevalence of depression in the samples recruited from different health care settings vary considerably.

In 11 studies,^{70–80} the authors did not specify if the physicians were blinded to the gold standard diagnosis (incomplete reporting). The rest of the studies included references to the blinding of physicians or the chart review.

To determine whether the quality of the studies may influence the conclusions of our literature review and meta-analysis, we performed a sensitivity analysis that explored the following aspects of study quality:

- Method of sampling—selecting patients in a nonrandom manner may lead to selection bias.
- Characteristics of the study sample (prevalence of depression, age and gender distribution)—the variation of these characteristics by study setting may introduce spectrum bias.⁸¹

3. Blinding of physicians—the physicians' awareness of the gold standard diagnosis may improve the accuracy of recognition, leading to incorporation bias.⁸²

Statistical Analysis

We calculated missing sensitivities, specificities, and crude odds ratios (OR) using data reported in the papers reviewed. We used the random effects model⁸³ to calculate the summary diagnostic odds ratios, sensitivity, and specificity. Because there was evidence of varying cut points (Spearman: -0.635 , $p=0.0005$), we also plotted our main measure and recalculated the overall sensitivity using receiver operating characteristic (ROC) curves.⁸⁴

We performed the Cochrane's Q test⁸⁵ to assess the presence of heterogeneity among results reported in studies included in our systematic literature review and meta-analysis. We performed a univariate metaregression analysis, testing 7 potential sources of heterogeneity: the method of documentation (chart review vs physician diagnosis), blinding of physicians (physicians were blinded vs blinding status was not specified), the gold standard diagnostic tools used (structured clinical interview vs diagnostic scale with a specific cut point), method of sampling (random vs other), age (55 and over vs all ages and younger patients only), gender of sample (62% and more vs less than 62% female), date of study publication (after 1998 vs 1998 and before) and prevalence of depression (less than 25% vs 25% or more). The categories of the last 3 covariates were created using the median value as cut point. These analyses were performed using the STATA statistical software (STATA/SE 8.1 for Windows).

RESULTS

We included 36 studies in our systematic review (Tables 1, 2, 3, and 4). Twenty-three papers reported only sensitivity^{72-74,78-80,86-102}. For 10 of these studies, we calculated the specificity and the diagnostic OR.^{72,73,80,87,89,91,93,99-101} In 8 papers, both the sensitivity and specificity of recognition were presented.^{75,77,103-108} Eleven papers reported other measures (such as positive predictive value, agreement expressed as percentage or kappa correlation coefficient, and identification index),^{70,76,77,79,103,104,106,107,109,110} and we calculated the sensitivity, specificity, and OR of recognition using data presented in 5 of them.^{70,76,77,79,109} One paper included in our systematic review¹⁰⁵ reported a sensitivity of 0% and a specificity of 98.9%. We considered these results outliers and decided to exclude them from our analysis.

Twenty-seven (75%) of the studies included in our systematic review and meta-analysis were conducted in primary care,^{70-79,80,86,88,89,94-100,102,103,107-110} 3 studies were conducted in the emergency department,^{90,104,105} three studies included patients admitted to the hospital,^{87,92,93} and three studies included outpatients attending specialty clinics.^{70,91,101}

Overall, we found high specificity (83.7%, 95% CI: 77.5–90.0, Table 5), but lower sensitivity (36.4%, 95% CI: 27.9–44.8) with a resulting diagnostic OR of 4.0 (95% CI: 3.2–4.9). The sensitivity varied across the method of assessment (Table 5), with physician diagnosis (PD) method having higher pooled

Table 2. Studies that used the Chart Review Method (Results)

Author/Year	Main results			
	Sens. (%)	Spec. (%)	DOR	Other†
Asch 2003 ⁸⁶	46			
Balestrieri 2002 ⁸⁷	32.5	92.1*	5.5*	
Bertakis 2001 ⁷³	27.7	89.1*	3.1*	
Callahan 1997 ⁸⁸	27.7			
Crawford 1998 ⁷⁶	25.8*	89.8*	3*	
Garrard 1998 ⁸⁹	52%	77.4*	3.7*	
Lichtenberg 1993 ⁹⁰	Overall: 27 Female: 40 Male: 10			
McCusker 1996 ¹⁰³	26	92	3.8*	PPV=50% Overall agreement:78%
Meldon 1997 ¹⁰⁴	13*	89*	1.2*	
Meldon 1997 ¹⁰⁵	0	98.9	0	
Nuyen 2005 ⁷⁸	28.8			
Perez-Stable 1990 ⁹¹	35.7	81.5*	2.4*	
Pouget 2000 ⁹²	16.7			
Rapp 1988 ⁹³	8.7	95.2*	1.9*	
Volkers 2004 ⁷⁴	20.8			
Whooley 1997 ⁹⁴	8.8			
Zung 1983 ⁹⁵	15			

*Sensitivities (Sens.), specificities (Spec.) and crude diagnostic odds ratios (DOR) not reported by the authors were computed from the data available.

†PPV = positive predictive value

sensitivity than those using chart review (CR). Papers published after 1998 had higher pooled sensitivity than those before 1998 (Table 5). When we calculated the overall sensitivity, based on summary ROC curves, we found a sensitivity of 42.3% (Fig. 2).

Our results were heterogeneous. On metaregression, the method of documentation, age of the sample, and date of publication showed a statistically significant effect on the summary sensitivity (Table 6). These results indicate that the summary sensitivity of studies that used as method of documentation the physician diagnosis, were published after 1998 and had a sample of younger or all ages of patients was higher by 14.5%, 11.8%, and 12.3%, respectively, compared to those that used as method of documentation the chart review, were published in 1998 or before, and had a sample of patients aged 55 and more.

Only age of the sample and date of publication explained the heterogeneity of our pooled odds ratios. Studies that were published after 1998 and had a sample of younger or all ages of patients reported higher odds ratios of recognition compared to studies that were published in 1998 or before and had a sample of patients aged 55 and more (Table 7).

We performed a multivariate metaregression, including in the model 3 variables that had a significant effect on summary sensitivity or the pooled odds ratios in the univariate analysis: method of documentation, age of the sample, and date of publication. The multivariate analysis showed that the summary sensitivity of studies that used as method of documentation the physician diagnosis was 12.5% higher compared to those that used as method of documentation the chart review, when controlled for age of the sample and date of publication (Table 7).

Table 3. Studies that Used the Physician Diagnosis Method (Sample Characteristics, Physician Specialties, and Methods)

Author/Year	Sample size (% depressed)	Age (years)	Physician specialties	Methods			
				Method of sampling	Blinding of physicians	Criterion measure for depression	Method and criteria of recognition of depression by physician
Aragones 2004 ⁹⁶	306 (39.2%)	18–70	General practitioners	Consecutive	Yes	SRDS—cut point of 55	<i>Criteria:</i> answers of “yes” and “possible yes” on the questionnaire asking about patients’ depression
Balestrieri 2004 ¹⁰⁹	2093 (13.5%)	Not specified	Primary care practitioners	Consecutive	Yes	PHQ—cut point=9	<i>Criteria:</i> the physician filled out a form indicating the patient’s current depression, current antidepressant treatment and previous episodes of depression
Becker 2004 ¹¹⁰	431 (19.9%)	18–80	Primary care doctors	Consecutive	Yes	PHQ-9 (cut point not specified)	<i>Criteria:</i> rated patients as cases of depression
Berardi 2005 ⁷²	361 (44.3%)	>14	Primary care physicians	Random	Yes	WHO ICD-10 Symptom Checklist for Depression	<i>Criteria:</i> clinical diagnosis of depression
Bowers 1990 ⁷⁵	101 (14.8%)	≥ 70	General practitioner	Consecutive	Not specified	Diagnostic interview for depression (DSM-III-R)	<i>Criteria:</i> moderate and severe depression
Coyne 1995 ⁹⁷	143 (100%)	Mean age 39.7	Family physicians	Consecutive	Not specified	SCID (DSM-III-R)	<i>Criteria:</i> affirmative response to the question regarding patients’ state of depression
Crawford 1998 ⁷⁶	318 (19.5%)	>65	General practitioners	Consecutive	Not specified	Short CARE scale—cut point of 6	<i>Criteria:</i> in a face- to-face interview physicians were asked if their patients were depressed or prone to depression
Christensen 2003 ¹⁰⁸	301 (30.2)	Mean age 38.8	General practitioners	Consecutive	Yes	SCAN	<i>Criteria:</i> GPs were asked whether the patient suffered from depression
Klinkman 1998 ⁷⁷	372 (21.7%)	Mean age 39.6	Family physicians	Consecutive	Not specified	SCID (DSM-III-R)	<i>Criteria:</i> clinically significant depression
Katon 2004 ⁹⁸	4385 (12%)	Mean age 59	Not specified	Consecutive	Yes	PHQ-9—cut point of 5	<i>Criteria:</i> – diagnosis of depression – antidepressant treatment – specialty mental health visit
McCusker 1996 ¹⁰³	94 (52.1%)	>60	Primary care physicians	Random	Yes	GDS—cut point of 11	<i>Criteria:</i> any of the following – note of depression – prescription of antidepressant medication
Meldon 1997 (a) ¹⁰⁴	101 (29.7%)	>65	Emergency physicians	Consecutive	Yes	SRDS—cut point of 4	<i>Criteria:</i> note of depressive symptoms.
Passik 1998 ⁷⁰	1105 (35.8%)	Majority 50–70	Oncologists	Not specified	Not specified	SRDS—cut point of 50	<i>Criteria:</i> scores 4 to 10 on the physician rating scale
Pfaff 2005 ⁹⁹	916 (23.8%)	>60	General practitioners	Consecutive	Yes	CES-D—cut point of 16	<i>Criteria:</i> presence of symptoms of depression
Pond 1990 ¹⁰⁰	133 (14.3%)	70–84	General practitioners	Random	Yes	GDS—cut point of 10	<i>Criteria:</i> note of depressive symptoms.
Shulman 2002 ¹⁰¹	101 (43.6%)	45–84	Neurologists	Consecutive	Not specified	BDI—cut point 10	<i>Criteria:</i> depressive symptoms reported by physicians

(continued on next page)

Table 3. (continued)

Author/Year	Sample size (% depressed)	Age (years)	Physician specialties	Methods			
				Method of sampling	Blinding of physicians	Criterion measure for depression	Method and criteria of recognition of depression by physician
Simon 1999 ⁷⁹	948 (100%)	<65	Primary care physicians	Random	Not specified	CIDI (ICD-10 criteria)	<i>Criteria:</i> both recognition of psychological “caseness” and assignment of an appropriate diagnosis
Sliman 1992 ¹⁰⁶	420 (25.2%)	16–94	Internal medical residents	Consecutive	Yes	BDI—cut point of 6	<i>Criteria:</i> a score of 7 or more on physician's rating
Stek 2004 ¹⁰²	77 (100%)	>85	General practitioners	Consecutive	Yes	15-item GDS-S: cut point of 5	<i>Criteria:</i> note of depressive symptoms.
Thompson 2001 ¹⁰⁷	18 414 (19.9%)	16–94	General practitioners	Consecutive	Yes	HAD scale—cut point of 8	<i>Criteria:</i> clinically significant depressive illness—mild, moderate or severe
Tiemens 1999 ¹¹⁶	709 (24.5%)	18–65	Primary care physicians	Random	Yes	CIDI-PHC	<i>Criteria:</i> score >2 on the Physician's Encounter Form
Wittchen 2001 ⁸⁰	19106 (11.5%)	15–99	Primary care physicians	Consecutive	Not specified	DSQ—cut point of 8 (ICD-10) or 10 (DSM-IV)	<i>Criteria:</i> presence of depression

*These studies presented results for both methods.

BDI = Beck Depression Inventory, CARE = Comprehensive Assessment and Referral Evaluation, CES-D = Center for Epidemiological Studies Depression Scale, CIDI = Composite International Diagnostic Interview, CIDI-PHC = Composite International Diagnostic Interview—Primary Care, DSQ = Depression Screening Questionnaire, GDS = Geriatric Depression Scale, HAD = Hospital Anxiety and Depression Scale, ICD = International Classification of Diseases, PHQ = Personal Health Questionnaire, PHQ-9 = Patient Health Questionnaire, SCAN = Schedules for Clinical Assessment in Neuropsychiatry, SCID = Structural Clinical Interview for Depression, SRDS = Zung Self Rating Depression Scale

DISCUSSION

Recognition of depression by nonpsychiatric physicians has received increased attention from researchers as depression became one of the most prevalent diseases of the 21st century and an important public health issue. The assessment of validity of recognition is challenged by the variety of methods used to document recognition and to diagnose depression. In this systematic literature review, we qualitatively and quantitatively summarized the accuracy of recognition of depression using data presented in the papers reviewed. Further, we tried to identify sources of heterogeneity in the results reported.

The summary sensitivity showed that less than half of the depressed patients are recognized by their physicians. This is consistent with rates of depression detection reported in previous studies.⁹ On the other hand, the summary specificities are reasonable (calculated using data reported in 22 studies) and are consistent with interrater agreement found in studies on the accuracy of psychiatrist interviews.¹¹¹

Studies that used as method of documentation the physician diagnosis had a higher summary sensitivity compared to studies that used the chart review. This result is consistent with the conclusion of a study that compared the 2 methods,¹⁰³ showing that either physicians tend to diagnose depression more frequently when they are specifically asked about this possible diagnosis or documentation of depression recognition (including

diagnosis, treatment, and referral to a mental health specialist) in patients' charts is low. Moreover, the same study found that even if the sensitivity of recognition by physician diagnosis was higher, the specificity of this method of documentation was lower than the specificity of recognition by chart review. This may explain why the method of documentation did not affect the summary odds ratio in our metaregression analysis.

We also found that the sensitivity and odds of recognition of depression by physicians are significantly higher in younger or unselected patients than in older ones. Data reported in several studies included in our literature review show no significant association between recognition of depression and age^{88,107} or higher recognition of depression in patients less than 35 compared to patients aged 65 and more,¹¹² which demonstrates the variability among studies. However, in the multivariate metaregression, the age and the sample had no effect on summary sensitivity and summary OR when controlled for method of documentation and date of publication.

Another notable finding of our meta-analysis is that studies published after 1998 tend to report higher sensitivities and odds ratios of recognition than those published in 1998 and before. This may suggest that the non-psychiatric physicians training in diagnosing depression has improved over the years. However, in the multivariate metaregression, the date of publication had no effect on summary sensitivity and summary OR when controlled for method of documentation and age of the sample.

Table 4. Studies that Used the Chart Review Method (Results)

Author/Year	Main results			
	Sens.	Spec.	DOR	Other†
Aragones 2004 ⁹⁶	72%			
Balestrieri 2004 ¹⁰⁹	38.8%*	93.7%*	9.4*	Identification index=0.3
Becker 2004 ¹¹⁰	48.8%*	89.5%*	8.2*	Agreement kappa=0.4
Berardi 2005 ⁷²	79.4%	48.2%*	3.6*	
Bowers 1990 ⁷⁵	20%	90%	2.1*	
Coyne 1995 ⁹⁷	27.9%			
Crawford 1998 ⁷⁶	51.6%*	71.8%*	2.7* (PD)	Kappa of agreement=0.19 (0.08–0.29)
Christensen§ 2003 ¹⁰⁸	40.6	94.7	6.76*	
Klinkman§ 1998 ⁷⁷	34.9%	92.9%	4.7*	PPV=44.6%
Katon 2004 ⁹⁸	51.1%			
McCusker 1996 ¹⁰³	67%	81%	8.8*	PD: PPV=55%
				Overall agreement:78% PPV=32%
Meldon 1997 (a) ¹⁰⁴	27%	75%	1.1	
Passik 1998 ⁷⁰	43.6%*	78.9%*	2.9*	Kappa of agreement=0.17
Pfaff 2005 ⁹⁹	39.9%	81.2%*	2.8*	
Pond 1990 ¹⁰⁰	21%	91.2%*	2.8*	
Shulman 2002 ¹⁰¹	35%	89.5%*	4.4*	
Simon 1999 ⁷⁹	36%			
Sliman 1992 ¹⁰⁶	46.2%	84.4%	4.6*	PPV=50% NPV=82.3% Pearson's correlation coefficient=0.42
Stek 2004 ¹⁰²	22%			
Thompson 2001 ¹⁰⁷	36.1%	91.5%	6.1*	Kappa=0.31 (0.28, 0.33)
Tiemens 1999 ¹¹⁶	40.2%*	85.8%*	4.0*	Total percentage agreement: 86.6% Kappa=0.29
Wittchen 2001 ⁸⁰	75% (DSM) 59% (ICD-10) 85.1%* (either DSM-IV or ICD-10)	60.4%* (either DSM-IV or ICD-10)	8.7* (either DSM-IV or ICD-10)	

*Sensitivities (Sens.), specificities (Spec.) and crude diagnostic odds ratios (DOR) not reported by the authors were computed from the data available.

†PPV = positive predictive value, NPV = negative predictive value

‡These studies presented results for both methods.

§The sensitivity, specificity, and OR reported by the authors were weighted estimates. We calculated the crude estimates of sensitivity, specificity, and OR using data reported in the papers (Christensen 2003: 41.7%, 85.4% and 4.2, respectively; Klinkman 1998: 61.7%, 88.3%, and 4.7, respectively). We used these crude estimates in our meta-analysis.

We identified several limitations of our literature review and meta-analysis. First, the literature search was restricted to two languages. Is it possible that articles written in other languages than English or French were missed. Second, it was difficult to

Table 5. Summary Sensitivities, Specificities, and ORs of Recognition, with 95% CI (Random Effects Model)

	Sensitivity		Specificity	OR
All studies	N=38* 36.4 (27.9, 44.8)	N=25* 39.2 (28.0, 50.6)	N=25* 83.7 (77.5, 90.0)	N=25* 4.0 (3.3, 4.9)
CR method	N=15 26.6 (18.4, 34.9)	N=8 28.2 (16.6, 39.8)	N=8 88.1 (82.2, 94.0)	N=8 3.5 (2.9, 4.1)
PD method	N=23 42.9 (31.6, 54.1)	N=17 44.7 (30.4, 58.9)	N=17 81.6 (73.4, 89.8)	N=17 4.4 (3.5, 5.6)
Age 55+	N=15† 28.7 (19.8, 37.5)	N=11 31.4 (20.6, 42.1)	N=11 85.1 (80.1, 90.0)	N=11 2.9 (2.4, 3.7)
All ages	N=20 41.1 (28.3, 53.9)	N=12 46.7 (28.5, 64.9)	N=12 80.6 (68.9, 92.3)	N=12 4.5 (3.5, 5.8)
Published after 1998	N=19 42.1 (29.8, 54.4)	N=11 45.9 (27.6, 64.1)	N=11 82.5 (72.0, 93.0)	N=11 5.2 (4.1, 6.6)
Published in 1998 and before	N=19 30.2 (22.5, 37.8)	N=14 33.8 (25.9, 41.7)	N=14 84.8 (80.8, 88.9)	N=14 3.2 (2.6, 3.8)

*Three studies^{76,103,104} contributed each with two sets of data.

†In three studies,^{73,86,109} data on the age of the sample were not available.

assess the quality of the papers included, as there was a great variability in methods used. We abstracted and presented data regarding the sites of the studies, the method of sampling, the blinding of outcomes, and the specialties of physicians involved in these studies, to offer a view of the methodological quality of these studies that may have affected the validity of our results. Third, our subgroup analysis did not cover all the possible sources of heterogeneity (for example, the categorization in subgroups by the type of gold standard diagnostic tool used does not account for the variability in type of depression scales and cut points used). Fourth, many studies included as "missed" patients those with subthreshold depression, a category of depression in which treatment and placebo both have

Figure 2. Summary ROC curve

similar high rates of remission¹¹³ and which tend to be transient.⁹ Fifth, there was no gradation on severity of depression. Studies have shown that patients with more severe forms of depression are more likely to be diagnosed.^{9,79,80,107,114,115}

CONCLUSION

The overall sensitivity of recognition of depression by non-psychiatric physicians reported in our systematic literature review and meta-analysis was low, although non-psychiatric physicians had good specificity. A number of variables, including the method of documentation of recognition, age of the sample, and date of study publication, had an impact on the summary sensitivity. Moreover, the last 2 factors had a significant effect on the summary odds ratio of recognition, a measure of diagnostic accuracy.

A large number of potential barriers to recognition and treatment of depression have been identified. The specific reason studies consistently find low rates of sensitivities is unclear. Given the high prevalence of the disease and its

Table 6. Effect of Sources of Heterogeneity on Summary Sensitivity of Recognition and OR of Recognition (Univariate Metaregression)

Covariate	Sensitivity of recognition N=38*		OR of recognition N=25*	
	β -coefficient	p value	β -coefficient	p value
Method of documentation (physician diagnosis vs chart review)	0.145	.004	0.350	.125
Blinding of physicians (blinded vs not specified)	-0.030	.603	-0.047	.835
"Gold standard" diagnosis test (structured clinical interview vs depression scale)	0.024	.663	0.017	.936
Method of sampling (random vs other)	-0.053	.386	-0.095	.739
Gender (62% and more vs less than 62% female)	-0.005	.932	-0.098	.649
(Missing)	(5)		(4)	
Age (55 and over vs all ages and younger patients only)	-0.123	.039	-0.499	.011
(Missing)	(4)		(3)	
Date of publication (after 1998 vs in 1998 and before)	0.118	.024	0.541	.003
Prevalence of depression (less than 25% vs 25% or higher)	0.028	.637	-0.352	.070

*Three studies^{76,103,104} contributed each with two sets of data.

Table 7. Effect of Sources of Heterogeneity on Summary Sensitivity of Recognition and OR of Recognition (Multivariate Metaregression)

Covariate	Sensitivity of recognition N=34*		OR of recognition N=22 *	
	β - coefficient	p value	β - coefficient	p value
Method of documentation (physician diagnosis versus chart review)	0.125	.038	-0.075	.748
Age (55 and over versus all ages and younger patients only)	-0.086	.131	-0.037	.079
Date of publication (after 1998 versus in 1998 and before)	0.052	.387	0.340	.096

*Three studies^{76,103,104} contributed each with two sets of data. In three studies,^{73,86,109} data on the age of the sample was not available.

significant impact on the overall health of patients, this question deserves further research. It would be helpful if there were a standardized method of documenting non-psychiatric physicians' recognition of depression. This method may be useful for studies that tests the efficacy of various educational programs designed to improve physicians' accuracy in diagnosing depression. Also, a standardized method of documenting recognition may improve the quality of studies that look at various factors which may affect recognition of depression, such as age, gender or ethnicity of patients, or physicians' characteristics.

Acknowledgments: This study was supported by a grant from Fonds de la recherche en santé du Québec (Grant number 25004-2560) and two grants from Institutes for Health Research (Grant numbers MOP-64462 and 11949).

Conflict of Interest: None disclosed.

Corresponding Author: Monica Cepoiu, MD, MSc; Department of Community Health Studies, University of Calgary, Rm G8, 3330 Hospital Drive, Calgary, AB, Canada, T2N 4N1 (e-mail: monica.cepoiucalgary.ca).

REFERENCES

1. Berardi D, Menchetti M, De Ronchi RD, Rucci P, Leggieri G, Ferrari G. Late-life depression in primary care: a nationwide Italian epidemiological survey. *J Am Geriatr Soc*. 2002;50(1):77-83.
2. Murphy JM, Laird NM, Monson RR, Sobol AM, Leighton AH. A 40-year perspective on the prevalence of depression: the stirling county study. *Arch Gen Psychiatry*. 2000;57(3):209-15.
3. Wells KB, Sturm R, Sherbourne CD, Meredith LS. *Caring for Depression*. Cambridge, MA, US: Harvard University Press; 1996.
4. Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause, 1990-2020: Global Burden of Disease Study. *Lancet*. 1997;349(9064):1498-504.
5. Jackson JL, Passamonti M, Kroenke K. Outcome and impact of mental disorders in primary care at 5 years. *Psychosom Med*. 2007;69(3):270-6.
6. Davidson JRT, Meltzer-Brody SE. The underrecognition and undertreatment of depression: what is the breadth and depth of the problem? *J Clin Psychiatry*. 1999;60(suppl. 7):4-11.

7. **Cole MG, Bellavance F.** Depression in elderly medical inpatients: a meta-analysis of outcomes. *Can Med Assoc J.* 1997;157(8):1055-60.
8. **Wittchen HU, Lieb R, Wunderlich U, Schuster P.** Comorbidity in primary care: presentation and consequences. *J Clin Psychiatry.* 1999;60(suppl 7):29-36.
9. **Jackson JL, O'Malley PG, Kroenke K.** Clinical predictors of mental disorders among medical outpatients. Validation of the "S4" model. *Psychosomatics.* 1998;39(5):431-6.
10. **O'Connor DW, Rosewarne R, Bruce A.** Depression in primary care. 1: Elderly patients' disclosure of depressive symptoms to their doctors. *Int Psychogeriatr.* 2001;13(3):359-65.
11. **Herran A, Vazquez-Barquero JL, Dunn G.** Recognition of depression and anxiety in primary care. Patients' attributional style is important factor. *BMJ.* 1999;318(7197):1558.
12. **Kessler D, Lloyd K, Lewis G, Gray DP.** Cross sectional study of symptom attribution and recognition of depression and anxiety in primary care. *BMJ.* 1999;318(7181):436-9.
13. **Docherty JP.** Barriers to the diagnosis of depression in primary care. *J Clin Psychiatry.* 1997;58(suppl 1):5-10.
14. **Goldman LS, Nielsen NH, Champion HC.** Awareness, diagnosis, and treatment of depression. *J Gen Intern Med.* 1999;14(9):569-80.
15. **Wittchen HU, Pittrow D.** Prevalence, recognition and management of depression in primary care in Germany: the Depression 2000 study. *Human Psychopharmacol.* 2002;17(suppl 1):1-11.
16. **Parchman ML.** Physicians' recognition of depression. *Fam Pract Res J.* 1992;12(4):431-8.
17. **Cooper LA, Brown C, Vu HT, Palenchar DR, Gonzales JJ, Ford DE, Powe NR.** Primary care patients' opinions regarding the importance of various aspects of care for depression. *Gen Hosp Psychiatry.* 2000;22(3):163-73.
18. **Carney PA, Eliassen MS, Wolford GL, Owen M, Badger LW, Dietrich AJ.** How physician communication influences recognition of depression in primary care. *J Fam Pract.* 1999;48(12):958-64.
19. **Kirmayer LJ.** Cultural variations in the clinical presentation of depression and anxiety: implications for diagnosis and treatment. [Review] [77 refs]. *J Clin Psychiatry.* 2001;62(suppl 13):22-8.
20. **McCall L, Clarke DM, Rowley G.** A questionnaire to measure general practitioners' attitudes to their role in the management of patients with depression and anxiety. *Aust Fam Physician.* 2002;31(3):299-303.
21. **Tai-Seale M, Bramson R, Drukker D, et al.** Understanding primary care physicians' propensity to assess elderly patients for depression using interaction and survey data. *Med Care.* 2005;43(12):1217-24.
22. **Hedayati SS, Grambow SC, Szczech LA, Stechuchak KM, Allen AS, Bosworth HB.** Physician-diagnosed depression as a correlate of hospitalizations in patients receiving long-term hemodialysis. *Am J Kidney Dis.* 2005;46(4):642-9.
23. **Buist A, Bilszta J, Barnett B, et al.** Recognition and management of perinatal depression in general practice—a survey of GPs and postnatal women. *Aust Fam Physician.* 2005;34(9):787-90.
24. **Richards HL, Fortune DG, Weidmann A, Sweeney SK, Griffiths CE.** Detection of psychological distress in patients with psoriasis: low consensus between dermatologist and patient. *Br J Dermatol.* 2004;151(6):1227-33.
25. **Furedi J, Rozsa S, Zambori J, Szadoczky E.** The role of symptoms in the recognition of mental health disorders in primary care. *Psychosomatics* 2003;44(5):402-6.
26. **Evers MM, Samuels SC, Lantz M, Khan K, Brickman AM, Marin DB.** The prevalence, diagnosis and treatment of depression in dementia patients in chronic care facilities in the last six months of life. *Int J Geriatr Psychiatry.* 2002;17(5):464-72.
27. **Harman JS, Schulberg HC, Mulsant BH, Reynolds CF, III.** The effect of patient and visit characteristics on diagnosis of depression in primary care. *J Fam Pract.* 2001;50(12):1068.
28. **Luber MP, Hollenberg JP, Williams-Russo P, et al.** Diagnosis, treatment, comorbidity, and resource utilization of depressed patients in a general medical practice. *Int J Psychiatry Med.* 2000;30(1):1-13.
29. **Lecrubier Y.** Is depression under-recognised and undertreated? *Int Clin Psychopharmacol.* 1998;13(suppl. 5):S3-S6.
30. **Schmidt LA, Greenberg BD, Holzman GB, Schulkin J.** Treatment of depression by obstetrician-gynecologists: a survey study. *Obstet Gynecol.* 1997;90(2):296-300.
31. **Strothers HS III, Rust G, Minor P, Fresh E, Druss B, Satcher D.** Disparities in antidepressant treatment in Medicaid elderly diagnosed with depression. *J Am Geriatr Soc.* 2005;53(3):456-61.
32. **Rout U, Rout J.** Diagnosis and treatment of depression by general practitioners in England. *Psychol Rep.* 1996;78(2):516-8.
33. **Crystal S, Sambamoorthi U, Walkup JT, Akincigil A.** Diagnosis and treatment of depression in the elderly medicare population: predictors, disparities, and trends. *J Am Geriatr Soc.* 2003;51(12):1718-28.
34. **Webber AP, Martin JL, Harker JO, Josephson KR, Rubenstein LZ, Alessi CA.** Depression in older patients admitted for postacute nursing home rehabilitation. *J Am Geriatr Soc.* 2005;53(6):1017-22.
35. **Eisses A-M, Kluiters H, Jongenelis K, Pot AM, Beekman ATF, Ormel J.** Care staff training in detection of depression in residential homes for the elderly: randomised trial. *Br J Psychiatry.* 2005;186:404-9.
36. **Preville M, Cote G, Boyer R, Hebert R.** Detection of depression and anxiety disorders by home care nurses. *Aging Ment Health.* 2004;8(5):400-9.
37. **Brown EL, Bruce ML, McAvay GJ, Raue PJ, Lachs MS, Nassisi P.** Recognition of late-life depression in home care: accuracy of the outcome and assessment information set. *J Am Geriatr Soc.* 2004;52(6):995-9.
38. **Dobalian A, Tsao JCI, Radcliff TA.** Diagnosed mental and physical health conditions in the United States nursing home population: differences between urban and rural facilities. *J Rural Health.* 2003; 19(4):477-83.
39. **Barton C, Miller B, Yaffe K.** Evaluation of the diagnosis and management of cognitive impairment in long-term care. *Alzheimer Dis Assoc Disord.* 2003;17(2):72-6.
40. **Leo RJ, Sherry C, Jones AW.** Referral patterns and recognition of depression among African-American and Caucasian patients. *Gen Hosp Psych.* 1998;20(3):175-82.
41. **Teresi JA, Abrams R, Holmes D, Ramirez M, Shapiro C, Eimicke JP.** Influence of cognitive impairment, illness, gender, and African-American status on psychiatric ratings and staff recognition of depression. *Am J Geriatr Psychiatry.* 2002;10(5):506-14.
42. **Watts SC, Bhutani GE, Stout IH, et al.** Mental health in older adult recipients of primary care services: is depression the key issue? Identification, treatment and the general practitioner. *Int J Geriatr Psychiatry.* 2002;17(5):427-37.
43. **Teresi J, Abrams R, Holmes D, Ramirez M, Eimicke J.** Prevalence of depression and depression recognition in nursing homes. *Soc Psychiatry Psychiatr Epidemiol.* 2001;36(12):613-20.
44. **Rincon HG, Granados M, Unutzer J, et al.** Prevalence, detection and treatment of anxiety, depression, and delirium in the adult critical care unit. *Psychosomatics.* 2001;42(5):391-6.
45. **Bagley H, Cordingley L, Burns A, et al.** Recognition of depression by staff in nursing and residential homes. *J Clin Nurs.* 2000;9(3):445-50.
46. **McDonald MV, Passik SD, Dugan W, Rosenfeld B, Theobald DE, Edgerton S.** Nurses' recognition of depression in their patients with cancer. *Oncol Nurs Forum.* 1999;26(3):593-9.
47. **Shah A, De T.** Documented evidence of depression in medical and nursing case-notes and its implications in acutely ill geriatric inpatients. *Int Psychogeriatr.* 1998;10(2):163-72.
48. **Boyle VL, Roychoudhury C, Beniak R, Cohn L, Bayer A, Katz I.** Recognition and management of depression in skilled-nursing and long-term care settings: evolving targets for quality improvement. *Am J Geriatr Psychiatry.* 2004;12(3):288-95.
49. **Bell M, Goss AJ.** Recognition, assessment and treatment of depression in geriatric nursing home residents. *Clin Excell Nurse Pract* 2001;5(1):26-36.
50. **Halloran E, Prentice N, Murray CL, et al.** Follow-up study of depression in the elderly. Clinical and SPECT data. *Br J Psychiatry.* 1999;175:252-8.
51. **Boey KW.** Detection of geriatric depression: knowledge and practice of hospital nurses. *Clin Gerontol.* 1999;20(2):47-56.
52. **Sullivan K, Dyck J, Culp KR, Buckwalter K.** Comparing the geriatric depression scale, minimum data set, and primary care provider diagnosis for depression in rural nursing home residents. *J Am Psychiatr Nurses Assoc.* 2005;11(5):270-75.
53. **Brown EL, McAvay G, Raue PJ, Moses S, Bruce ML.** Recognition of depression among elderly recipients of home care services. *Psychiatr Serv.* 2003;54(2):208-13.
54. **Koenig HG, Goli V, Shelp F, Kudler HS, Cohen HJ, Blazer DG.** Major depression in hospitalized medically ill older men: documentation, management, and outcome. *Int J Geriatr Psychiatry.* 1992;7(1):25-34.
55. **Bowler C, Boyle A, Branford M, Cooper S-A, Harper R, Lindesay J.** Detection of psychiatric disorders in elderly medical inpatients. *Age Ageing.* 1994;23(4):307-11.
56. **Huffman JC, Smith FA, Blais MA, Beiser ME, Januzzi JL, Fricchione GL.** Recognition and Treatment of Depression and Anxiety in Patients With Acute Myocardial Infarction. *Am J Cardiol.* 2006;98(3):319-24.

57. **Helmes E, Duggan GM.** Assessment of depression in older adult males by general practitioners. Ageism, physical problems and treatment. [see comment]. *Aust Fam Phys.* 2001;30(3):291-4.
58. **Tylee A, Freeling P, Kerry S, Burns T.** How does the content of consultations affect the recognition by general practitioners of major depression in women? *Br J Gen Pract.* 1995;45(400):575-8.
59. **Passik SD, Donaghy KB, Theobald DE, Lundberg JC, Holtsclaw E, Dugan J.** Oncology staff recognition of depressive symptoms on videotaped interviews of depressed cancer patients: implications for designing a training program. *Journal of Pain and Symptom Management.* 2000;19(5):329-38.
60. **Stoppe G, Sandholzer H, Huppertz C, Duwe H, Staedt J.** Family physicians and the risk of suicide in the depressed elderly. *J Affect Disord.* 1999;54(1-2):193-8.
61. **Penn JV, Boland R, McCartney JR, Kohn R, Mulvey T.** Recognition and treatment of depressive disorders by internal medicine attendings and housestaff. *Gen Hosp Psychiatry.* 1997;19(3):179-84.
62. **Hasin D, Link B.** Age and recognition of depression: implications for a cohort effect in major depression. *Psychol Med.* 1988;18(3):683-8.
63. **Gerrity MS, Cole SA, Dietrich AJ, Barrett JE.** Improving the recognition and management of depression: is there a role for physician education? *J Fam Pract.* 1999;48(12):949-57.
64. **Dowrick C, Buchan I.** Twelve month outcome of depression in general practice: does detection or disclosure make a difference? *Br Med J.* 1995;311(7015):1274-6.
65. **Gallo JJ, Bogner HR, Straton JB, et al.** Patient characteristics associated with participation in a practice-based study of depression in late life: the spectrum study. *Int J Psychiatry Med.* 2005;35(1):41-57.
66. **Bellantuono C, Mazzi MA, Tansella M, Rizzo R, Goldberg D.** The identification of depression and the coverage of antidepressant drug prescriptions in Italian general practice. *J Affect Disord.* 2002;72(1):53-9.
67. **Bogner HR, Ford DE, Gallo JJ.** The role of cardiovascular disease in the identification and management of depression by primary care physicians. *Am J Geriatr Psychiatry.* 2006;14(1):71-8.
68. **Weintraub D, Moberg PJ, Duda JE, Katz IR, Stern MB.** Recognition and treatment of depression in Parkinson's disease. *J Geriatr Psychiatry Neurol.* 2003;16(3):178-83.
69. **Deeks JJ.** Systematic reviews in health care: systematic reviews of evaluations of diagnostic and screening tests. *BMJ.* 2001;323(7305):157-62.
70. **Passik SD, Dugan W, McDonald MV, Rosenfeld B, Theobald DE, Edgerton S.** Oncologists' recognition of depression in their patients with cancer. *J Clin Oncol.* 1998;16(4):1594-600.
71. **Schulman E, Gairola G, Kuder L, McCulloch J.** Depression and associated characteristics among community-based elderly people. *J Allied Health.* 2002;31(3):140-6.
72. **Berardi D, Menchetti M, Cevenini N, Scaini S, Versari M, De RD.** Increased recognition of depression in primary care: comparison between primary-care physician and ICD-10 diagnosis of depression. *Psychother Psychosom.* 2005;74(4):225-30.
73. **Bertakis KD, Helms LJ, Callahan EJ, Azari R, Leigh P, Robbins JA.** Patient gender differences in the diagnosis of depression in primary care. *J Women's Health Gend-Based Med.* 2001;10(7):689-98.
74. **Volkers AC, Nuyen J, Verhaak PFM, Schellevis FG.** The problem of diagnosing major depression in elderly primary care patients. *J Affect Disord.* 2004;82(2):259-63.
75. **Bowers J, Jorm AF, Henderson S, Harris P.** General practitioners' detection of depression and dementia in elderly patients. *Med J Aust.* 1990;153(4):192-6.
76. **Crawford MJ, Prince M, Menezes P, Mann AH.** The recognition and treatment of depression in older people in primary care. *Int J Geriatr Psychiatry.* 1998;13(3):172-6.
77. **Klinkman MS, Coyne JC, Gallo S, Schwenk TL.** False positives, false negatives, and the validity of the diagnosis of major depression in primary care. *Arch Fam Med.* 1998;7(5):451-61.
78. **Nuyen J, Volkers AC, Verhaak PFM, Schellevis FG, Groenewegen PP, Van den Bos GAM.** Accuracy of diagnosing depression in primary care: the impact of chronic somatic and psychiatric co-morbidity. *Psychol Med.* 2005;35(8):1185-95.
79. **Simon GE, Goldberg SD, Tiemens BG, Ustun TB.** Outcomes of recognized and unrecognized depression in an international primary care study. *Gen Hosp Psychiatry.* 1999;21(2):97-105.
80. **Wittchen H-U, Hofler M, Meister W.** Prevalence and recognition of depressive syndromes in German primary care settings: poorly recognized and treated? *Int Clin Psychopharmacol.* 2001;16(3):121-35.
81. **Goehring C, Perrier A, Morabia A.** Spectrum bias: a quantitative and graphical analysis of the variability of medical diagnostic test performance. *Stat Med.* 2004;23(1):125-35.
82. **Ransohoff DF, Feinstein AR.** Problems of spectrum and bias in evaluating the efficacy of diagnostic tests. *N Engl J Med.* 1978;299(17):926-30.
83. **DerSimonian R, Laird N.** Meta-analysis in clinical trials. *Control Clin Trials.* 1986;7(3):177-88.
84. **Moses LE, Shapiro D, Littenberg B.** Combining independent studies of a diagnostic test into a summary ROC curve: data-analytic approaches and some additional considerations. *Stat Med.* 1993;12(14):1293-316.
85. **Cooper H, Hedges LV.** *The Handbook of Research Synthesis.* New York, NY, US: Russell Sage Foundation; 1994.
86. **Asch SM, Kilbourne AM, Gifford AL, et al.** Underdiagnosis of depression in HIV: who are we missing? *J Gen Intern Med.* 2003;18(6):450-60.
87. **Balestrieri M, Bisoffi G, Tansella M, Martucci M, Goldberg DP.** Identification of depression by medical and surgical general hospital physicians. *Gen Hosp Psychiatry.* 2002;24(1):4-11.
88. **Callahan EJ, Bertakis KD, Azari R, Helms LJ, Robbins J, Miller J.** Depression in primary care: patient factors that influence recognition. *Fam Med.* 1997;29(3):172-6.
89. **Garrard J, Rolnick SJ, Nitz NM, et al.** Clinical detection of depression among community-based elderly people with self-reported symptoms of depression. *J Gerontol Ser A Biol Sci Med Sci.* 1998;53(2):M92-M101.
90. **Lichtenberg PA, Gibbons TA, Nanna M, Blumenthal F.** Physician detection of depression in medically ill elderly. *Clin Gerontol.* 1993;13(1):81-90.
91. **Perez-Stable EJ, Miranda J, Munoz RF, Ying Y-W.** Depression in medical outpatients. Underrecognition and misdiagnosis. *Arch Intern Med.* 1990;150(5):1083-8.
92. **Pouget R, Yersin B, Wietlisbach V, Burnand B, Bula CJ.** Depressed mood in a cohort of elderly medical inpatients: Prevalence, clinical correlates and recognition rate. *Aging Clin Exp Res.* 2000;12(4):301-7.
93. **Rapp SR, Walsh DA, Parisi SA, Wallace CE.** Detecting depression in elderly medical inpatients. *J Consult Clin Psychol.* 1988;56(4):509-13.
94. **Whooley MA, Avins AL, Miranda J, Browner WS.** Case-finding instruments for depression: two questions are as good as many. *J Gen Int Med.* 1997;12(7):439-45.
95. **Zung WKK, Magill M, Moore JT, George DT.** Recognition and treatment of depression in a family medicine practice. *J Clin Psychiatry.* 1983;44(1):3-6.
96. **Aragones E, Pinol JL, Labad A, Folch S, Melich N.** Detection and management of depressive disorders in primary care in Spain. *Int J Psychiatry Med.* 2004;34(4):331-43.
97. **Coyne JC, Schwenk TL, Fechner-Bates S.** Nondetection of depression by primary care physicians reconsidered. *Gen Hosp Psychiatry.* 1995;17(1):3-12.
98. **Katon WJ, Simon G, Russo J, et al.** Quality of depression care in a population-based sample of patients with diabetes and major depression. *Med Care.* 2004;42(12):1222-9.
99. **Pfaff JJ, Almeida OP.** A cross-sectional analysis of factors that influence the detection of depression in older primary care patients. *Aust N Z J Psychiatry.* 2005;39(4):262-5.
100. **Pond CD, Mant A, Bridges-Webb C, et al.** Recognition of depression in the elderly: a comparison of general practitioner opinions and the geriatric depression scale. *Fam Pract.* 1990;7(3):190-4.
101. **Shulman LM, Taback RL, Rabinstein AA, Weiner WJ.** Non-recognition of depression and other non-motor symptoms in Parkinson's disease. *Parkinsonism Relat Disord.* 2002;8(3):193-7.
102. **Stek ML, Gussekloo J, Beekman ATF, Van TW, Westendorp RGJ.** Prevalence, correlates and recognition of depression in the oldest old: The Leiden 85-plus study. *J Affect Disord.* 2004;78(3):193-200.
103. **McCusker J, Karp E, Yaffe MJ, Cole M, Bellavance F.** Determining detection of depression in the elderly by primary care physicians: chart review or questionnaire? *Prim Care Psychiatry.* 1996;2(4):217-21.
104. **Meldon SW, Emerman CL, Schubert DSP.** Recognition of depression in geriatric ED patients by emergency physicians. *Ann Emerg Med.* 1997;30(4):442-7.
105. **Meldon SW, Emerman CL, Schubert DSP, Moffa DA, Etheart RG.** Depression in geriatric ED patients: prevalence and recognition. *Ann Emerg Med.* 1997;30(2):141-5.
106. **Sliman RJ, Donohue TA, Jarjoura D, Ognibene AJ.** Recognition of depression by internal medicine residents. *J Commun Health.* 1992;17(3):143-52.

107. **Thompson C, Ostler K, Peveler RC, Baker N, Kinmonth A.** Dimensional perspective on the recognition of depressive symptoms in primary care: the Hampshire depression project 3. *Br J Psychiatry*. 2001;179:317–23.
108. **Christensen KS, Toft T, Frostholm L, Ornbol E, Fink P, Olesen F.** The FIP Study: a randomised, controlled trial of screening and recognition of psychiatric disorders. *Br J Gen Pract*. 2003;53(495):758–63.
109. **Balestrieri M, Carta MG, Leonetti S, Sebastiani G, Starace F, Bellantuono C.** Recognition of depression and appropriateness of antidepressant treatment in Italian primary care. *Soc Psychiatry Psychiatr Epidemiol*. 2004;39(3):171–6.
110. **Becker SM.** Detection of somatization and depression in primary care in Saudi Arabia. *Soc Psychiatry Psychiatr Epidemiol*. 2004;39(12):962–6.
111. **Spitzer RL, Forman JB, Nee J.** DSM-III field trials: I. Initial interrater diagnostic reliability. *Am J Psych* 1979;136(6):815–7.
112. **Borowsky SJ, Rubenstein LV, Meredith LS, Camp P, Jackson-Triche M, Wells KB.** Who is at risk of nondetection of mental health problems in primary care? *J Gen Intern Med*. 2000;15(6):381–8.
113. **Barrett JE, Williams JW Jr, Oxman TE, et al.** Treatment of dysthymia and minor depression in primary care: a randomized trial in patients aged 18 to 59 years. *J Fam Pract*. 2001;50(5):405–12.
114. **Ronalds C, Creed F, Stone K, Webb S, Tomenson B.** Outcome of anxiety and depressive disorders in primary care. *Br J Psychiatry*. 1997;171:427–33.
115. **Pini S, Berardi D, Rucci P, et al.** Identification of psychiatric distress by primary care physicians. *Gen Hosp Psych*. 1997;(6):411–8.
116. **Tiemens BG, Vonkorff M, Lin EHB.** Diagnosis of depression by primary care physicians versus a structured diagnostic interview: understanding discordance. *Gen Hosp Psych*. 1999;21(2): 87–96.