Published in final edited form as:

Neurobiol Learn Mem. 2008 January ; 89(1): 81–85.

Age-Related Changes in Contextual Associative Learning

Trinh T. Luu¹, Eva Pirogovsky², and Paul E. Gilbert^{1,2}

¹ Department of Psychology, San Diego State University, San Diego, CA

² San Diego State University-University of California San Diego-, SDSU-UCSD Joint Doctoral Program in Clinical Psychology, San Diego, CA

Abstract

The hippocampus plays a critical role in processing contextual information. Although age-related changes in the hippocampus are well documented in humans, nonhuman primates, and rodents, few studies have examined contextual learning deficits in old rats. The present study investigated age-related differences in contextual associative learning in young (6 mo) and old (24 mo) rats using olfactory stimuli. Stimuli consisted of common odors mixed in sand and placed in clear plastic cups. Testing was conducted in two boxes that represented two different contexts (Context 1 and Context 2). The contexts varied based on environmental features of the box such as color (black vs. white), visual cues on the walls of the box, and flooring texture. Each rat was simultaneously presented with two cups, one filled with Odor A and one filled with Odor B in each context. In Context 1, the rat received a food reward for digging in the cup containing Odor A, but did not receive a food reward for digging in the cup containing Odor B. In Context 2, the rat was rewarded for digging in the cup containing Odor B, but did receive a reward for digging in the cup containing Odor A. Therefore, the rat learned to associate Context 1 with Odor A and Context 2 with Odor B. The rat was tested for eight days using the same odor problem throughout all days of testing. The results showed no significant difference between young and old rats on the first two days of testing; however, young rats significantly outperformed old rats on Day 3. Young rats continued to maintain superior performance compared to old rats on Days 4–8. The results suggest that aging results in functional impairments in brain regions that support memory for associations between specific cues and their respective context.

Keywords

Aging; Context; Hippocampus; Olfactory; Associative Learning; Fischer 344 Brown Norway

The contextual features of an episode play a critical role in learning and memory. Research has suggested that the hippocampus may be essential for processing contextual information. For example, disruptions of hippocampal function impair contextual fear conditioning (Anagnostaras, Maren, & Fanselow, 1999) and appetitive conditioning to contextual stimuli (Good & Honey, 1991). Although fear conditioning to the environmental context relies on the hippocampus, fear conditioning to a discrete stimulus (e.g. a tone) is not hippocampal-dependent (Phillips & LeDoux, 1992). Recent studies utilizing novelty-preference and paired-associate learning paradigms also have demonstrated that hippocampal damage impairs the

Please address all correspondence to Dr. Paul Gilbert, SDSU-UCSD Joint Doctoral Program in Clinical Psychology, 6363 Alvarado Court, Suite 103, San Diego, CA 92120-4913. Tel: (619) 594-7409; Fax: (619) 594-3773; Email: pgilbert@sciences.sdsu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

acquisition and retention of contextual stimuli (Mumby, Gaskin, Glenn, Schramek, & Lehmann, 2002; Rajji, Chapman, Eichenbaum, & Greene, 2006). In addition, spatial firing patterns in hippocampal neurons demonstrate significant changes when the context is altered, including changes in the spatial layout, color, and odor of an environment as well as changes in non-environmental features such as task demands (Anderson & Jeffery, 2003; Smith & Mizumori, 2006; Wood, Dudchenko, Robitsek, & Eichenbaum, 2000).

There is an extensive literature from human and animal research implicating the medial temporal lobes, and particularly the hippocampus, in age-related deficits in learning and memory (for review see Gallagher & Rapp, 1997). Recent studies have reported that there are similar numbers of principal neurons in the hippocampus in young and aged rats (Rasmussen, Schliemann, Sorensen, Zimmer, & West, 1996); however, aged animals exhibit changes in the functional connectivity of the hippocampus (for review see Barnes, 1994). In addition, normal aging is associated with impairments on tasks that rely on intact functioning of the hippocampus and surrounding regions. For example, aged nonhuman primates and rodents demonstrate parallel impairments to animals with hippocampal damage in a variety of memory tasks, including tasks measuring spatial memory (Barnes, 1994; Gallagher, Burwell, & Burchinal, 1993), delayed recognition memory (Dunnett, Evenden, & Iversen, 1988; Rapp & Amaral, 1991), odor memory (Rapp, Kansky, & Eichenbaum, 1996), and transitive inference (Rapp et al., 1996).

A number of studies have shown age-related changes in fear conditioning to context (Houston, Stevenson, McNaughton, & Barnes, 1999; Moyer & Brown, 2006; Oler & Markus, 1998). Similar to animals with hippocampal lesions, older rats show impaired contextual fear conditioning compared to younger rats, but normal fear conditioning to a tone (Houston et al., 1999; Oler & Markus, 1998). However, few if any studies have investigated age-related changes in contextual memory using an appetitive-based paradigm. The present study investigated contextual associative learning in young and aged rats using an appetitive task developed by Rajji et al. (2006).

Twenty Fischer 344/Brown Norway (Harlan Laboratories) male rats 6 mo of age ($n=10$) and 24 mo of age ($n=10$) were used as test subjects. All rats were housed in individual cages and all testing took place during the light portion of a 12-hr light-dark cycle. The rats had unlimited access to water, but were maintained on a restricted diet in order to maintain 85–90% of their free-feeding weight. All procedures complied with guidelines established by the Institution of Animal Care and Use Committee.

Testing was conducted in two clear Plexiglas boxes ($18 \times 16 \times 15$ cm) that represented two different contexts, termed Context 1 and Context 2. A context was defined as the total of all of the environmental cues in the apparatus, including floor texture, color of walls, and visual cues on the walls. In Context 1, the flooring was black and textured. The walls were black with thin white stripes (see Figure 1). In Context 2, the flooring was white and smooth. The walls were white with a different black geometric shape (i.e., star, triangle, square, circle) on each wall (see Figure 1).

Prior to testing, each rat was shaped in the homecage to dig in a clear plastic cup filled with unscented sand to receive a food reward. Following the shaping period, each rat was tested on a contextual learning task developed by Rajji et al. (2006). In this task, rats must learn to associate one odor with one context and a second odor with a different context. Olfactory stimuli consisted of common odors: cinnamon, cumin, ginger and baby powder. Odors were mixed in sand and placed in small clear plastic cups (6.5 cm diameter and 6 cm high) as described in previous studies (Bunsey & Eichenbaum, 1996; Gilbert & Kesner, 2003). In each context, the rat was presented simultaneously with two cups, each filled with a different odorant

(Odor A and Odor B). The cups were presented adjacent to one another with an 8 cm space between each cup as shown in Figure 1. In Context 1, the rat received a food reward for digging in the cup containing Odor A, but did not receive a food reward for digging in the cup containing Odor B. In Context 2, the rat received a food reward for digging in the cup containing Odor B, but did not receive a food reward for digging in the cup containing Odor A. If the rat dug in the cup containing the rewarded odor, the trial was scored as a correct response. However, if the rat dug in the cup containing the unrewarded odor, the trial was scored as an incorrect response and the rat was not allowed to dig in the other cup to receive a reward. Therefore, each rat must learn to associate Context 1 with Odor A and Context 2 with Odor B. The left/right position of each cup and the order in which rats were presented with each context followed a pseudorandom order. Each rat performed 10 trials per day with five trials in each context across eight consecutive days. The same odor problem was presented throughout all days of testing. Each trial was separated by a 15–20 min intertrial interval.

The mean ($\pm$ SE) number of correct responses on the contextual learning task as a function of days (1–8) for the 6 mo and 24 mo old rats are shown in Figure 2. A 2×8 analysis of variance (ANOVA) with group (6 mo, 24 mo) as the between-group factor and day (days 1–8) as the within group factor was used to analyze the data. The analysis revealed a statistically significant main effect of group, $F(1, 18) = 27.58, p < .001$ and day $F(7, 126) = 4.61, p < .001$. In addition, the analysis revealed a statistically significant group $\times$ day interaction, $F(7, 126) = 3.66, p < .01$. A Newman-Keuls post hoc comparison test of the statistically significant group $\times$ day interaction did not detect significant differences between young and old rats on the first two days of testing. However, young rats significantly ($p < .05$) outperformed old rats on Day 3. Young rats continued to perform better than old rats on Days 4–7 and significantly outperformed old rats on Day 8 ($p < .05$). The post hoc analysis of the group $\times$ day interaction also indicated that 6 mo old rats performed significantly better on Days 3–8 compared to Day 1 ($p < .05$), indicating learning across the eight days of testing. However, there were no significant differences among any days of testing in the 24 mo old rats.

Each rat then was tested on an olfactory discrimination task to rule out that any deficits in 24 mo old rats on the contextual learning task were due to an inability to discriminate between odors. Each rat was tested on the task in a neutral context with gray walls and a gray smooth floor. Odor discrimination was assessed using a two-choice discrimination task described by Brushfield et al. (2006). Each rat was trained to discriminate between the two odors used during the contextual learning task. For each rat, one odor was randomly assigned as the rewarded odor and the other odor was assigned as the non-rewarded odor. On each trial, the two odors were presented simultaneously side-by-side and the rat was allowed to choose between the two odor cups. If the rat dug in the odor cup containing the rewarded odor, the rat received a food reward. However, if the rat dug in the odor cup containing the non-rewarded odor, the rat did not receive a reward and was not allowed to dig in the odor cup containing the rewarded odor. The position of each odor varied pseudo-randomly on each trial with respect to the left and right position to eliminate position bias. Each rat received 12 trials per day and was tested until the animal reached a criterion of nine correct choices out of a sliding block of 10 consecutive trials within the 12 trials.

The mean ($\pm$ standard error) number of trials to criterion on the olfactory discrimination task were 24.9 (3.22) for the 6 mo old rats and 25.3 (4.33) for the 24 mo old rats. A one-way ANOVA with group (6 mo, 24 mo) as the between-group factor and mean trials to criterion as the dependent measure did not reveal a statistically significant main effect of group, $F(2, 17) = .47, p = .85$. Therefore, the data suggest that 24 mo old rats could discriminate between the odors used in the contextual learning task as well as 6 mo old rats.

The present study investigated age-related changes in associative memory for context using an appetitive task. The data revealed that 6 mo old rats performed at chance on the first day of testing but significantly outperformed the 24 mo old rats by the third day of testing. The 24 mo old rats performed just above chance across all eight days of testing. These results suggest that associative memory for contextual information may be particularly sensitive to age-related changes in the brain. The results of the present study are consistent with prior reports of age-related impairments in contextual fear conditioning (Houston et al., 1999; Moyer & Brown, 2006; Oler & Markus, 1998). However, to the authors' knowledge, this is the first study to demonstrate impaired contextual memory in aging rats using an appetitive task.

Many previous behavioral studies involving aged rats have used Fisher 344 rats (F344). The present study was conducted using a Fischer 344/Brown Norway (F344/BN) hybrid strain. The F344/BN strain has been shown to live longer than Fischer 344 rats. The 50% survival age for F344/BN male rats is 34 mo, whereas the 50% survival age for F344 male rats is 24 mo (National Institute on Aging). Therefore, the rats in the present study showed significant learning impairments at 24 months of age despite having longer average longevity than other strains of rats used in prior behavioral experiments.

Performance of the present contextual associative learning task required that rats were able to discriminate between the two odors used during testing. Studies have reported that olfactory sensitivity is decreased in aged rats relative to young rats (Kraemer & Apfelbach, 2004). Therefore, it is important to demonstrate that any deficit in learning the contextual associative learning task is not the result of an inability to discriminate between the two olfactory stimuli presented in each context. Similar to prior studies, the results of the olfactory discrimination task demonstrate that 24 mo old rats can discriminate between the odors as well as 6 mo old rats (Brushfield et al., 2006; Kraemer & Apfelbach, 2004; Schoenbaum, Nugent, Saddoris, & Gallagher, 2002). Therefore, age-related impairments in contextual learning observed in present task were not solely due to an inability to discriminate between the olfactory stimuli. Furthermore, a prior study in the laboratory using the same strain of rats demonstrated that 24 mo old rats can perform visual discriminations as well as 6 mo old rats (Brushfield et al., 2006). Therefore, it is unlikely that the deficits on the contextual associative learning task in 24 mo old rats were due solely to difficulties in visually discriminating between the two contexts. Thus, it is suggested that the age-related deficits observed in the present task are the result of difficulty forming associations between a stimulus and a context.

The deficits observed in 24 mo old rats in associative memory for context may be due to age-related dysfunction in the medial temporal lobes and particularly in the hippocampus. The behavioral paradigm used in the present experiment was developed by Rajji and colleagues (2006). The paradigm was used to examine the effects of CA3 NR1 gene deletions in the hippocampus on learning novel paired associations between olfactory cues and their context. The results showed that NR1 gene deletions in less than 30% of the dorsal hippocampus CA3 subregion were sufficient to disrupt learning the contextual associations (Rajji et al., 2006). Therefore, the task appears to be quite sensitive to hippocampal dysfunction. As mentioned previously, a number of other studies also have shown that the hippocampus may be critical for processing contextual information (Anagnostaras et al., 1999; Anderson & Jeffery, 2003; Good & Honey, 1991; Smith & Mizumori, 2006; Wood et al., 2000). In addition, there are age-related functional changes in the hippocampus (Barnes, 1994; Driscoll et al., 2003; Mizumori, Lavoie, & Kalyani, 1996; Tanila, Sipila, Shapiro, & Eichenbaum, 1997) and aged rats demonstrate impairments on a variety of tasks that are considered to rely on the functional integrity of the hippocampus (Barnes, 1994; Dunnett et al., 1988; Gallagher et al., 1993; Rapp & Amaral, 1991). Therefore, age-related changes in the hippocampus may be responsible for the impairment found in associative memory for context in the present study.

The encoding of contextual information may be critical to the formation of episodic memories due to the need to form associations among stimuli, actions, and places that compose an event (for a review see Eichenbaum, 2004). The hippocampus may be particularly important for the formation of associations between a place and a stimulus (Gilbert & Kesner, 2003) and may be important for forming configural associations (Sutherland & Rudy, 1989). Therefore, age-related changes in associative memory for contextual information may contribute to the episodic memory deficits reported in aged humans and animals. Oler & Markus (2000) suggest that age-related deficits in episodic memory may result from an inability of the hippocampal network to respond to meaningful environmental changes. Numerous studies have shown age-related episodic memory impairments in healthy older humans (Craig, 1990; Rand-Giovannetti et al., 2006). Since hippocampal damage in humans has been shown to result in impaired episodic memory (Tulving & Markowitsch, 1998), episodic memory impairments observed in older adults likely result from age-related changes in the hippocampus. The results from the current study add to the literature suggesting that normal aging results in impairments in associative memory for contextual stimuli, which may contribute to age-related changes in episodic memory. The results also suggest that age-related changes in contextual memory can be observed using an appetitive-based paradigm. Therefore, the present paradigm may be useful for studying age-related contextual memory impairments in an animal model of aging.

In summary, the present study found age-related impairments in associative memory for context using an appetitive task. These results suggest that normal aging results in functional impairments in brain regions that support memory for associations between specific cues and their respective context. The findings may have important implications for understanding age-related changes in episodic memory.

Acknowledgements

This research was supported by NIH grant #AG026505 from NIA to Paul E. Gilbert. The authors thank Danielle Fellman for her assistance with data collection and Dr. Jeffrey Long for his technical advice.

References

- Anagnostaras SG, Maren S, Fanselow MS. Temporally graded retrograde amnesia of contextual fear after hippocampal damage in rats: within-subjects examination. *The Journal of Neuroscience* 1999;19:1106–1114. [PubMed: 9920672]
- Anderson MI, Jeffery KJ. Heterogeneous modulation of place cell firing by changes in context. *The Journal of Neuroscience* 2003;23:8827–8835. [PubMed: 14523083]
- Barnes CA. Normal aging: regionally specific changes in hippocampal synaptic transmission. *Trends in Neuroscience* 1994;17:13–18.
- Brushfield AM, McDonald C, Luu T, Moreland C, Callahan B, Penso L, Pavlik D, Robinson L, Wirkus J, Gilbert PE. The effects of normal aging on odor memory using an animal model. *Society for Neuroscience Abstracts*. 2006
- Bunsey M, Eichenbaum H. Conservation of hippocampal memory function in rats and humans. *Nature* 1996;379:255–257. [PubMed: 8538790]
- Craig FI. Changes in memory with normal aging: a functional view. *Advancements in Neurology* 1990;51:201–205.
- Dunnett SB, Evenden JL, Iversen SD. Delay-dependent short-term memory deficits in aged rats. *Psychopharmacology (Berl)* 1988;96:174–180. [PubMed: 3148143]
- Driscoll I, Hamilton DA, Petropoulos H, Yeo RA, Brooks WM, Baumgartner RN, Sutherland RJ. The aging hippocampus: cognitive, biochemical and structural findings. *Cerebral Cortex* 2003;13:1344–1351. [PubMed: 14615299]
- Eichenbaum H. Hippocampus: cognitive processes and neural representations that underlie declarative memory. *Neuron* 2004;44:109–120. [PubMed: 15450164]

- Gallagher M, Burwell R, Burchinal M. Severity of spatial learning impairment in aging: development of a learning index for performance in the Morris water maze. *Behavioral Neuroscience* 1993;107:618–626. [PubMed: 8397866]
- Gallagher M, Rapp PR. The use of animal models to study the effects of aging on cognition. *Annual Review of Psychology* 1997;48:339–370.
- Gilbert PE, Kesner RP. Localization of function within the dorsal hippocampus: the role of the CA3 subregion in paired-associate learning. *Behavioral Neuroscience* 2003;117:1385–1394. [PubMed: 14674856]
- Good M, Honey RC. Conditioning and contextual retrieval in hippocampal rats. *Behavioral Neuroscience* 1991;105:499–509. [PubMed: 1930720]
- Houston FP, Stevenson GD, McNaughton BL, Barnes CA. Effects of age on the generalization and incubation of memory in the F344 rat. *Learning and Memory* 1999;6:111–119. [PubMed: 10327236]
- Kraemer S, Apellbach R. Olfactory sensitivity, learning and cognition in young adult and aged male Wistar rats. *Physiological Behavior* 2004;81:435–442.
- Mizumori SJ, Lavoie AM, Kalyani A. Redistribution of spatial representation in the hippocampus of aged rats performing a spatial memory task. *Behavioral Neuroscience* 1996;110:1006–1016. [PubMed: 8919003]
- Moyer JR Jr, Brown TH. Impaired trace and contextual fear conditioning in aged rats. *Behavioral Neuroscience* 2006;120:612–624. [PubMed: 16768613]
- Mumby DG, Gaskin S, Glenn MJ, Schramek TE, Lehmann H. Hippocampal damage and exploratory preferences in rats: memory for objects, places, and contexts. *Learning and Memory* 2002;9:49–57. [PubMed: 11992015]
- Oler JA, Markus EJ. Age-related deficits on the radial maze and in fear conditioning: hippocampal processing and consolidation. *Hippocampus* 1998;8:402–415. [PubMed: 9744425]
- Oler JA, Markus EJ. Age-related deficits in the ability to encode contextual change: a place cell analysis. *Hippocampus* 2000;10:338–350. [PubMed: 10902903]
- Phillips RG, LeDoux JE. Differential contribution of amygdala and hippocampus to cued and contextual fear conditioning. *Behavioral Neuroscience* 1992;106:274–285. [PubMed: 1590953]
- Rajji T, Chapman D, Eichenbaum H, Greene R. The role of CA3 hippocampal NMDA receptors in paired associate learning. *The Journal of Neuroscience* 2006;26:908–915. [PubMed: 16421310]
- Rand-Giovannetti E, Chua EF, Driscoll AE, Schacter DL, Albert MS, Sperling RA. Hippocampal and neocortical activation during repetitive encoding in older persons. *Neurobiology of Aging* 2006;27:173–182. [PubMed: 16298252]
- Rapp PR, Amaral DG. Recognition memory deficits in a subpopulation of aged monkeys resemble the effects of medial temporal lobe damage. *Neurobiology of Aging* 1991;12:481–486.
- Rapp PR, Kansky MT, Eichenbaum H. Learning and memory for hierarchical relationships in the monkey: effects of aging. *Behavioral Neuroscience* 1996;110:887–897. [PubMed: 8918992]
- Rasmussen T, Schliemann T, Sorensen JC, Zimmer J, West MJ. Memory impaired aged rats: no loss of principal hippocampal and subicular neurons. *Neurobiology of Aging* 1996;17:143–147. [PubMed: 8786797]
- Schoenbaum G, Nugent S, Saddoris MP, Gallagher M. Teaching old rats new tricks: age-related impairments in olfactory reversal learning. *Neurobiology of Aging* 2002;23:555–564. [PubMed: 12009505]
- Smith DM, Mizumori SJ. Learning-related development of context-specific neuronal responses to places and events: the hippocampal role in context processing. *The Journal of Neuroscience* 2006;26:3154–3163. [PubMed: 16554466]
- Sutherland RJ, Rudy JW. Configural association theory: the role of the hippocampal formation in learning, memory and amnesia. *Psychobiology* 1989;17:129–144.
- Tanila H, Sipila P, Shapiro M, Eichenbaum H. Brain aging: impaired coding of novel environmental cues. *The Journal of Neuroscience* 1997;17(13):5167–5174. [PubMed: 9185554]
- Tulving E, Markowitsch HJ. Episodic and declarative memory: role of the hippocampus. *Hippocampus* 1998;8:198–204. [PubMed: 9662134]

Wood ER, Dudchenko PA, Robitsek RJ, Eichenbaum H. Hippocampal neurons encode information about different types of memory episodes occurring in the same location. *Neuron* 2000;27:623–633. [PubMed: 11055443]

Figure 1.
Photographs of the two contexts used in the contextual associative learning task.

CONTEXTUAL MEMORY

Figure 2.

The mean ($\pm$ SE) percent correct performance of 6 mo and 24 mo old rats on the contextual associative learning task as a function of days (1–8).