reserve this type of therapy for cases of undulant fever that have resisted other forms of treatment.

We should like to express our thanks to Messrs. May and Baker, Ltd., for generously supplying the amount of M & B 693 required for our experiments.

REFERENCES

- Antopol, W., and Robinson, H. (1939). *Proc. Soc. exp. Biol.*, N.Y., **40**, 428.
 Chinn, B. D. (1939). *J. infect. Dis.*, **64**, 78.
 Colebrook, L. C. (1939). *Lancet*, **2**, 158.
 Gross, P., and Cooper, F. B. (1939). *British Medical Journal*, **1**, 1058.
 Marshall, E. K., and Long, P. H. (1939). *J. Amer. med. Ass.*, **112**, 1671.
 Menefee, E. E., and Poston, M. A. (1939). *J. Bact.*, **37**, 269.
 Wien, R. (1938). *Quart. J. Pharm. and Pharmacol.*, **11**, 217.
 Wilson, G. S., and Maier, I. (1939). *British Medical Journal*, **1**, 8.

PLATELET-REDUCING EXTRACTS OF THE SPLEEN

BY

FELICITY C. G. HOBSON, B.A.

AND

L. J. WITTS, M.D., F.R.C.P.

(From the Radcliffe Infirmary, Oxford)

When Troland and Lee (1938a, 1938b) reported the preparation from spleens removed from patients with thrombocytopenic purpura of an extract which reproduced the condition in animals hope was felt that we might be approaching an understanding of this very puzzling disease. It subsequently appeared that Torrioli and Puddu (1938) had been working on similar lines since 1933, and they issued a warning that the results were by no means so clear-cut or so specific as Troland and Lee had claimed. More recently Pohle and Meyer (1939) were unable to duplicate Troland and Lee's results in three cases of thrombocytopenic purpura. The rarity of the disease must be our excuse for reporting results in a single case.

A married woman aged 48 presented the classical features of thrombocytopenic purpura. She had had the disease for a year before coming to us, and we kept her under observation for a further three months, until menorrhagia made splenectomy essential. Her spleen showed the negative picture described in thrombocytopenic purpura, and her platelets, which we had never found above 45,000 per c.mm. before operation, rapidly rose to over 200,000. She promptly recovered from all her symptoms.

The spleen, which weighed 250 grammes, was extracted with acetone for seventeen days and, after removing the acetone and making the extract up to 100 c.cm. with distilled water, 5 c.cm. and 20 c.cm. respectively were injected intravenously into two rabbits. Little effect was produced by 5 c.cm., but after 20 c.cm. there was a sharp fall in the number of platelets. We then injected another 45 c.cm. into the first rabbit, and the platelets fell in thirty hours from 525,000 to 109,000 per c.mm.; they had returned to normal by the fifth day. This was, of course, a relatively enormous dose, corresponding to about 112 grammes of fresh spleen. No abnormal symptoms were noted.

Our first control was an acetone extract prepared in the same way from a spleen removed post mortem from a man aged 42 who died of a fractured skull. After intravenous injection into Rabbit No. 3 of 40 c.cm., corresponding to 44 grammes of fresh spleen, the platelets fell

Acetone Extract of Thrombocytopenic Spleen

Hours	Platelets per c.mm.	
	Rabbit No. 1 (Wt. 2.62 kg.)	Rabbit No. 2. (Wt 2.71 kg.)
Before Injection		
66	424,000	553,000
48	462,000	525,000
40	398,000	486,000
24	476,000	501,000
1	484,000	576,000
After Injection	Inject 5 c.cm. I.V.	Inject 20 c.cm. I.V.
1	472,000	368,000
4	408,000	405,000
22	406,000	442,000
24	476,000	471,000
30	495,000	482,000
46	549,000	572,000
70	526,000	
94	525,000	
After Injection	Inject 45 c.cm. I.V.	
1	329,000	
3	321,000	
6	386,000	
22	167,000	
24	130,000	
30	109,000	
48	164,000	
54	214,000	
70	270,000	
72	400,000	
77	391,000	
96	390,000	
120	568,000	

Acetone Extract of Normal Spleen (Rabbit No. 3, Weight, 2.52 kg.)

Hours	Platelets per c.mm.	Hours	Platelets per c.mm.
Before Injection		After Injection	Inject 40 c.cm. I.V.
24	400,000	2	306,000
18	447,000	5	385,000
2	454,000	22	460,000

Ringer's Extract of Normal Spleen

Hours	Platelets per c.mm.	
	Rabbit No. 4. (Wt. 2.55 kg.)	Rabbit No. 5. (Wt. 2.34 kg.)
Before Injection		
42	230,000	422,000
25	254,000	421,000
19	283,000	433,000
1	219,000	351,000
After Injection	Inject 5 c.cm. I.V.	Inject 20 c.cm. I.V.
3	211,000	300,000
7	217,000	246,000
22	231,000	329,000
24	187,000	267,000
30	270,000	315,000
48	256,000	400,000

in two hours from 454,000 to 306,000 per c.mm., but were back to normal within twenty-four hours. As the choice of acetone by Troland and Lee appeared to be purely arbitrary we then tried the effect of an extract made by emulsification of normal spleen in Ringer's solution in the cold for twenty-four hours and subsequent centrifuging. This spleen was obtained post mortem from a child aged 3 who died from cerebellar decompression. Of this extract 5 c.cm. had no obvious effect, but 20 c.cm., corresponding to 10 grammes of fresh spleen, reduced the platelets appreciably.

Summary

The acetone extract of a thrombocytopenic spleen reduced the platelets of a rabbit more powerfully than an acetone extract of a normal spleen. A Ringer extract of normal spleen was more effective in reducing platelets than the acetone extract of normal spleen, and it appears desirable to determine the optimum conditions for extracting platelet-lowering substance from normal spleen before comparing the conditions in health and disease.

REFERENCES

- Pohle, F. J., and Meyer, O. O. (1939). *J. clin. Invest.*, **18**, 537.
 Torrioli, M., and Puddu, V. (1938). *J. Amer. med. Ass.*, **111**, 1455.
 Troland, C. E., and Lee, F. C. (1938a). *Johns Hopk. Hosp. Bull.*, **62**, 85.
 ——— (1938b). *J. Amer. med. Ass.*, **111**, 221.

TREATMENT OF INFECTED BURNS

BY

**B. C. MURLESS, M.B., B.Ch., F.R.C.S.Ed.
 M.R.C.O.G., Surgeon Lieutenant R.N.V.R.**

Registrar and Pathologist, the Hospital for Women, Soho Square; Gynaecological Registrar, Royal Waterloo Hospital

The literature on the treatment of burns is legion. A great number of publications followed the introduction of the use of tannic acid by E. C. Davidson of the Henry Ford Hospital, Detroit, in 1925, which has revolutionized this branch of surgery. Many writers since have published modifications of the original technique described, with reports showing excellent results. It is therefore with some trepidation that I venture to add to a subject so extensively investigated. The tremendous increase of burns in war-time surgery may, however, render this short article of interest now, and the somewhat rare opportunity of treating and watching the progress of eighteen cases of severe burns from one of His Majesty's ships seems worth recording.

The Use of Tannic Acid

The functions of tannic acid are twofold. First, in the early stages it prevents the absorption of toxins by combining with proteins and their metabolites to form a coagulum; and, secondly, the coagulum formed exercises a protective action. Suffering and shock are prevented by the protection of exposed nerve endings, but equally important is the effect of the coagulum in rendering the surface free from outside disturbance and guiding the growth of the new epithelium.

For practical purposes a burn may be considered like any flesh wound, healing proceeding most rapidly in the absence of bacterial infection. Such conditions are of course ideal for tannic acid treatment, the burnt area being rendered aseptic by cleaning before the tannic acid is applied. In war time the difficulties of thoroughly cleansing burns are many, and delay must often occur.

Severe shock may contraindicate anaesthesia or any active treatment; transport difficulties are bound to delay the arrival of cases at the theatre. Consequently, though they may have received preliminary treatment, many burns met with under war-time conditions must be of the septic variety, and it is with such cases that this article is concerned.

Tannic Acid in Septic Cases

Burns which have been left for forty-eight hours or more may be considered septic, as no amount of treatment with ether or spirit will render them surgically clean. Difference of opinion exists as to the use of tannic acid in septic cases. Some surgeons prefer picric acid dressings unless the burn has been cleaned within forty-eight hours. Hamilton Bailey (1938) states that tannic acid treatment may be instituted as late as seventy-two hours. The cases reviewed here may help to support the contention that there is no time limit after which tannic acid treatment should be abandoned, and that good results may be obtained by its use where sepsis is fully established.

When applied to a septic surface tannic acid will form a coagulum more readily than on a surgically clean area. A crust soon dries which, if it stays in place and can be kept hard, will supply the rest and protection essential for the healing of the inflamed tissues beneath. Such treatment may be compared with that adopted by Winnett Orr for open fractures and by Dr. J. Trueta for infected wounds in Spain. The essential advantages of this method are that the surface remains undisturbed, the covering allowing epithelium to grow in from the edges, at the same time preventing the formation of fibrous tissue and piled-up granulations, which are nearly always seen when infected burns have been treated by constantly changed dressings. The speed of epithelization with such treatment is infinitely greater than when the surface is disturbed daily, and the saving of pain and discomfort to the patient beyond estimation.

Pus Formation

Pus will usually collect beneath the crust and may form in patches, or almost the whole crust may be slightly raised and fluctuation may be present. The usual teaching of removal of or incision into the coagulum under such circumstances was not adopted in these cases. If an expectant attitude is taken, within twelve hours the pus will always find its way out at some point. It may come through cracks in the coagulum, especially at joint flexures, or may ooze out at the edges. After discharging for a short time suppuration will cease, and any part of the crust which has become softened may be resprayed with tannic acid and dried. Not unnaturally the smell during the suppurative phase is somewhat unpleasant, but it is not nearly so bad as that encountered in the closed treatment of infected fractures.

Fever and Pain

Nearly all cases register a rise in temperature during this treatment. It may reach as high as 104°, but this should not cause undue anxiety, and should not be considered an indication for removal of the coagulum. In most cases the fever subsides in from two to three days.

The absence of pain during tannic acid treatment is well known to be remarkable, and the same fact applies when it is used for septic burns. Following the first application and drying little pain is experienced. A feeling of tenseness and heat sometimes exists for a few hours during the suppurative phase, before the pus has