SURGICAL PROCEDURES IN GENERAL PRACTICE

This is one of a series of articles contributed by invitation

THE AFTER-TREATMENT OF FRACTURES

BY

H. OSMOND CLARKE, F.R.C.S.

In the treatment of fractures there are three main objectives: (1) to secure bony union; (2) to restore full movement in those joints which have been temporarily immobilized; and (3) to restore the maximum power in the muscles which move those joints. This communication is concerned only with the latter two aims. All three must, however, be visualized from the earliest stage in the treatment of every fracture. Thus it is possible to discuss the preservation and restoration of function—which is after all the prime consideration in the treatment of all injuries and most diseases—under two main headings: (a) during the period of fixation; (b) after the splints have been discarded.

Treatment during Period of Fixation

Adequate fixation of a fracture of a long bone demands the immobilization of the joint above and the joint below the site of fracture. In a large number of fractures this immobilization is best provided by plaster-of-Paris applied directly on the skin without padding. This affords such security that it is possible for the remaining joints of the extremity to be used freely without risk of disturbing the fragments—for example, patients with fractures of the forearm bones in which the elbow and wrist are immobilized can use the fingers and shoulder freely. And it is of the greatest importance that free use of the joints not splinted should be insisted on. Not only does it entail a full range of movement in these joints from the very beginning, but it means that the muscles covered by plaster-of-Paris are being contracted constantly even though they are unable to move the splinted joints. In this way venous stagnation and muscle atrophy are reduced to a minimum, and thus wasting and oedema, so often seen after the abandonment of splints in the days when it was the custom to immobilize the whole extremity, are nowadays rarely seen. Tendons no longer become bound down by adhesions in their grooves, and one of the most gratifying features is the amount of movement possible in the joints immediately after the fixation apparatus is removed. By this present-day method of so-called "functional use" the active power and mobility of an injured extremity at the time when the fracture has united far exceed what could be obtained by the older "extremity immobilization" methods. So much so that the early removal, while the fracture is still not soundly united, of retentive apparatus for massage and passive or active movements has been for the most part discarded.

There are, however, certain fractures in which early movement and physical treatment are justified. Broadly speaking, these are impacted fractures in the region of the surgical neck of the humerus, in which very effective fixation is supplied by the surrounding muscular attachments and in which adhesions of torn folds of the highly voluminous capsule and synovial membrane of the shoulder-joint are very prone to occur; and some other fractures in old people—for example, "bumper" fractures of the external tuberosity of the tibia.

Treatment after Discarding of Splints

The fracture has now united and may be sufficiently consolidated to withstand the cross and shearing strains of muscular action alone, as in the upper extremity, but not the strain of weight-bearing, as in the lower extremity. Consequently fractures of the femur or tibia may require the support of a calliper or walking steel for some months after the more complete fixation has been discarded. Indeed, some mild support for all fractures after discarding the retentive apparatus is greatly appreciated by all but very young patients—for example, a few turns of strapping around the wrist when a Colles's fracture has united gives the patient a sense of security and encourages him to start movements of his wrist more readily.

Restoration of joint movement and of muscle power proceeds concomitantly. Children and young adults can usually mobilize their joints without any special help other than appropriate instruction. Older people are often apprehensive and tend to "nurse" an injured joint after removal of the splint. Physiotherapy is most helpful for the latter group of patients as a supplement to their own efforts. The following regime, which can be modified to suit individual cases, has proved valuable.

1. Thorough "baking" of the affected joint by the application of heat in any of the well-recognized forms (radiant heat, infra-red, or a rubber hot-water bottle). The resulting hyperaemia constitutes an excellent local analgesic.

2. Massage and faradic stimulation of the muscles which move those joints.

3. Encouragement in voluntary movements. It must be emphasized that the constant active efforts of the patient himself are by far and away the most important factor in restoring muscle power and joint movement. There is, unfortunately, a mistaken impression among the lay population, and indeed among many doctors, that complete restoration of function follows automatically on the application of the various forms of physiotherapy. Consequently the patient does not feel it imperative, nor is he urged, to perform active movements at regular intervals. A definite routine of muscular contractions and relaxations which entails movement of the joint or joints in all directions should always be instituted for a period of five or ten minutes every hour during the day. At first the patient, especially if elderly, may require the sympathetic but firm encouragement of a doctor or masseuse, and his position should be arranged so that he is not called upon immediately to carry out movements against the resistance of gravity. It is better, too, for each patient in the early stages of his rehabilitation to receive individual treatment. Once the joints begin to mobilize and confidence has been restored, class exercises can be started and a spirit of competition encouraged. As muscle power and bulk increase and the joints approach the limits of movement more vigorous exercises will help to complete the cure, and will be valuable in assessing the amount of stress and strain to which the fractured region can be subjected.

Passive Movements

Following prolonged immobilization of a joint there is always a certain amount of stiffness when the splint is removed, especially if there has been a fracture in the neighbourhood of the joint. This limitation of movement is due to "adhesions," a word commonly used and rarely understood. Adhesions are simply bands or masses of scar tissue formed by the organization of blood clot and the sero-fibrinous exudate which follow on an injury. In this way folds or plications of the joint capsule

become adherent, and painful limitation of movement results. In the vast majority of fractures such adhesions are most effectively stretched gradually by the patient's own active efforts on removal of the splint. It cannot be emphasized too often that passive movements carried out frequently and enthusiastically are most prejudicial to the restoration of joint movement. They cause, by repeatedly tearing, as opposed to stretching, adhesions, further exudate and more adhesions, and may result in the formation of plaques of new bone in the capsule or surrounding muscles. As the late Sir Robert Jones repeatedly stressed, they have no place in the early stages of joint mobilization after injury, as they always result in increasing stiffness.

Manipulation

Occasionally, after a prolonged trial of active mobilization, a manipulation under anaesthesia may be advisable to overcome residual stiffness in a joint situated close to a fracture, or for the cure of pain persisting after an apparently full range of movement has been achieved. In the absence of arthritis this pain is often due to adhesions, and on careful examination will be found to occur when the very extremes of a certain range are forced. The success of a manipulation will depend on the nature of the adhesions present (whether localized or diffuse), on the timing and technique of the manipulation, and on the character of the after-treatment. The appropriate time at which a manipulation should be carried out is most difficult to decide, and much experience and careful judgment are required. It can be stated emphatically that no manipulation should be attempted: (a) In the early stages after a fracture has united when the neighbouring joints are still irritable and manifest their irritability by muscle spasm if movements, active or passive, are attempted beyond the limit of pain. Manipulation at this stage may cause re-fracture and will increase capsular scarring and damage joint cartilage, and may produce nerve injury and myositis ossificans. It is still unfortunately too common to see patients whose elbows, within two or three weeks following a supracondylar fracture or dislocation, have been "pump-handled" at two or three day intervals under an anaesthetic and are permanently stiff. (b) If movement is increasing, however slowly, as the result of the patient's own active effort. Sometimes impatience on the part of the patient and the doctor will lead to an unnecessary and prejudicial manipulation. (c) In children. It has already been stressed that young people are capable of restoring full movement by their own efforts alone. The worst tragedies of injudicious manipulations are exhibited in fractures around the elbow-joint in children. (d) In the presence of vascular or nervous complications or arthritis.

There is no magic in the technique of manipulation. It consists essentially in putting a joint through its fullest range of movement in all directions. But this cannot be done safely by any doctor. Certain safeguards of technique, of touch—a certain "sense"—are developed with experience which enable the surgeon to avoid using too little or too much force, to avoid breaking bones and dislocating joints, to avoid doing too much or too little at one sitting. It is often better and will cause less reaction to stretch adhesions by a series of careful incomplete manipulations at intervals of six weeks rather than to perform one very thorough manipulation.

Finally, the good effects of a manipulation will be jeopardized if it is followed by injudicious after-treatment—that is, repeated forcible passive stretchings.

Rehabilitation

The majority of patients—children, housewives, and sedentary workers—treated efficiently require no special rehabilitation. There is, however, a small minority of patients, particularly those suffering from industrial injuries, for whom the gap between the stresses of such remedial activity as a hospital massage department can provide and the stresses of full heavy work should be bridged by a final period of re-educational treatment. This would aim at subjecting the patient through graduated work to the full stresses of his original occupation while under constant medical supervision. This constitutes the argument for the creation of rehabilitation centres in suitable areas throughout the country. In the final report of the Home Office Interdepartmental Committee on the Rehabilitation of Persons injured by Accidents (1939) it is appreciated that "reconditioning" is necessary for a small minority of patients, and various schemes are suggested.

The L.M.S. Railway rehabilitation hospital at Crewe receives a considerable number of patients with old injuries of all kinds who, after long periods of disability and long periods of treatment in physiotherapeutic departments of general hospitals, remain unfit for work and seemingly incapable of improvement. Of 165 consecutive patients of this type, 115 were returned to their former employment after an average period of seventeen days of rehabilitating treatment in the Crewe hospital. An effort to incorporate rehabilitation as a routine in organized fracture treatment is to be seen in the new fracture clinic founded by the Seamen's Hospital Society at the Albert Dock Hospital, London. This clinic is a highly specialized one, for it deals only with industrial injuries in sailors.

So far as possible—and this is the practice in some progressive massage departments—exercises given to the patient should be of an occupational character and correspond closely to those which he is constantly carrying out in the process of his work. Another excellent method of rehabilitation is through the medium of light work—a method employed with the very greatest success by Dr. O'Neill Sherman of the Carnegie Illinois Steel Trust. He is fortunate, however, in being able to insist, as soon as he considers a man fit for any form of work, that suitable work is provided. In this country it is regrettable that similar facilities, though occasionally available, are few. The ability of an injured workman to do "light work" provides more often than not an excuse for depriving him of a substantial amount of compensation without the provision of suitable employment. As the Workmen's Compensation Act stands at the moment it is often difficult to prevent the development of an anxiety neurosis, with consequent persistent disability in the injured workman. It is to be hoped that suggestions for the alteration of this aspect of the Act will be made by the Royal Commission on Workmen's Compensation now considering the subject. If altered effectively rehabilitation would no doubt become an easier problem.

Conclusions

The process of preserving and restoring function begins with the initial treatment of the fracture.

Active movements by the patient are the most important factor in restoring joint mobility and muscle power. Physiotherapy is helpful for adults but is unnecessary for children.

Passive movements are likely to do more harm than good. Manipulation of a joint which has resisted the patient's active efforts is occasionally justifiable. When and how it should be done require experienced judgment.