tions, just as we are looking for syphilis and rheumatic fever.

A prominent feature of gonococcal infection is its obstinate persistence—that is, the tendency to formation of chronic foci. This fact seems to reflect itself to some extent in the course of the complement-fixation reaction. Investigation into this reaction is now being made in co-operation with Dr. Peter Krag of the State Serum Institute. One of the results may be mentioned in this connexion—namely, the surprisingly high incidence of positive reactions found among patients without apparent signs of recognized sequels to gonococcal infection. The Wassermann reaction has aided in the evaluation of many a syphilitic manifestation that would scarcely have been diagnosed as such without this serologic evidence. Perhaps in future the gonococcal complement-fixation reaction will help to reveal that chronic gonococcal infection plays a part in one or more diseases not hitherto attributed to this cause. Certain heart diseases may be among them.

Summary of Conclusions

Apart from the myocardial involvement in the bacterial endocarditis of gonococcal septicaemia, myocarditis is seldom established as a complication in gonococcal infection. In view of this it seems surprising that a series of six instances of what is regarded as gonorrhoeal myocarditis was encountered in the course of one year in an ordinary medical ward. It seems probable that this condition often escapes detection, since suggestive symptoms may be absent.

The diagnosis was made on the ground that all six cases had changes in the electrocardiogram occurring in the course of acute or chronic complicated gonococcal infection. In some instances the myocarditis was slight and transitory, as may be the case in many simple infections—for example, pneumonia. However, in two cases incomplete bundle-branch block developed and persisted, and in a third case the anginal pain still troubled the patient considerably when he was re-examined a year later. Such a course suggests the possibility that some of these patients will return to hospital sooner or later with chronic heart disease. It is possible, also, that some of the chronic “degenerative” heart diseases which we see to-day arise from gonococcal infection.

The gonococcal complement-fixation reaction has been instrumental in pointing out how great a proportion of so-called rheumatic arthritis cases were really gonorrhoeal. It may be worth while to search the anamnesis of heart patients for gonorrhoea as we search it for syphilis and rheumatic fever, and the complement-fixation reaction may reveal that some instances of heart lesion attributed to rheumatic infection ought to be regarded as following on persistent gonococcal infection.

REFERENCES

- Councilman, W. T. (1893). *Amer. J. med. Sci.*, **106**, 277.
Dittrich, O. (1930). *Jadassohn's Handb. Haut. u. Geschl.*, **20**, pt. 2, J. Springer, Berlin.
Gillespie, J. O., and Thompson, R. M. (1937). *Milit. Surg.*, **80**, 418.
Guldberg, G. (1935). *Norsk Mag. Laegevidensk.*, **96**, 576.
Larsen, K. H., and Warburg, E. J. (1937). *Hospitalstidende*, **80**, 276.
Neumann, H., and Bingenheimer, E. (1937). *Derm. Wschr.*, **105**, 1099.
Rowlands, R. A., and Simpson, S. L. (1937). *Brit. J. ven. Dis.*, **13**, 215.
Williams, R. H. (1938). *Arch. intern. Med.*, **61**, 26.

Bronchography.—The authors describe a method by which radiographs may be obtained showing the lipiodol-filled bronchi standing out in relief. Some excellent examples are included in the article.—Bronchography in Relief. M. R. Castex, E. Alvarez, and E. S. Mazzei.—*Arch. méd.-chir. Appar. resp.*, No. 6 (1938), **13**, 441.

THE PREVENTION OF ANXIETY

BY

R. G. GORDON, M.D., F.R.C.P.Ed.

Physician, Royal United Hospital, Bath

A young happily married woman aged 30 with two healthy children is the victim of intense anxiety if she goes out alone. While she is working about the house she is all right. If she goes out in the company of certain people she is practically all right. The best “protection” in this respect is her husband, next her sister, then her doctor—the general practitioner who looks after the family, but any doctor is “some use.” Other people help but do not abolish the anxiety.

Physically she is healthy and any organic disease can be excluded, but in addition to the anxiety she complains of uncertain sleep, dull headache, poor memory, and inability to concentrate. Asked to describe the anxiety, she states that it seems that something dreadful is going to happen, probably that she is going to die. Her husband would prevent it happening and doctors would see that it didn't. The anxiety was not always a fear of death; it used to be a fear of going insane, but that has now changed to a fear of death. The anxiety is obsessional in nature—that is, intellectually she realizes that it is irrational, but emotionally she is quite powerless to overcome the feeling.

When asked what to her is the significance of death, she says she is not sure, but states vaguely: being separated from those she knows and loves, being “shut up in a box.” She hopes that she would be liberated and reunited to her loved ones after death, but is very uncertain and muddled about this. Questioned about going insane, she replies with a heated diatribe against the iniquity of the system by which doctors can shut people up for no adequate reason. Pressed on this point she said, “Of course, good doctors would not do this, but it is scandalous how easily it could be done.”

The foundation for this attitude seemed to be that an aunt of her husband's had been “put in an asylum when she was really quite sane.” This was interesting, for this aunt was clearly a psychopathic individual, was an incorrigible thief, and, by what possibly was somewhat of a “wangle,” had been represented in court as irresponsible for her actions and confined in a mental hospital instead of in a prison, a procedure which apparently was regarded as a lesser evil by most of her relatives but as a scandalous crime by our patient.

The real basis of the present anxiety seemed, therefore, to be a fear of going out of her mind and of being shut up away from her friends. The fear had later been transferred to death as a lesser evil, since if she was dead she herself would not know about it. A light is thrown on this by her previous medical history. She had been quite healthy, apart from one or two exanthemata in childhood, till a very short time before her marriage, when she had an attack of appendicitis which necessitated operation and the postponement of her wedding. This was attended by no anxiety whatever, only exasperation at the upset of arrangements. A severe attack of influenza after her marriage left her similarly emotionally unscathed. Her next illness six months later, however, had very different results. At the time it seemed trivial: she knocked her head on a low door and sustained slight concussion. She was not rested enough and began to have headache, loss of memory, and poor concentration. These symptoms,

with gradually increasing obsessive anxiety, had persisted for nearly seven years. It was quite clear that the original concussion was not so severe as to cause permanent organic changes; why, then, did she remain persistently ill? Obviously this illness alarmed her more than any other had done, and what had particularly struck her about it was the failure of memory concentration and general mental capacity.

Further investigation elicited that she thought this might mean insanity; so the fear of mental illness was partly explained, and this was later intensified by the fate of her husband's aunt, but, as is usual in such cases, there were other more important determinants further back in her history. Her sister was in her eyes much more attractive in looks and in every other way. She was gay, a good mixer, and made friends with the greatest ease. The patient was shy, diffident, and had few friends, but she was clever. Her parents always expected her to do well at school work, and she did: intelligence was her one real asset. Actually she became engaged before her sister did, and, although she is an attractive, nice-looking girl, she was sure that it was her mind that had drawn her husband—the one strong emotional attachment of her life—to her side. So if she were to “lose her mind” as the result of this accident all was lost.

This was not all, however; she was emotionally insecure and she feared violent accidents. It may be noted that her husband was far the best person to relieve her of her anxiety if she went out, and he was her one deep emotional episode, as has been said. As she expressed it, her parents had brought up their children to respect them rather than to love them. Her father had been an austere and rigid man, and had died during her childhood without much psychic trauma on her part, since he had never been close to her. Her mother had died of cancer in the patient's early adult life after a painful illness nursed by the patient, whose sister was away from home. The mother's death had been rather a relief, and the patient had maintained herself alone in a flat for two years before her marriage. She had hardly had time to feel seriously lonely, however, since her husband had appeared on the scene not long after her mother's death. She had never known what it was to be enfolded and protected by parental love, and therefore was not well safeguarded against the buffets of fate, but, if these took the form of violent injury, an early experience had intensified the fear.

By far the most vivid memory of childhood was that of air raids and of being wakened in the night, taken downstairs, and told to stay under the dining-room table with her sister while her parents went to the front door to see what was happening. She said that her sister, who was three years younger than herself, did not seem to mind; but she herself was terrified. That it made a deep impression was evident from the fact that it seemed to her that this happened over and over again, whereas from the part of the country where she lived actual raids could not have occurred more than twice. Her parents, however, were very agitated about raids and were always frightened lest they should occur, and their going out to see what was happening, leaving the children to shiver under the table, was not an expression of courage or bravado but of an agitation too great to allow them to stay still.

Such, then, was the background of this case of pathological anxiety—not an unusual or a remarkable case, but one which may serve as a text for a discussion on the genesis of this condition, which may be all too common if the war develops a more active phase.

The Physiology of Anxiety

Anxiety is the emotion engendered by insecurity. It differs from fear in that it is not the response to a sudden actual danger but to a continually present potential danger. The stimulus comes, therefore, almost entirely from fantasy, from the thought of the subject as to what might happen and not directly from what does happen in the environment.

If we are to understand the physiology of anxiety we must first understand the physiology of such an experience as fear. In this case the stimulus arrives through the sensory receptors, let us say the sight of a charging lion. The image of a lion is formulated first in the occipital cortex, but as other association paths are quickly traversed by the impulse which has been set going through the cortex, meaning is attached, danger is recognized, and the well-facilitated pattern of escape is set in action with the accompanying emotion of fear and the motor phenomenon of flight. The latter is activated by means of the motor cortex, but the impulses pass not only forward in the brain to the effector areas responsible for action of voluntary muscles, but also downwards to the thalamus and hypothalamus, where autonomic changes are initiated and endocrine secretion is procured through the controlling medium of the pituitary. Such changes “set” the body for violent muscular effort and provide the necessary chemical changes to maintain this. The resultant sensory pattern, returned, as it were, to the cortex and so to consciousness by these autonomic and chemical changes, constitutes what we experience as emotion—in this case the emotion of fear.

We know that sympathetic activity and excess of adrenaline are associated with fear; the emotional pattern might, therefore, be described in terms of an equation involving sympathetic +, vagal −, and endocrine (adrenaline +) values. As yet, however, we cannot get very far in writing such an equation, since the equation for anger must also be sympathetic +, vagal −, and endocrine (adrenaline +); but fear is quite distinct and different from anger. Still, for fear the process is: environmental danger → cortical perception → thalamic and hypothalamic activity → autonomic and endocrine pattern set in motion → return cortical perception of this, which is the conscious experience of fear. This will persist just so long as environmental danger persists, but when it stops fear stops. Motor activity by removing the subject from danger will hasten the cessation of fear; but if motor activity is prevented—for example, by tying up the subject—the impulse tends to flow more and more through hypothalamic paths with more and more derangement of autonomic and endocrine activity, which may become so intense that functions of the body, including that of the cerebral cortex, are all much upset, with the result that the subject actually loses consciousness. The reactions to environmental danger associated with fear are fairly constant in their form, since the pattern is inherently facilitated, being instinctive and therefore innate. Experience, suggestions, and education will determine which objects or situations act as most effective stimuli, but the pattern “flight-fear” is always there ready to be set going.

The infant at birth can experience fear but probably not anxiety; anxiety requires thought and reflection, at least in their primitive form. The infant may experience anxiety if he thinks the breast is going to be denied him. We do not experience anxiety when we are aware that the lion is there, but only when we think he might be there. The anxiety pattern, then, is not innate but acquired, and therefore is not so set and so facilitated.

Moreover, there is very little if any motor-cortex-voluntary-muscle component, and so the autonomic endocrine activity may be very severely disturbed. It is noticeable, however, that it is not nearly so definitely sympatheticotonic. Vagotonic symptoms (intestinal spasm, diarrhoea, etc.) are often met with, while thyroid overactivity is much more characteristic than adrenaline overactivity. The anxiety equation will read, therefore, sympathetic +, vagal +, endocrine (thyroid +), though again we cannot specify this with any exactitude. Although the anxiety pattern is not innate like the fear pattern, we are all capable of experiencing anxiety, and it may be that in some of us there is an inherent disposition which makes us more liable to anxiety than others; but almost certainly the tendency to anxiety depends much more on early experience. As has been said, anxiety is the emotion that is associated with a feeling of insecurity, and for the infant and young child security depends on the enjoyment of parental and perhaps especially maternal love. If the infant is deprived of parental love he will tend to be anxious, and there is no doubt that the broken or quarrelsome home makes for anxiety in the child. If perfect love casteth out fear, most certainly imperfect love generates anxiety. It is probably true to say that the anxious parent is incapable of perfect love, and undoubtedly his child is most suggestible in respect of anxiety. Such a symptom as enuresis, which in the majority of cases is psychogenic, illustrates this; for the more anxious and fussed the mother the more anxious the child and the more frequent the wet bed.

Basis of Pathological Anxiety

In our case the basis of the subsequent pathological anxiety was laid by the fact that the patient did not enjoy perfect parental love, for her parents wished to be respected above everything. The anxiety was intensified enormously by the suggestion given by the parents' anxiety over air raids, perhaps more than by the actual experience of them, for, as has been said, they were not actually frequent but very rare. It is significant in this case that the more attractive younger sister was the parents' favourite and has not developed a pathological anxiety, so there may also have been the factor that the elder sister's nose was put out of joint by the arrival of the younger, though this did not come out during treatment.

The basis of all neurotic anxiety, then, is deprivation of parental love because the parent (*a*) is not capable of unselfish love for his or her child, or (*b*) is not there to give it, or (*c*) has displaced it partially or wholly on to another child. This may well be intensified by anxiety on the part of the parents, and may do something to determine the nature of the anxiety and its objects. Violent accidents as a cause of anxiety were so determined in our case.

More important, however, in shaping the psychic pattern of the anxiety are the patient's subsequent experience and development. Insecurity will be more readily felt if one single attribute is regarded by the subject as all-important, whether because of its own inherent significance or because the patient regards it as all he or she has got. In our case it was intelligence, but more often it is some physical attribute which brings power to the patient or which he thinks brings him power over his friends or relatives. This attitude may be in reality or fantasy, may be enjoyed or only wished for. The boy may wish for physical strength sufficient to overcome his oppressive father, the daughter may wish for beauty to

outshine her mother, and each of them may develop anxiety about anything which threatens to deprive him or her of these desirable attributes. In analysis, given such a situation, it is not difficult to discover early memories related to an Oedipus situation.

Again, where an insecure girl wishes for power over her apparently more secure brothers she may easily feel that she has been deprived of something which they have, and if the power she wishes is sexual then the castration complex situation can easily be discovered. But these are more likely to be "retrojections" of ideas from present experiences than real past experiences. In the boy, ideas of physical or sexual insecurity may easily be associated with anxiety regarding possible impotence. Such anxieties may be attached to masturbation or venereal disease, which may be supposed to determine the impotence.

In other cases the insecurity is social or economic rather than physical or sexual, and the physical pattern is so shaped. Given the tendency to anxiety acquired in early childhood and the general object of anxiety acquired as a rule in adolescence, the actual onset of the pathological anxiety is precipitated by some incident which threatens the cherished attribute of the patient. In our case this was the concussion which threatened the integrity of her mind.

Establishment of the Pathological Condition

Once the anxiety is started its persistence will be determined by the pre-existing sense of security or its absence, and what we may call pathological anxiety will not be established unless the threat to security is continuous and repeated or unless there is a serious predetermined sense of insecurity, initiated as a rule in early childhood.

These two latter factors are generally in inverse proportion—the greater the predetermined sense of insecurity the more trivial need be the precipitating cause; but once the anxiety is established the patient is apt to perpetuate it because any untoward symptom or external event is interpreted as fresh evidence of potential danger. Since the anxiety itself is a source of autonomic dysfunction such symptoms are all too common, and such autonomic dysfunctions are hard to describe, for even the patient's doctor often finds it difficult to explain them to the patient and to reassure him as to their significance or insignificance. This means that these symptoms and feelings are mysterious and so tend to intensify the anxiety.

This was well illustrated by a patient who was recovering from phthisis and developed a pathological anxiety quite unconnected with his lung disease. This was accompanied by marked psychically determined palpitation, which terrified him exceedingly and intensified his conviction that he was on the point of death. As he expressed it, he was never bothered about the tuberculosis—he always knew he would get better of that; but this pounding in his head was more than he could stand: he knew that it would kill him.

The longer the anxiety persists the more does the pattern become set and the more difficult does it become to treat, and such an anxiety pattern is common to many psychotic and psychoneurotic conditions. It may persist in more or less the original form, or the object may be displaced so that a definite phobia or obsession develops. In our patient the fear of insanity was displaced by a fear of death, since the latter was a lesser evil to her; but much more elaborate distortions and displacements occur even to the systems of mental maladjustments of the schizo-

phrenic. So far as treatment is concerned the more complex is the pattern and the longer it has persisted the more difficult will it be to treat, but if it is not too rigid and there is still a possibility of reorientation and reassociation with other mental processes recovery is possible. This is the state of affairs in the psychoneurotic, and by patient treatment of an analytical type, not necessarily by orthodox psycho-analysis, the pattern can be unravelled, the various components separated, and each given its proper value and time relations. In this way the pathological anxiety is resolved and the patient is restored to a reasonable sense of security.

In the psychotic schizophrenic, however, the pattern has, for some reason which is obscure, become extremely rigid and association with other mental processes has become very difficult or impossible; the patient has lost insight, and, having retired from reality into fantasy, is no longer particularly interested in being restored to security in relation to the real world. It would not appear that the explanation of this is wholly to be found in the intensity of the anxiety or the extreme degree of the insecurity; there would seem to be some other factor, possibly chemical, in the sense of cell or synapse integrity, which intervenes, but as to the nature of which we are ignorant. In such "closed" systems, however, psychotherapy is of very little avail, but modern convulsive therapy may afford relief by breaking them up, leaving the patient less anxious and less enclosed in his retreat from reality, but often at least more simple-minded and at a lower level of mental capacity.

Roizin (1939) showed that convulsive therapy produced small areas of cerebral necrosis, and these would interrupt and break up the facilitated "circulation" through the anxiety pattern.

Prevention of Pathological Anxiety

If pathological anxiety is to be prevented the manner in which the pattern is formed is important. At present our concern is with prevention under war conditions, both in the Services and in civil life. Ideally, it might be held, the way to prevent this condition is to exclude all those who owing to unfavourable circumstances in early life are potentially insecure. As a matter of fact, however, this is not only quite impracticable but would unnecessarily exclude a great number of people who could and would perform useful service even in positions of considerable danger. This is because, as we have seen from the example quoted, in the genesis of pathological anxiety not only is there a substratum of insecurity determined in the early years, but there is also a paucity of weapons wherewith the individual can maintain himself in the struggle for existence. The early conditions of our patient and her sister were probably very similar, but the patient later came to depend on her intelligence, and her intelligence alone, whereas her sister had many more strings to her bow. When her intelligence was threatened, then came the persistent anxiety, whereas in the case of her sister quite a lot of attributes would have had to be threatened before her feeling of security in life would have been seriously undermined.

Of course, if it is possible to recognize in time the individual who has been insecure in childhood, and also has in his own eyes too narrow an equipment with which to meet life, that person would be better evacuated from the danger area if he is a civilian or removed from the obligation of active general service if he is a combatant. Only occasionally, however, can such people be recognized in time for complete prevention of breakdown, but

after an individual has been subjected to strain, and is clearly showing signs of serious anxiety, his previous reactions to life should be carefully investigated. Not only the competent psychiatrist but also every understanding physician with even a little psychological knowledge will be able to appraise the situation and will see the wisdom of removing such a person as is described above from the strain of a continued feeling of insecurity before he has completely broken down into a useless liability to the nation and later a continuous burden on the taxpayer.

REFERENCE

Roizin, L. (1939). *Riv. Patol. nerv. ment.*, **53**, 491.

MANAGEMENT OF RESPIRATORY DEPRESSION DURING ANAESTHESIA

BY

**F. BARNETT MALLINSON, M.R.C.S.,
L.R.C.P., D.A.**

Anaesthetist to Princess Beatrice Hospital, Royal Waterloo Hospital, and Woolwich Memorial Hospital; Honorary Assistant Anaesthetist to the National Dental Hospital

The rapidly increasing use of the intravenously administered "light" barbiturates in modern anaesthesia has brought into greater prominence problems involving respiratory depression. All the barbiturates used in this way are apt to produce respiratory depression, particularly following heavy premedication; but if this disadvantage is realized, and is faced frankly by the modern anaesthetist and countered by appropriate measures, it presents no serious bar to the employment of a technique which is undoubtedly one of the outstanding advances of recent years in anaesthetics.

The barbiturate which seems to be generally acknowledged to produce by far the best results is sodium pentothal (Tovell and Thompson, 1936; Heard, 1936; Murphy, 1936; Mallinson, 1937), but it can also give rise to the most pronounced degree of respiratory depression, especially if not handled by an experienced administrator. Very occasionally—but much less frequently than its predecessor, evipan—pentothal causes severe laryngeal spasm, though this happening is usually due to errors of technique, such as too slow or too rapid a rate of injection. Evipan causes considerably less respiratory depression, but it is definitely inferior to pentothal in its effects and less reliable in its results. What follows here is therefore chiefly concerned with the use of pentothal as the intravenous agent.

Methods of Correcting Respiratory Depression

There are two chief ways to correct respiratory depression severe enough to cause anoxaemia, arising during anaesthesia.

1. STIMULATION OF THE RESPIRATORY CENTRE

This can be done by drugs injected or by carbon dioxide inhaled. The latter is useful in some circumstances and can if necessary be combined with the second method to be described; but it has its contraindications, particularly where the anaesthetic technique includes spinal analgesia (Mallinson, 1938). The most useful and reliable