BRITISH MEDICAL JOURNAL

LONDON SATURDAY JANUARY 27 1940

GONORRHOEAL MYOCARDITIS

BY

OLE BANG, M.D.

(From Medical Department III, Kommunehospitalet, Copenhagen; Chief Physician, Poul Iversen, M.D.)

Heart lesions resulting from gonococcal infection are well known so far as septicaemia is concerned. Most frequently the ulcerating endocarditis is described, though it was emphasized many years ago (Councilman, 1893) that both the myocardium and the pericardium are often involved, so that a pancarditis ensues. These cases are rare. Rowlands and Simpson (1937), going through the London Hospital material from the years 1912 to 1936, were able to find only six such cases. Gillespie and Thompson (1937) report three instances of endocarditis among 1,917 infected persons; while Neumann and Bingenheimer (1937) among 5,262 cases saw only one case with septicaemia. Williams (1938), on the other hand, holds that gonococcal infection is responsible for 26 per cent. of all cases of bacterial endocarditis. In this connexion it must be pointed out, however, that while a true septicaemia, or bacterial endocarditis, is a rare occurrence in gonococcal infection, a transitory bacteraemia seems to be a fairly common event, judging from the frequency with which gonococci can be cultivated from the joints, bursae, etc., and even sometimes from the blood in complicated cases. A benign endocarditis has also been described, appearing as transitory murmurs, palpitations, and praecordial pain in ambulant patients with apparently uncomplicated urethritis. It is looked upon as a "toxic" manifestation.

Apart from the myocardial involvement in septic endocarditis, myocarditis is seldom established as a complication in gonococcal infection. Dittrich (1930) refers to two instances of isolated myocarditis complicating gonorrhoeal arthritis, showing tachycardia and arrhythmia and leaving a state of tachycardia after exertion; but reference to the literature of recent years reveals no other mention of isolated myocarditis except one. This, by Guldberg (1935), in Oslo, concerns a man aged 35 who had signs of myocarditis three weeks after urethritis; a partial auriculo-ventricular block lasted four months without any fever or arthritis. Besides, gonococcal infection is never mentioned in the current textbooks on heart diseases as a possible cause of myocarditis, though almost every other common infectious disease is recognized as such.

In view of this it is rather surprising that in the course of one year in an ordinary medical ward we should see the following series of cases in which gonorrhoeal myocarditis may be regarded as a probable diagnosis. It may mean that this condition, escaping detection since

it is not searched for, is more common than ordinarily believed. This, with some questions arising, is discussed below.

Case Records

[All negative findings are left out.

GR = Gonococcus complement-fixation reaction, carried out in the State Serum Institute.

SR = Sedimentation rate (millimetres in one hour).

Ecg = Electrocardiogram; the chest leads occasionally employed are those advocated by Larsen and Warburg (1937); in this modification the main deflections should be upright as in the usual leads.]

Case 1.—Man aged 33. Ten years ago primary urethritis, uncomplicated. Six years ago urethritis (probably recurrence) with severe complications: arthritis, tenosynovitis, metastatic conjunctivitis. GR +11. Subsequently treated with compligon and streptomid, as he was suffering from chronic urethral discharge and arthralgia. Now complaining of attacks of anginal pain and palpitations, and so admitted to hospital. Heart: auscultation and radiograph reveal no definite abnormality. Ecg (Fig. 1): P-Q = 0.25 sec.; P-Q remained pro-

FIG. 1.—Case 1.

longed in subsequent tracings, with small variations. Normally the P-Q interval is less than 0.18 second and never greater than 0.2 second. GR +5. SR 14.

Comment.—Signs of myocarditis are found in a case of chronic complicated gonorrhoea.

Case 2.—Man aged 52. Twenty-five years ago gonorrhoea accompanied by arthritis, treated in hospital. Sixteen and ten years ago attacks of arthralgia. A fortnight ago new attack

FIG. 2.—Case 2.

of arthralgia following a neglected sore throat and fever; admitted to hospital. A diagnosis of chronic arthritis, especially in a hip-joint and the knees, was established. Heart: a slight murmur heard; radiograph negative. Ecg (Fig. 2): P-Q more than 0.4 sec.; incomplete bundle-branch block. GR +1, 0, 0. SR above 100, falling.

Comment.—Signs of myocarditis are found in a case of chronic gonorrhoeal rheumatism.

Case 3.—Man aged 33. Admitted to eye department with severe double conjunctivitis (gonococci in the pus) three weeks after the primary urethritis and treated with streptomycin and milk injections. After a week transferred to medical department with very severe arthritis, especially of knees and ankles, fever (max. 39.8° C.), and anaemia (haemoglobin 45 per cent., red cells 2,500,000 per c.mm.). Heart: tachycardia; questionable murmur; questionable enlargement in x-ray film. Ecg (Fig. 3): Incomplete bundle-branch block (Bailey's Type 2). GR +8, +2, 0. SR 152, falling.

FIG. 3.—Case 3.

Comment.—Severe myocarditis develops in a case of acute gonorrhoeal arthritis.

Case 4.—Man aged 25. Uncomplicated gonorrhoea two years ago. Now admitted with arthritis, starting in an ankle and subsequently attacking several other joints, especially a hip-joint and the wrists. Auscultation and radiograph reveal a fresh transitory pleurisy on the right side. Temperature up to 39.9° C. Heart: passing pain in the praecordium; soft systolic murmur heard; radiograph of heart negative. Ecg (Figs. 4a, 4b, 4c): Changes are noted as follows: (a) low T₁; (b) P-Q above 0.2; T₃ now negative; Q₃ questionably enlarged; (c) P-Q and T summits now normal. GR +1, +1, +2, +1, +1. SR 116, falling subsequently.

Comment.—There can be little doubt that the arthritis is gonorrhoeal, and so is probably the pleurisy. Signs of an acute transitory myocarditis are demonstrated.

FIG. 4a.—Case 4. Low T₁. FIG. 4b.—Case 4. P-Q above 0.20; T₃ now negative; Q₃ questionably enlarged.

FIG. 4c.—Case 4. P-Q and T summits now normal.

Case 5.—Man aged 36. Diphtheria four years old. Rheumatic fever (?) about the age of 9 or 12, leaving no ill-health. Uncomplicated gonorrhoea as a youth. Three years ago "rheumatic fever," treated in another hospital; arthritis of a shoulder, elbow, and knee were found, and the GR was positive (+3, +2, +1); SR 117, falling. There were albumin, erythrocytes, leucocytes, and epithelial cells in the urine. The prostate was normal. A month ago "rheumatic" pains in the chest and the right wrist, subsequently in the right hip, following a sore throat and a motor-car trip during which he was exposed to cold and draught. Admitted with arthritis, especially of the right wrist, and fever (max. 39.3° C.). Anaemia developed (haemoglobin 75 per cent.). The urine was normal. Heart: pain in the sternal region for about two weeks (coinciding with changes in the electrocardiogram). Ecg (Figs. 5a, 5b); In Fig. 5b, a month after Fig. 5a, T₁ is flattened and T₂ is negative; three weeks later the electrocardiogram (not shown) is exactly like Fig. 5a. GR +2, +7. SR 109, -127, -19.

Comment.—The rheumatic infection is dubious, while it is known that gonococcal infection has taken place. During the attacks of arthritis observed in hospitals the GR was positive; so there is little doubt that the gonococcal infection is the cause

of the recurrent arthritis. The changes in the electrocardiogram point to transitory myocardial damage. It cannot be denied that the anaemia may be instrumental in producing

FIG. 5a.—Case 5.

FIG. 5b.—Case 5; a month later than Fig. 5a.

such features in the electrocardiogram, even though frequent readings did not show less than 75 per cent. of haemoglobin. It seems most probable, however, that this is another instance of transitory myocarditis occurring in a case of gonorrhoeal arthritis.

Case 6.—Man aged 26. Diphtheria with croup when 4 years old, treated with serum and tracheotomy. Scarlet fever at age 6 to 7; gonorrhoea at 18 and again five months ago. One month ago tenderness of left heel and a red and swollen Achilles tendon. Ten days ago fever ($39^{\circ}\text{C}.$) and pain in the left hip-joint and a toe. Admitted to hospital with slight

FIG. 6.—Case 6.

arthritis in the hip and toe, and fever ($38^{\circ}\text{C}.$); in a few days all symptoms had disappeared. Heart: no murmur; no subjective symptoms. Ecg (Fig. 6): In the first electrocardiogram

(not shown) the P-Q was 0.23 to 0.25 sec.; in the next one, three weeks later, the P-Q was 0.31 to 0.32 sec. GR 0, 0. SR 33, subsequently falling.

Comment.—The negative GR does not eliminate gonococcal infection, as several cases of complicated gonorrhoea (salpingitis; arthritis with gonococci in the pus) are known to occur with a negative GR. The fact that the P-Q is changing seems to exclude the possibility of a long-standing myocarditis. A gonorrhoeal myocarditis affords the most reasonable explanation.

Discussion

The common feature in these six instances of myocarditis is the occurrence of the condition in carriers of gonococcal infection. Though there are no means available for the establishment of definite proof, there is a very strong suggestion indeed that gonococcal infection is the cause. Endocarditis seems to accompany myocarditis in some of the cases, but we shall go no further into that, since it is not, as is myocarditis, demonstrable by objective methods. The fact that some of the patients did not show any signs of heart involvement on routine examination seems to support the view that myocardial damage may escape notice if not looked for. Accordingly the incidence of this complication may be higher than is assumed at present.

The *pathogenesis* of the myocardial lesion is not clear. In septicaemia or bacteraemia there may be propagation from the endocardium or from bacterial foci in the myocardium. It is more probable, however, that in the majority of cases the myocardial damage is brought about by some agent originating in a distant focus—for example, in the genital sphere or in a joint.

The *prognosis* is varying. In Cases 4, 5, and probably 6, the myocarditis seems to be a passing event, corresponding with the banal changes often met with in other infectious diseases, as, for instance, influenza and pneumonia. There is little doubt, however, that the outlook is much graver with regard to the two patients (Cases 2 and 3) who left hospital with a bundle-branch block. A follow-up examination of Case 1, carried out almost a year after the onset of his cardiac symptoms, revealed that he was still out of work owing to his attacks of pain and palpitations, and the electrocardiogram showed lengthening of the P-Q, besides a slurring of the QRS; the complement-fixation reaction has remained positive (+ 5).

There is, in other words, the possibility that some of these patients will return to hospital sooner or later with chronic "degenerative" heart disease. The question is: Will the aetiology suggest itself by that time? The question may be reversed: When we treat to-day a series of cases with degenerative (and valvular) heart disease of obscure origin, must we not suspect some of them to be the result of a remote but probably somewhere persisting gonococcal infection?

When the gonococcal complement-fixation reaction was introduced into the routine examination of all acute arthritis cases we learnt that a great proportion of seeming rheumatic fever cases were in reality gonorrhoeal rheumatism. *A priori*, it seems not improbable that some of the "rheumatic" heart diseases should be unmasked in the same way. In making the diagnosis of rheumatic heart disease we may have been placing too much confidence in the information we were getting about rheumatic arthritis in the anamnesis; no doubt a good many cases of gonorrhoeal arthritis have sailed under that false flag. It may prove to be worth while to search the anamnesis for gonorrhoea and its complica-

tions, just as we are looking for syphilis and rheumatic fever.

A prominent feature of gonococcal infection is its obstinate persistence—that is, the tendency to formation of chronic foci. This fact seems to reflect itself to some extent in the course of the complement-fixation reaction. Investigation into this reaction is now being made in co-operation with Dr. Peter Krag of the State Serum Institute. One of the results may be mentioned in this connexion—namely, the surprisingly high incidence of positive reactions found among patients without apparent signs of recognized sequels to gonococcal infection. The Wassermann reaction has aided in the evaluation of many a syphilitic manifestation that would scarcely have been diagnosed as such without this serologic evidence. Perhaps in future the gonococcal complement-fixation reaction will help to reveal that chronic gonococcal infection plays a part in one or more diseases not hitherto attributed to this cause. Certain heart diseases may be among them.

Summary of Conclusions

Apart from the myocardial involvement in the bacterial endocarditis of gonococcal septicaemia, myocarditis is seldom established as a complication in gonococcal infection. In view of this it seems surprising that a series of six instances of what is regarded as gonorrhoeal myocarditis was encountered in the course of one year in an ordinary medical ward. It seems probable that this condition often escapes detection, since suggestive symptoms may be absent.

The diagnosis was made on the ground that all six cases had changes in the electrocardiogram occurring in the course of acute or chronic complicated gonococcal infection. In some instances the myocarditis was slight and transitory, as may be the case in many simple infections—for example, pneumonia. However, in two cases incomplete bundle-branch block developed and persisted, and in a third case the anginal pain still troubled the patient considerably when he was re-examined a year later. Such a course suggests the possibility that some of these patients will return to hospital sooner or later with chronic heart disease. It is possible, also, that some of the chronic “degenerative” heart diseases which we see to-day arise from gonococcal infection.

The gonococcal complement-fixation reaction has been instrumental in pointing out how great a proportion of so-called rheumatic arthritis cases were really gonorrhoeal. It may be worth while to search the anamnesis of heart patients for gonorrhoea as we search it for syphilis and rheumatic fever, and the complement-fixation reaction may reveal that some instances of heart lesion attributed to rheumatic infection ought to be regarded as following on persistent gonococcal infection.

REFERENCES

- Councilman, W. T. (1893). *Amer. J. med. Sci.*, **106**, 277.
Dittrich, O. (1930). *Jadassohn's Handb. Haut. u. Geschl.*, **20**, pt. 2, J. Springer, Berlin.
Gillespie, J. O., and Thompson, R. M. (1937). *Milit. Surg.*, **80**, 418.
Guldborg, G. (1935). *Norsk Mag. Laegevidensk.*, **96**, 576.
Larsen, K. H., and Warburg, E. J. (1937). *Hospitalstidende*, **80**, 276.
Neumann, H., and Bingenheimer, E. (1937). *Derm. Wschr.*, **105**, 1099.
Rowlands, R. A., and Simpson, S. L. (1937). *Brit. J. ven. Dis.*, **13**, 215.
Williams, R. H. (1938). *Arch. intern. Med.*, **61**, 26.

Bronchography.—The authors describe a method by which radiographs may be obtained showing the lipiodol-filled bronchi standing out in relief. Some excellent examples are included in the article.—*Bronchography in Relief*. M. R. Castex, E. Alvarez, and E. S. Mazzei.—*Arch. méd.-chir. Appar. resp.*, No. 6 (1938), **13**, 441.

THE PREVENTION OF ANXIETY

BY

R. G. GORDON, M.D., F.R.C.P.Ed.

Physician, Royal United Hospital, Bath

A young happily married woman aged 30 with two healthy children is the victim of intense anxiety if she goes out alone. While she is working about the house she is all right. If she goes out in the company of certain people she is practically all right. The best “protection” in this respect is her husband, next her sister, then her doctor—the general practitioner who looks after the family, but any doctor is “some use.” Other people help but do not abolish the anxiety.

Physically she is healthy and any organic disease can be excluded, but in addition to the anxiety she complains of uncertain sleep, dull headache, poor memory, and inability to concentrate. Asked to describe the anxiety, she states that it seems that something dreadful is going to happen, probably that she is going to die. Her husband would prevent it happening and doctors would see that it didn't. The anxiety was not always a fear of death; it used to be a fear of going insane, but that has now changed to a fear of death. The anxiety is obsessional in nature—that is, intellectually she realizes that it is irrational, but emotionally she is quite powerless to overcome the feeling.

When asked what to her is the significance of death, she says she is not sure, but states vaguely: being separated from those she knows and loves, being “shut up in a box.” She hopes that she would be liberated and reunited to her loved ones after death, but is very uncertain and muddled about this. Questioned about going insane, she replies with a heated diatribe against the iniquity of the system by which doctors can shut people up for no adequate reason. Pressed on this point she said, “Of course, good doctors would not do this, but it is scandalous how easily it could be done.”

The foundation for this attitude seemed to be that an aunt of her husband's had been “put in an asylum when she was really quite sane.” This was interesting, for this aunt was clearly a psychopathic individual, was an incorrigible thief, and, by what possibly was somewhat of a “wangle,” had been represented in court as irresponsible for her actions and confined in a mental hospital instead of in a prison, a procedure which apparently was regarded as a lesser evil by most of her relatives but as a scandalous crime by our patient.

The real basis of the present anxiety seemed, therefore, to be a fear of going out of her mind and of being shut up away from her friends. The fear had later been transferred to death as a lesser evil, since if she was dead she herself would not know about it. A light is thrown on this by her previous medical history. She had been quite healthy, apart from one or two exanthemata in childhood, till a very short time before her marriage, when she had an attack of appendicitis which necessitated operation and the postponement of her wedding. This was attended by no anxiety whatever, only exasperation at the upset of arrangements. A severe attack of influenza after her marriage left her similarly emotionally unscathed. Her next illness six months later, however, had very different results. At the time it seemed trivial: she knocked her head on a low door and sustained slight concussion. She was not rested enough and began to have headache, loss of memory, and poor concentration. These symptoms,