to Mr. M. Raeside, L.D.S., senior school dental officer, City of Coventry Education Department, for his helpful criticism.

REFERENCES

- Cartin, B. (1938). *Dent. Items*, 61, 315.
 Cartwright, F. F. (1939). *British Medical Journal*, 1, 1031.
 Feldman, M. H., and Cartin, S. (1938). *J. Amer. dent. Ass. dent. Cosmos*, 25, 616.
 Goldman, V. (1936). *British Medical Journal*, 2, 122.
 — (1937). *Ibid.*, 2, 1265.
 Gunter, J. H., Beach, E. W., and Looby, J. P. (1936). *Int. J. Orthod. oral Surg.*, 22, 746.
 Lyons, S. S., and Frank, H. P. (1939). *J. Amer. dent. Ass.*, 26, 580.
 Ramage, G. (1937). *Medical Officer*, 57, 45.
 Roche, L. S. C. (1937). *J. R.A.M.C.*, 69, 399.
 Rosamilia, P. A. (1938). *J. Amer. dent. Ass. dent. Cosmos*, 25, 1465.

AN INQUIRY INTO THE HOUR OF BIRTH

BY

V. SPILLER, M.D., M.R.C.S., L.R.C.P., D.P.H.

Obstetrical and Gynaecological Unit, Royal Free Hospital, London

An investigation into the relative frequency of deliveries by day to deliveries by night has been carried out in a small series of 2,225 patients delivered per vaginam during the years 1934-7 at the Royal Free Hospital. The following table shows the number born during these years, arranged according to the time at which the birth occurred.

TABLE I
Births by Day

	8-10 a.m.	10-12	12-2 p.m.	2-4 p.m.	4-6 p.m.	6-8 p.m.	Total
Primiparae:							
1934 ..	11	29	27	28	25	26	146
1935 ..	27	42	36	26	42	26	199
1936 ..	33	39	44	56	53	45	270
1937 ..	43	42	39	38	41	41	244
Total ..	114	152	146	148	161	138	859
Multiparae	32	25	44	36	52	36	225
Total ..	146	177	190	184	213	174	1,084

Births by Night

	8-10 p.m.	10-12	12-2 a.m.	2-4 a.m.	4-6 a.m.	6-8 a.m.	Total
Primiparae:							
1934 ..	29	23	16	23	24	17	132
1935 ..	29	36	42	46	40	23	216
1936 ..	48	42	45	44	41	37	257
1937 ..	51	42	43	42	49	35	262
Total ..	157	143	146	155	154	112	867
Multiparae	47	43	46	43	45	50	274
Total ..	204	186	192	198	199	162	1,141

If we reckon "night" as from 8 p.m. to 8 a.m. there are over the total series 5.25 per cent. more births by night than by day, the preponderance of night births over day births being smaller in the primiparae than in the multiparae. Only 0.9 per cent. more primiparae are delivered by night than by day, whereas the corresponding figure for multiparae is 22 per cent. It would be unwise, however, to draw any inference from this variation, considering the small numbers in our group of multiparae.

The following table shows the comparison between night and day figures for primiparae only.

TABLE II

1934 ..	146	by day and 132	by night	(100 day, against 90.4 night)
1935 ..	199	" " 216	" "	(" " 108.5 ")
1936 ..	270	" " 257	" "	(" " 95.2 ")
1937 ..	244	" " 262	" "	(" " 107.4 ")
Total ..	859	" " 867	" "	(" " 100.9 ")

It will be seen that the yearly figures vary slightly around 100 (-9.6 per cent. to +8.5 per cent.), and for 1934 and 1936 show a smaller number of births during the night than during the day. These variations are well within the limits of probable error, and suggest a comparatively even distribution of births over night and day. When, however, the records as to the hour of birth in other centres [as published by Guthmann and Bienhüls, 1936, Dyroff and Mulhäuser (Ritala, 1931), Lane (Halban and Seitz, 1927, 1929), Meyer, 1935, Knapp, 1909-10, Lynch, 1907, Veit (Nathanson, 1935), White (Puppel, 1932), Birlinski (Quigley, 1931)] are studied, it is found that in over 180,000 cases there is a slight predominance in the number of cases delivered by night over those delivered by day. It must be noted here, however, that the definition of "night" varies between 7 p.m. to 7 a.m. and 10 p.m. to 10 a.m., according to the different authors. Meyer (1936), Jenny (1933), and Guthmann and Bienhüls (1936) find that year by year the number of children born by day is increasing. In our small series no such increase is found.

May one perhaps assume that the great diversity of opinion in the literature as to the hour when most births occur is some support for our contention that the distribution throughout night and day is relatively even? The following is a list of the most usual and the least usual incidence of the hour of birth according to different writers:

	Maximum	Minimum
Royal Free Hospital ..	4 p.m.	8 a.m.
Knapp ..	11-12 p.m.	6-7 p.m.
Jenny ..	5 p.m.	5 a.m.
Veit ..	1 a.m.	5 p.m.
Guthmann and Bienhüls ..	11 a.m.	—
Birlinski ..	4 a.m. and 1 p.m.	7 a.m. and 8 p.m.
Lynch ..	8-9 a.m.	1-2 p.m.
Henry (1932) ..	3 a.m.	2 p.m.

Summary and Conclusion

An investigation into the relative frequency of deliveries by day to deliveries by night has been carried out in a small series of 2,225 patients delivered per vaginam.

The variations are well within the limits of probable error, and suggest a comparatively even distribution of births over night and day.

I am deeply indebted to Professor Amy Fleming for her constant help and encouragement during the course of this work.

REFERENCES

- Guthmann, H., and Bienhüls, M. (1936). *Msehr. Geburts. Gynäk.*, 103, 337.
 Halban, J., and Seitz, L. (1927). *Biol. Path. Weibes*, Berlin, 7, 38.
 — (1929). *Ibid.*, 8, 1906.
 Henry, J. R. (1932). *Marseille méd.*, 1, 203.
 Jenny, E. (1933). *Msehr. Kinderheilk.*, 56, 165.
 Knapp, C. B. (1909-10). *Bull. Lying-in Hosp. N.Y.*, 6, 69.
 Lynch, F. W. (1907). *Surg. Gynec. Obstet.*, 5, 677.
 Meyer, M. (1935). *Med. Welt*, 9, 1856.
 — (1936). *Ibid.*, 10, 907.
 Nathanson, J. N. (1935). *Amer. J. Obstet. Gynec.*, 30, 159.
 Puppel, E. (1932). *Zbl. Gynäk.*, 56, 3067.
 Quigley, J. K. (1931). *Amer. J. Obstet. Gynec.*, 21, 234.
 Ritala, A. M. (1931). *Acta obstet. gynec. scand.*, Supp. 1, 11, 1.

Dr. Lucien Dautrebande, professor of pharmacology at Liège University and member of the Royal Academy of Belgium, has been awarded the Montyon prize for his work on the toxicology of war gases and the vapours of solvents used in industry.