my own ulcer treatment it merely implies that this is the way in which we manage it in Denmark, at Bispebjerg Hospital, in my department. It consists of rest in bed, though not absolute, for four to five weeks, a soft non-irritating diet with increasing variety during the same period, frequent meals—that is to say, every second or third hour—alkali, and belladonna; tobacco is forbidden. How much importance attaches to the last restriction I shall not go into. We endeavour to make the diet qualitatively sufficient, but I must admit that in certain ways it perhaps leaves something to be desired, and it is conceivable that we ought to pay more attention to remedying possible vitamin deficiencies.

On discharge from the hospital the patients are urged to avoid the most irritating kinds of food for six months, to continue taking frequent meals, to restrict the use of tobacco and not to let "a smoke" take the place of a snack between meals, and also to try to adopt a calmer attitude towards life and its incidents.

Summary

The treatment of bleeding ulcer is first discussed because experience with a free diet has influenced the outlook on ulcer treatment in general.

The results of treating 491 cases of profusely bleeding ulcer are given.

The conclusion drawn is that the more severe the bleeding the more the need for fluid and food. Operative treatment with a view to haemostasis is not advised.

In non-bleeding ulcer it is maintained that use of a freer diet gives improved results; treatment is supplemented with alkalis and belladonna.

The view is held that the basic principles of the general medical treatment of ulcer should be rest, relaxation, and adequate food taken at frequent meals.

REFERENCES

- Davies, D. T., and Wilson, A. T. M. (1937). *Lancet*, 2, 1353.
 Meulengracht, E. (1935). *Ibid.*, 2, 1220.
 Spriggs, E. I. (1909). *British Medical Journal*, 1, 825.
 Witts, L. J. (1937). *Ibid.*, 847.

Public Social Services is the title of a handbook produced by the National Council of Social Service with the assistance and approval of various Government Departments, describing in a simple and understandable form the many Acts and Regulations which provide State services affecting the well-being of the general public, their method of administration, and the names and addresses of the respective authorities to whom inquiries and appeals may be addressed. This handbook was first published as a pamphlet in 1917, and it has been revised and enlarged several times since then. The principal differences of administration in Scotland are noted, and information as to how the many social enactments apply to that country has been prepared in co-operation with the various departments of the Scottish Office. Sir Ronald Davison writes on the part played by voluntary organizations and voluntary social workers in the administration of the public services. The chapter on public health deals with infectious diseases, housing, town and country planning, abatement of nuisances, maternity and child welfare, mental deficiency, the welfare of the blind, and the prevention of blindness. The National Health Insurance and Pensions Acts are fully explained. The chapter on the Public Education Services includes sections on elementary and higher education and school medical services, and special reference to neglected and delinquent children. Copies of *Public Social Services* may be obtained through any bookseller or direct from the National Council of Social Service, 26, Bedford Square, W.C.1, price 2s. post free.

FOOD CONSERVATION IN RELATION TO NATIONAL FOOD SUPPLY*

BY

SIR JOSEPH BARCROFT, F.R.S.

Late Professor of Physiology, University of Cambridge

In this paper I deal with the ways in which food conservation can help the nation in times of emergency. With matters of policy I have no concern. How much of this or that the Government has seen fit to store is not my affair: what I can do is to say something about how science has increased the selection of food at their disposal.

In my own mind foodstuffs for the present purpose fall into three classes: (1) those which can be kept indefinitely—in practice, say ten years; (2) those which can be kept for a season, say ten months; (3) those which can be kept on any ordinary voyage, say two months or less.

Foods that can possibly be Kept Over Ten Years

The first food in this group to occur to anyone would be wheat. Has wheat any valid title to come into this class? The storage of wheat as against emergencies is as old as the book of Genesis. You remember that Joseph had his seven-year plan, which of course was really a fourteen-year plan. This fourteen-year plan is just a little interesting, because not all wheat will keep for fourteen years—in fact, not all wheat will keep for ten years; and, remember, Joseph kept the wheat in Egypt, as we would say under tropical conditions. The higher the temperature at which the food is kept the more difficult it is to store. Speaking generally, most chemical changes are doubled in velocity, or more than doubled, for every 10° rise in temperature. That law, perhaps, does not apply rigidly to a live foodstuff like wheat. Thus, if we, on the one hand, can store without refrigeration at 12° C., and Joseph, on the other hand, had to compete with 42° C., we should be able to keep our stocks eight times as long as Joseph. If he could keep wheat ten years, we should be able to keep the same wheat eighty years; or if he could keep something one year, we should be able without refrigeration to keep it eight years or, by reducing it to two, sixteen years. I mention this not as an idle speculation, but because these considerations are the basis of what we call accelerated tests. If you ask me how I know that a particular sample of food can be kept eight years when it has never actually been kept that length of time, the answer is: I cannot say positively, but if it can be kept one year at 42° C. the chance is sufficiently in favour of our being able to store it for eight years at 12° C. (53° F.) to make that chance worth taking. We are not absolved from the duty of carrying out tests over long periods of time, but meanwhile we have a good deal to go upon by the performance of such accelerated tests as I have described. But to return to Joseph and his wheat. If he was handicapped by the temperature at which he had to store the wheat he had another factor very much in his favour—namely, the dryness of the atmosphere in which the wheat was stored. Not all wheat keeps indefinitely: the critical characteristic appears to be the water content. In the best wheats the water content is low and the wheat will keep. Fortunately some of the best-keeping wheats from this point of view are grown in one of the Dominions. The problem of growing wheat of high keeping property in this country is primarily

* Read in opening a discussion in the Section of Medical Sociology (Nutrition) at the Annual Meeting of the British Medical Association, Aberdeen, 1939.

one for the scientific farmer, and not for the Food Investigation Board; but in the meantime it does not matter very much that the storable wheat comes from over-seas, because the demand for grain so far outruns the home supply that home wheat can be eaten at once.

I wonder whether there were weevils in Egypt. These now constitute a major problem in the storage of wheat over long periods, but they do not, I hope, present one which is insoluble. There is one phase of the subject which is immediately within our purview. Wheat is very bulky. Is there any way in which we can store all that is worth storing in a smaller space and possibly in a more accessible form? This brings me to the storage of flour. I also wonder in what form Joseph's wheat was eaten. It is one thing for wheat to keep, to all appearances, well, but before it deteriorates in appearance it will go back as a material from which to make first-rate flour. Canadian wheat can be kept seven years in this country, and Canadian wheat in Canada has, when sown after ten and a half years, yielded 90 per cent. of grains which have germinated.

Wheat contains about 15 per cent. of water. It is possible to reduce the water content of flour to about 11 per cent., in which case the flour may be stored in tins. In that way it will keep a long time, and, what is very important, is inaccessible not only to water but to weevils. The general conception of adding to the length of storage life by drying has been extended to potatoes. Dried potatoes come into the ten-year category, but of them I treat later. So far I have not touched on the storage of fats in bulk. In this connexion I need only mention that whale oil and vegetable oil, with all their calories, are eminently storable substances. Whale oil can be kept in bulk if air be excluded, perhaps not for ten years, but for something much more than a season, say, three or four years, and if hydrogenated—that is, passed through the first operation in making margarine—the period would be extended.

Quite a different class of material that falls into the category of foods which can be kept for more than ten years is that which embraces canned meat of various sorts. There is no question that if canned meat is properly packed it is entirely suitable for food for a very long time. Of course, such meat has been sterilized by heat, and to that extent its vitamin value has probably been reduced: the comparison is not between canned meat and raw meat, but between canned meat and cooked fresh meat, so that the loss of vitamins does not come into the picture. The subject of flavour raises a point of great interest. We have four senses which matter as regards our judgment of meat—taste, smell, sight, and touch. So far as we know, if the meat will pass all of these its nutritive value is maintained. Perhaps the sense of taste is the most exacting, especially for the fat of meat, or indeed the fats of anything. A very trifling degree of oxidation of fat will alter the flavour of food, be it meat, milk, fish, or even food containing so little fat as biscuits; and in the case of fat the taste will tell the tale before the fat loses either in calorific value or in vitamin value or produces substances irritant to the digestive system.

There are a great number of canned foodstuffs that might be reviewed one by one if space permitted. One in particular is of interest because of its mixed character—a commodity defined in the Concise Oxford Dictionary as “tinned stew or part of the Army rations” (canned meat and vegetable rations). Here I will only mention the very satisfactory result, namely, that whereas the calorific value of the commodity was all that could be

desired, it was found to be deficient in vitamin—a very important research—and this deficiency can easily be made good by putting in new potatoes. Much work has been done on this subject of the nutritive value of the protein in preserved foods. On the general subject of the nutritive value of canned foods we are fortunate in having Mr. W. J. Godden here to-day to address us.

The nutritive value of canned foods generally seems to be covered by a sentence or two from a statement issued by the Food Investigation Board (F.I.B. Report, 1937, pp. 3–4), on the authority of the Medical Research Council: “The available evidence therefore suggests that modern methods of storing foods cause little depreciation in their nutritive value—in fact, it may be said that food of good initial quality that has been stored by the best modern methods is likely to be superior in many respects to similar food that, though still technically fresh, is in reality stale.” A caveat is then put forward with regard to the destruction of vitamins C and B₁ by heat; but this caveat becomes less urgent, though the vitamins do not become less necessary, if the comparison is between canned food and cooked fresh food.

Seasonal Storage

I must pass to foods which can be stored over a season. The point of course is this: Nature produces spasmodically; man eats at a steady rate. Nature produces in the summer; man eats all the year round—in fact, theoretically he eats most in the winter. Therefore, apart from what we may import from southern climates, one or both of two things must happen: either production must take place on a scale vastly greater than is required so that at its minimum it suffices, or else the excess of summer must be laid up for the winter months. As the former policy is eminently one of waste, it follows that, emergency or no emergency, it should be the aim of science to replace it by the latter; but of course the greater the emergency the greater the urgency. During the last war it was in the months of March, April, and May that the pinch came.

MILK

Of all the foodstuffs which call for consideration in this category the first place must undoubtedly be given to milk, for four reasons: (1) its importance—it contains practically all essential nutritive elements and food factors in an easily digestible form; (2) the principles enunciated above apply so appositely to it; (3) “sciencing” has already gone far to remove the seasonal trouble; (4) the nutritive value of milk has been studied so completely that it may serve for purposes of scientific discussion as a sample of other foods which have as yet been subjected to much less careful scrutiny.

The paramount importance of an ample supply of milk all the year round I need not emphasize. “All the year round” means, or has till recently meant, that the cows of this country must produce daily, even in winter, a quantity of milk ample for the needs of the community: apart altogether from the inferior quality of winter milk, the implication is that in summer there is a superfluity. This milk is made into cheese—perhaps the oldest way of processing milk—and butter, to say nothing of collar-studs, shoe-horns, and the like. Experimental evidence has been adduced that the best milk powders made from summer milk are of greater nutritive value than fresh winter milk. If this be once accepted as a fact its implications are enormous and far-reaching.

But how long can it be kept? Am I justified in placing the best forms of dried milk in the “seasonal” class?

The experiments that have been carried out by quite a number of observers, including the Low Temperature Station, the Dairy Institute at Reading, and Dr. Wright and his staff at the Hannah Institute of Dairy Research, leave no doubt upon this point. In fact, they raise another—namely, would I be justified in putting some dried milks in the ten-year class? The answer, I think, at the present time is that I would not, unless perhaps the milk was held at what I may call refrigeration temperatures. Were I to do so I would have to compromise on the flavour of the milk, though not, I believe, on its nutritive value.

Here I should say that the keeping power of the milk depends upon the way in which it is stored. In fresh milk the onset of *sourness* is the test, but that is not so in the case of dried milk. The fat is the tell-tale, and the chemical change which “turns” the fats is an oxidative one; therefore the secret in keeping sprayed milk powder is in the canning of it in such a way as to exclude air with the required degree of rigidity. Before leaving the question of milk I must not forget that, whatever importance or unimportance winter milk may have, cows and other animals must be kept through the winter. This presents another problem—namely, the seasonal preservation of grass. This matter I merely touch on in passing because it is only remotely of interest to the doctor, but it is one to which the Agricultural Research Council has given a great deal of attention. Apart from making hay—a wasteful process, viewed from the angle of the maintenance of nutritive value—there are two main processes of preserving grass: the making of silage and the desiccation of the grass. The former process has been greatly facilitated and cheapened by the introduction of silos made of paper suitably treated and reinforced; you buy it in rolls, and in a short time can put up a silo anywhere, given the necessary drainage.

POTATOES AND VEGETABLES

Of human foods, considered with regard to seasonal storage, perhaps the next most important to milk is potatoes; but potatoes as a merely seasonal commodity do not offer a problem to such a body as the Food Investigation Board. There is, however, this point: Can potatoes be so preserved as to place them in the ten-year class? Great advances have been made in this direction, and the probable answer is “Yes.” The fundamental method is drying till the water content has been reduced to about 8 per cent. Dried potato products, however, present this difficulty—namely, that when an effort is made to reconstitute them into washed potato they come out largely as starch paste. The guiding principle is that in the process of preservation the cells of which the potato is made must be preserved intact, and there are on the market preparations, notably in the form of powder, which are quite satisfactory as regards their keeping quality and the palatability of the meal they afford.

Carrots are particularly important because of their combined calorific and vitamin value. Here again the responsibility lies jointly with the agriculturist and with the food preserver; to the former the problem is that of saving the crop from the devastation of fly; on the latter lies the responsibility for the preservation of the crop. Allusion will be made to green vegetables later.

FRUIT AND EGGS

Among the most obvious and most important of seasonal products is fruit. Dr. Kidd has perhaps done more to put the storage of fruit on a sound basis than anyone else. The improvement in the quality of fruit

obtainable the year round and the increase in the variety are among the most astonishing features in our time

Among the important foodstuffs which can be made seasonal in the fresh state are eggs. The housewife was primarily responsible for the elevation of the egg from being a commodity which could be kept for days or weeks to one which could be stored over a season. The water-glass egg should keep at least a year; but at the end of that time, while you cannot call it a bad egg, it surely has severe limitations. True, it is good for cooking purposes, and that is saying a great deal, because the majority of eggs are, after all, used in this way; and it is good for scrambling. However, when it comes to frying, poaching, and above all boiling, something better is required. Moreover, as a large-scale commercial product you have to move the water-glass as well as the eggs. Recent work has produced a great change. The whole question of the effect of storage methods in the various constituents of the egg has been gone into. The three important factors are: (1) the temperature; (2) the moisture; and (3) the carbonic acid content of the air. An index of the improvement is the fact that in the Low Temperature Station at Cambridge they can actually hatch out 25 per cent. of the eggs in a setting after more than a month's storage. The results of a nine-months test may interest you.

Condition of Eggs at End of Nine Months' Storage

Cold Storage between 29° and 32° F.	Cold Storage + 2–2.5% CO ₂ + 85% Relative Humidity (29°–32° F.)	Cold Storage + 60–100% CO ₂ + 100% Relative Humidity (29°–32° F.)
Shell smooth	Shell slightly rough	Shell slightly rough
Air-cell large	Air-cell large	Air-cell small
White distinctly yellow	White faintly yellow	White faintly yellow
Consistency of white “runny”	Consistency of white satisfactory	White rather “runny”
Yolk flabby	Yolk firm	Yolk firm
Storage taste	No storage taste	No storage taste

The best method, then, is that of cold storage in an atmosphere containing 2 to 2.5 per cent. of CO₂ and with the air highly but not completely saturated with aqueous vapour. Eggs so kept, as well as being satisfactory for cooking purposes, will poach and fry well and, except for those who are so fastidious as to demand a creamy white, are satisfactory for boiling.

Short-term Storage

Lastly, we must consider the storage of foodstuffs over the period of a voyage, and two months will cover a journey from the Antipodes.

The importance of two months' storage is as follows: First, there are many commodities, such as meat and fruit, of which not enough is produced in this country to supply the needs of the nation, whether in peace or war, and to obtain sufficient of these things we must import them from distant quarters of the globe. But in emergency there is this further point: we cannot ignore the possibility of a reduction in our supplies from European countries—even friendly ones. It is important, therefore, to be able to make the most of what we can obtain from sources less accessible to those concerned with the curtailment of our supplies.

FISH AND MEAT

Fish offers one of the most interesting problems. Perhaps there is no food the freshness of which is more

important. Yet much of the fish arriving in our ports has deteriorated a good deal even before it is set on shore. On a winter's day I once watched the catch laid out for auction in a great fishing port. Stormy weather had kept some of the boats for as long as nineteen days at sea. The fish, of course, was ice-packed, but, even at that, much had deteriorated palpably. It was to me a pathetic sight to consider the expenditure of effort, of money, of hard-ship, perhaps even of life, that lay behind that catch of now indifferent fish. That, however, can be improved upon and all our principal food fishes can be brought into the season class. All important species can be kept in good condition for six months—the fatty fishes perhaps for no more, but the white fishes for longer.

The kipper, again, has normally a life of about a week; but arrangements could be made for herrings landed in Aberdeen to be sold as perfectly good kippers in Australia; and if they can be sold as good kippers in Australia, so they could be here after the same interval.

Another interesting fact which attaches to the preservation of fish is that among foodstuffs at present it stands out as one of which the deterioration can be assayed by chemical methods of the utmost delicacy.

Half of our meat supply comes from over-seas. This fact is so familiar that I need say very little about it, except this: in theory meat can be kept for a very long time. A Siberian physiologist claims to have made the heart of a dog beat thirty-one days after the death of the dog, the body being kept cold. Believe that or not as you please, there is no doubt that American workers have stored meat under sterile conditions—chilled but not frozen—for a year; the flavour had deteriorated, but the chemical changes that had taken place were only very trifling. The storage of meat, therefore, whether mutton or beef, is really a problem in cleanliness. Two factors should be realized: (1) that the meat should throughout its history come in contact with the fewest organisms; (2) that it should be kept under conditions as unfavourable as possible to the growth of such organisms as come its way. Both conditions are very much the business of the Food Investigation Board and of its associates in the Dominions: the second has been met by the gas storage of chilled meat.

Preservation over Shorter Periods

The subject of preservation over shorter times and shorter journeys, the insurance that lettuces or strawberries should reach the customer as fresh as when they were picked, is one of which I will only treat very lightly. First let me say that I have been taken to task for putting strawberries and lettuces in the same category. From the point of view of transport they are similar, but in importance they differ greatly: strawberries are a luxury; lettuces are the very type of green vegetable so important for its vitamin value. The storage of hundreds of tons of lettuces merely to provide a given number of grammes of vitamin seems the wrong policy. Surely the ultimate aim must be the manufacture and storage of the vitamins or vitamin concentrates. Meanwhile the provision of green vegetables out of season seems to leave a good deal to be desired in times of emergency.

The word "emergency" brings me to one final sentence. Although the statement I have just made is definitely a review of the position in the light of a possible emergency, the review has much wider implications. The possible implications have been put forward with all the authority of Maynard Keynes at the Cambridge meeting of the

British Association; it would be an impertinence for me to do more than allude to them. They are, however, in brief, that much work could be saved, prices could be stabilized, and speculation discouraged by a policy of storing foodstuffs which could be released in the correct quantities at the appropriate moments.

Summary

This paper deals with the contribution which food investigation has made to the facilities for the supply of commodities to this country, especially in time of emergency.

Foodstuffs are discussed under three main headings, according to the length of time they can be kept.

It is considered that the ultimate goal would be the manufacture and storage of the vitamins.

THE COMMONER TYPES OF SUPERSENSITIVENESS AND THEIR MANAGEMENT

BY

J. GIBSON GRAHAM, M.D., F.R.F.P.S.G.

Clinical Tutor and late Dispensary Physician, Western Infirmary, Glasgow

AND

J. D. OLAV KERR, M.B., Ch.B.

Clinical Tutor and Extra Dispensary Physician, Western Infirmary, Glasgow

Supersensitiveness has been defined as the liability of an individual to suffer severe effects of a special nature on exposure to certain agents which are harmless to others, at least in the doses used (Browning, 1934). The terms "allergy," "idiosyncrasy," and "hypersensitiveness" have been employed as equivalent terms. This condition may show itself in connexion with many of the body tissues—for example, the respiratory tract, the gastrointestinal tract, and the cutaneous structures, depending on the susceptibility of the individual. Thus one patient may display catarrhal manifestations of supersensitiveness affecting the mucous membranes of the nasopharyngeal region (hay fever), while in another, bronchial spasm occurs with resultant asthma. In others, varieties of cutaneous manifestations may be seen—for example, angioneurotic oedema, urticaria, dermatitis, etc.; also nausea and colic may be due to gastro-intestinal supersensitiveness.

Manifestations of Supersensitiveness

Asthma.—Owing to its wide and somewhat indiscriminate use "asthma" should be defined. Asthma, as distinct from the dyspnoea of simple bronchospasm, is a type of paroxysmal dyspnoea occurring in individuals in whom certain primary essentials are present. Bray (1937) has pointed out that these are, firstly, a family tendency to supersensitiveness, and, secondly, tissue damage following on a severe illness often pneumonic in nature, coupled in either case with a superimposed specific sensitization. In these individuals a paroxysm may be initiated by such secondary factors as reflex nasal irritation, psychic and endocrine disturbances, and non-specific factors often alimentary in origin. In the asthmatic attack there is a spasmodic constriction of the bronchial muscles