RENAL THRESHOLD FOR GLUCOSE : NORMAL AND IN DIABETICS

BY

R. D. LAWRENCE, M.D., F.R.C.P.

(From the Diabetic Clinic, King's College Hospital)

The renal threshold for glucose is of great interest to physiologists in relation to kidney function and of importance to clinicians in the study and treatment of glycosuria and diabetes. The term is applied to the critical concentration of sugar in the blood which determines the presence or absence of glycosuria. Below this level sugar is absent in the urine except in such minimal amounts as are always present and are not appreciated by the usual tests, Fehling or Benedict; while above this level easily appreciable amounts of sugar appear in the urine and gradually increase both in total amount and in percentage concentration as the blood sugar becomes progressively higher. The renal threshold is usually found to be between 170 and 200 mg. per 100 c.cm., depending rather on whether the analysis is made on venous or capillary blood. It is a common statement in medical literature that "the normal renal threshold is 170 mg. per 100 c.cm.," and a common implication that any deviation from this level is an abnormality, implying disorder of function if not disease of structure on the part of the kidney.

It has been recognized for forty years that deviations from this normal figure occur in both directions—that many individuals pass sugar in the urine when the blood sugar is below 170 mg. per 100 c.cm., and that many diabetics show no glycosuria when the blood sugar is considerably higher; in other words, that the kidney is more or less permeable to sugar than usual. Are these deviations to be considered as abnormalities or has the usual conception of the normal to be reconsidered?

Physiologists are not entirely agreed on the mechanisms of sugar excretion, but the commonest explanation is that glucose from the blood diffuses readily through the glomeruli, passes down the tubules in the glomerular filtrate, and is reabsorbed there to avoid the waste of a valuable substance. It is not thought probable that the tubular area actively secretes or excretes sugar into the urine, and so the presence or absence of sugar depends mainly on the reabsorptive power of the tubules. Whatever the detailed mechanism, it is clear from clinical observation that the kidneys are often more or less permeable to sugar than the accepted normal.

Determination of Renal Threshold

This is simple and consists of a *simultaneous* comparison of glycaemia and glycosuria, which is most satisfactorily carried out by the standard glucose-tolerance test where the blood sugar is estimated every half-hour for two hours after a dose of glucose. It is essential, if the threshold is to be ascertained with any accuracy, that the bladder should be emptied and separate samples of urine tested every half-hour, a point that is often neglected. Indeed, tests at more frequent intervals, say every fifteen minutes, are likely to catch the threshold more accurately, but it is impossible for many people to pass urine so often. Less elaborate and often equally clear is a single estimation of blood sugar and the testing of two samples of urine, the first passed immediately before (the bladder must be completely emptied) and the second as soon as possible after the blood sugar has been estimated. The *second* specimen

is therefore secreted at a known blood-sugar concentration and gives information on the threshold. For instance, in insulin-diabetic patients whom I suspect of a low threshold because the glycosuria persists even when the blood sugar has been reduced the blood is tested and the bladder emptied at the height of insulin activity (usually four hours after breakfast) and a second urine obtained as soon as possible and certainly before more food is taken. If this second specimen contains sugar and if the blood is below the normal threshold—say between 100 and 140 mg.—then a low threshold is clearly established. A point of possible error to note is that if the first voiding contains a heavy amount of sugar the second may contain traces without actual sugar being secreted from the kidneys, presumably through even a minimal "wetness" remaining in the bladder. I had this proved many years ago on a series of non-diabetic cystoscopy cases by filling the bladders with 5 per cent. glucose, draining this completely, and testing a subsequent sample passed naturally. About a quarter of these samples showed traces of sugar.

Permanent Deviations of Threshold

Variations of the threshold from the accepted normal are very common, but it is impossible to state in what percentage of the population they occur, as no extensive investigations have been undertaken.

A *low renal threshold* is often found in life assurance and other routine tests, and if the glycosuria is proved to accompany a normal blood-sugar concentration it can be regarded as a harmless anomaly consistent with perfect health and indicating neither disturbed carbohydrate metabolism nor renal disease. The condition is usually and best called *renal glycosuria*, but *diabète rénal* and *orthoglycaemic glycosuria* are other names for it. I have met it at all ages from 2½ to 80, and in my experience it generally persists throughout life. This renal anomaly has a strong hereditary tendency, and it runs in many families as a dominant Mendelian characteristic. The condition has been recognized for years, and the affected individuals are now accepted by most insurance companies as first-class lives.

Two points of error on the subject still occur in some textbooks. It is often stated that the degree of glycosuria is *independent of carbohydrate intake*. This in my experience is quite wrong. If the threshold is only moderately low, say 130 mg., then glycosuria appears only after a large carbohydrate meal. If the threshold is under 90 mg., sugar is also excreted even in the fasting condition, but more is excreted when the blood sugar rises after carbohydrate food or a glucose test. If more sugar is not excreted at the higher blood-sugar levels at the peak of a curve it is probable that the reducing substance is not glucose but lactose, laevulose, or pentose, all non-threshold substances which filter through the kidneys in constant small amounts. Again, it is frequently stated—indeed by Joslin (1937)—that a case of renal glycosuria should be watched and the diet be somewhat restricted for a few years before the conclusion is reached that the condition is harmless, although that author admits that he has never seen a case develop into true diabetes. I have seen two cases become diabetic, but this is not unnatural in over 300 cases discovered. If normoglycaemic glycosuria is proved by a thorough investigation, such as the finding of a normal blood-sugar curve, then I think the whole condition can be neglected at once and for ever. At any rate, I have never regretted acting on this opinion.

A *high renal threshold* has naturally been noted mostly in *diabetics*, because they alone have blood sugars which exceed the normal accepted renal threshold of 170 to 200 mg. per 100 c.cm. It is common to find that a diabetic becomes sugar-free on treatment and yet his blood sugar is found to be between 200 and 250 mg. or occasionally even higher. This occurs most frequently, I think, in elderly patients, but I have seen it in a few children and several young diabetics with healthy kidneys and showing normal renal function tests. A high renal threshold is very common in chronic sclerosing nephritis (Castaigne and Chaumerliac, 1939), and it is well recognized that the development of progressive sclerotic lesions may bring about the disappearance of glycosuria at a blood-sugar concentration which produced abundant glycosuria before the onset of kidney disease.

There is clear evidence that *non-diabetics* may also have an unusually high renal threshold. During investigations of the hyperglycaemia produced by anaesthesia it has been observed that the blood sugar may exceed 200 mg. without the appearance of glycosuria. That this is not the effect of anaesthesia on the threshold is shown by the fact that other cases do pass sugar during anaesthetic hyperglycaemia. Mann (1925) has shown that often the threshold is high in non-diabetic psychotics.

The frequency with which deviations of the threshold occur in the population is not known, as probably only a very small proportion of renal glycosurics are ever discovered, and persons with a high threshold never. From diabetics with normal kidney function we can get a rough idea of the percentage involved, from which we can draw conclusions, in my opinion fair ones, about the population at large. In a series of 500 diabetics studied up to 1929 (Lawrence, 1929) a lowered renal threshold was noticed in about 2 per cent. of cases and no doubt was missed in others; and in my experience a raised renal threshold is much commoner.

Deviations from the usual threshold lead to difficulties in the treatment of diabetes. A low threshold in diabetes has two disadvantages. If the condition is not recognized there is a tendency to raise the insulin too high in an attempt to abolish glycosuria, and serious hypoglycaemia may be produced in a vain attempt to free the urine from sugar. Secondly, the heavy loss of sugar with only slight degrees of hyperglycaemia wastes so much sugar from the carbohydrate of the diet that weight and energy may be lost and serious ketosis occur on a carbohydrate diet which would be adequate were the threshold normal. The lowering of the threshold in pregnancy—very common—is the chief cause in my experience of serious ketosis in diabetic pregnancies, and can only be overcome by a very high carbohydrate intake. A high renal threshold, on the other hand, generally results in a diabetic being allowed to live with a constant and often considerable hyperglycaemia, and in this type of case such complications as arterial disease, retinitis, and neuritis are unusually common. It is not certain, however, whether these complications are directly due to this hyperglycaemia or are merely common in elderly diabetics, who so often have a rather high threshold.

Changes in the Individual's Threshold

Although statements are made in medical writings that variations occur in the threshold from time to time, the evidence is not convincing, and my experience leads me to believe that low or high thresholds persist unchanged for many years if not for life. This is certainly true

of renal glycosuria in non-diabetics. The fact that the leak point is usually lower on the fall than on the rise of a blood-sugar curve does not invalidate this. Temporary changes, however, certainly occur in diabetics, and have happened in the following circumstances in my cases.

TEMPORARILY LOWERED THRESHOLD

Polyuria itself may produce this, perhaps by allowing insufficient time for reabsorption of glucose in the tubules. It is not uncommon to find a low threshold at the beginning of treatment in a patient with intense polyuria and after ten days the disappearance of glycosuria at a blood-sugar concentration practically the same as before. The most striking example I have seen occurred in a patient with only one kidney: in this case an initial blood sugar of 210 mg. per 100 c.cm. was accompanied by a total excretion of 212 grammes (8.5 per cent.) of sugar, whereas after a week of low carbohydrate diet only 2.5 grammes (0.2 per cent.) was excreted at a similar level in the blood, a threshold which has persisted for seven years and seems to be his permanent threshold.

Pregnancy often produces a temporary lowering of the threshold, and this was proved in twenty-nine (85 per cent.) out of thirty-four of my pregnant diabetics in whom the threshold was accurately determined. It is said to be equally common in normal women. The cause of this is obscure, but it is perhaps due to an effect of changed pituitary activity on their kidney function. It has long been known that phloridzin, which produces glycosuria by lowering the threshold, has more effect on pregnant than on non-pregnant women.

Posterior pituitary injections (pituirine) certainly lowered the threshold in one of my diabetic cases, and Mann states that he has repeatedly noted this effect of pituitrin in blood-sugar experiments on psychotics (personal communication).

French writers (Castaigne and Chaumerliac, 1939) state that subacute glomerular nephritis may produce a lowering of the threshold. I have not noticed this.

In one case of severe *toxaemia* from a huge carbuncle I observed a temporary lowering of the threshold to under 100 mg. per 100 c.cm. and a rise to its usual level on recovery.

TEMPORARILY RAISED THRESHOLD

This may occur: (1) in *circulatory failure* with oliguria; (2) in *diabetic coma* with its curious kidney "block" in which the blood contains a large excess of sugar and ketone bodies, but these may not appear in the urine and even sodium chloride may be retained; and (3) in *nephrosclerosis* (ischaemic contracture), which gradually makes the kidney less permeable to sugar and the threshold progressively higher.

Summary and Conclusion

The study of the renal threshold in diabetics and normals with healthy renal function shows frequent upward and downward deviations from the accepted "normal" figure of some 170 mg. per 100 c.cm.

The conception of a "normal" threshold is false. There is an *average* renal threshold, just as there is an average blood pressure, and the many deviations from it should be looked upon as physiological—a matter of individual idiosyncrasy, and of no pathological significance.

High and low thresholds are frequent and usually persist at the same level throughout life. A few factors

outlined above affect the threshold temporarily or even permanently.

The stigma of something abnormal (and hence pathological) that has been imputed to a high or low renal threshold should be removed.

REFERENCES

- Castaigne, J., and Chaumerliac, J. (1939). *Nutrition*, 9, 19.
 Joslin, E. P. (1937). *Treatment of Diabetes Mellitus*, p. 644, Philadelphia.
 Lawrence, R. D. (1929). *British Medical Journal*, 1, 196.
 Mann, S. A. (1925). *J. ment. Sci.*, 71, 443.

PREVENTION OF THROAT INFECTION

BY

HELEN PIRELL GOODRICH, M.A., D.Sc.

AND

GEOFFREY L. WAY, M.R.C.S., L.R.C.P.

Lieutenant R.A.M.C.

The toll in time and fitness levied by repeated epidemics of colds, sore throats, and inflamed tonsils arising among war workers is well known. When people are crowded together "droplet" infection is of course especially liable to occur.

Thymol as a Mouth-wash

During the last war, after investigating the action of various antiseptics on the organisms occurring in the mouth and tonsils, it was suggested by one of us (Goodrich, 1917a) that a solution of thymol was an excellent mouth-wash. This opinion has since been confirmed not only by our own daily experience but by that of many others. A distinguished biologist told us recently that he had used a saturated solution of thymol as a gargle as well as a mouth-wash ever since 1917, and to this attributed the fact that he had since had no throat infection, not even a common cold. Many people prefer to dilute the solution. No doubt there is considerable individuality with regard to the taste of thymol, some people finding it rather peppery and inclined to be irritating if strong. It should be remembered, too, that the solubility rises rapidly with the temperature, so that a really cold solution (0.05 to 0.06 per cent.) is much weaker than one saturated in a warm room.

For popular use the solution is best made by keeping the thymol in a suitable bottle filled with water; if in small crystals these may be enclosed in muslin to prevent their escape when the solution is poured off. The bottle may be refilled with cold* water as the solution is used, and one can rely on having a saturated solution after a short time. When required as a mouth-wash or gargle it may be added to water (preferably warm) until the taste is right for the individual but not strong enough to "bite the tongue," as children say. The thymol is so slightly soluble that an ounce, costing a few pence, will make about nine gallons of solution; the price is therefore negligible. The saturated solution is also very useful for disinfecting rooms and articles in them, and especially for spraying windows and mosquito nets in the Tropics.

It has elsewhere been pointed out (Goodrich, 1917b) that although the addition of glycerin enables much more thymol to dissolve in water there is no corresponding increase of antiseptic power, and that therefore the use of glycerin in this way is a great waste of a substance required for munitions. This criticism does not apply to the use

of glycerin as a demulcent in throat paints, but only to mixtures intended to be diluted before being used, in which case, therefore, the glycerin has practically no action except the reduction of the antiseptic properties of the thymol.

The Case for Oral Hygiene

It may perhaps be well to repeat here that in every mouth left undisturbed for some hours—for example, during sleep—a varying amount of white substance grows on the tongue and other surfaces and accumulates round the teeth. This consists chiefly of the thread-like plant *Leptothrix*, in the meshes of which may be numerous micrococci and other organisms and cells. The leptothrix, if not removed, will grow rapidly and calcify, passing from the so-called soft tartar ("materia alba" of Leeuwenhoek) to the well-known hard tartar. Goodrich and Moseley (1916) also recorded how frequently organisms from the mouth are to be found growing in the crypts or pockets of the tonsils. These folds of epithelium, when filled with calcified colonies of leptothrix as well as numerous bacteria, show up as white spots on the surface of the tonsil (so-called ulcerated throat). This is the unhealthy condition that must be cleaned up: if neglected the whole tonsil may become much hypertrophied, whitish, and thoroughly septic, with organisms and pus oozing from the surface as in some of the excised tonsils we have examined in the laboratory.

All this points to the advisability of keeping the throat and tonsils, as well as the mouth, clean. It seems strange that some people who would never think of missing their daily bath resent giving, by way of oral hygiene, more than a perfunctory rub to their teeth with a brush (possibly teeming with bacteria); and yet with a suitable antiseptic at hand it is so easy to sterilize a soft brush and to kill and remove most of the organisms trying to establish themselves in the mouth or throat. It would seem that the mere mechanical action in gargling must be of use in helping to empty the tonsil pockets so that the antiseptic can wash away the contents. Most children will take to gargling, as to the use of a toothbrush, if they have an inkling that their parents and other grown-ups use these hygienic measures, which really take so little time. Cleanliness can, of course, be gauged by the absence of tartar from the teeth and the healthy appearance of tongue and tonsils. Usually nothing more will be required to ensure reasonable freedom from infection.

One has, however, if necessary, a further line of defence in the practice of painting the throat and tonsils: here again the mechanical action doubtless helps to remove organisms lodged in the pockets or multiplying on the mucous surface of the throat. In the standard iodine-and-glycerin mixture used as a paint one can well afford to have the powerful antiseptic properties of iodine reduced (see below), and the glycerin ensures that the organisms are submitted to the antiseptic for a longer time than in mere rinsing.

In the report published by the Ministry of Health (Straker, Hill, and Lovell, 1939) on the bacteria of the nasopharynx surprisingly high percentages of meningococci, haemolytic streptococci, and other pathogenic organisms are recorded from some groups of the population. It is very much to be hoped that these investigations will be continued and some correlation attempted with the general hygienic conditions of the mouths and throats tested. Perhaps the results would throw some light on a fact for which the authors seem unable to account at present—namely, that in groups with similar occupations consider-

* Hot water must not be added to the solid thymol, since it melts at 51.5° C.