except the sterno-mastoid. No other physical abnormalities were present.

The symptom came on gradually three years previously while she was engaged. She first felt she could not face people and must look away. This difficulty became gradually worse until her head was turned to the right all the time, making her still more sensitive about meeting people. She married, thinking this would help her to get better, but was unable to go away on her honeymoon, so she and her husband have continued to live with her parents. She had a baby, and again hoped this would help her. From the fourth month of pregnancy until the child was 3 months old she had considerable relief from symptoms, but later they returned worse than before. She looked after the child and helped in the home, but was too self-conscious to go out alone, and would only venture out a short distance accompanied by her mother.

Previous History.—She was an only child and had been brought up very strictly; no sexual instruction was given nor was the subject ever referred to. At the age of 4 she went into her mother's bedroom and saw her mother standing naked. She was horrified at the sight of her mother's pubic hair, thinking it was some terrible abnormality, and rushed from the room feeling that she had seen something she ought not to have seen. The incident was completely repressed, but from then on she became very particular to keep her body entirely covered. At the age of 12 to 13 she noticed her own pubic hair developing. She was afraid of this, but did not know why. She felt that she was different from other girls, but dared neither to ask anyone nor to try and find out if her fears were justified. She kept it all to herself as a "terrible secret," but led a fairly normal life. Analysis showed a repressed fear that she had inherited a horrible abnormality from her mother. When she became engaged she was very anxious because she thought that when married her secret would be discovered. She became more conscious of a difference from other women and gradually began to feel she could not face other people. Her head seemed to be pulled to the right, and as the wryneck became more established her anxiety diminished. After marriage she was completely frigid and was terrified by intercourse: frequently an acute spasm of her neck would come on and prevent it. Her temporary improvement coincided with the period when coitus was impossible because of pregnancy.

Results and Comments.—Analysis produced freedom from symptoms. She became able to go out alone, was free from embarrassment in company, and her sexual life became normal. Her symptom symbolized her attitude of turning away from her sexual role as a woman.

Conclusions

It is a most striking feature that, although the psychogenic origin of spasmodic torticollis has so long been recognized, there was such an extended history of the condition in these cases before they were referred for psychotherapy. In three of them it had been present for from two to five years. Much inconvenience had been entailed and much valuable time wasted by previous methods of treatment, which in two cases had been prolonged in spite of the patient showing no response. As long ago as 1907 Janet wrote in reference to hysteria: "Do not count the number of arms cut off, of muscles of the neck incised for cricks, of bones broken for mere cramps, of bellies cut open for phantom tumours, and especially of women made barren for pretended ovarian tumours." It is regrettable that even now psychotherapy should have been considered for these patients only as a last resort, instead of being advised as soon as they came under medical observation. Belief in physical instead of psychotherapeutic lines of treatment for conditions in which the psychogenic origin is well recognized dies hard.

In three of these cases there was a period of mild generalized anxiety, which came on first but subsided

when the torticollis became established. The symbolic nature of the symptom seems to be significant. The patients each had some factor which they felt they could not face or from which they felt compelled to turn away. This bears out the importance Halliday (1937) gives to the symbolic significance of symptoms in his studies of the psychological factors in rheumatism. The more carefully we study the cases the more we see that the symptom is not an isolated factor but the response of an inadequately adjusted personality to a period of particular stress. Our task is to discover what real satisfaction lies behind the symptom. Our efforts to correct the condition must therefore be directed not to the torticollis alone, nor simply to a disclosure of its genesis, but to a fundamental attempt to readjust the personality to life.

Summary

The literature of spasmodic torticollis is reviewed and the psychogenic factors are stressed.

Four cases treated by psychotherapy are recorded; three of these recovered.

The importance of early psychological investigation is demonstrated.

I wish to thank Dr. T. Wishart Davidson, medical superintendent of the City Mental Hospital, Leicester, for permission to publish the cases.

REFERENCES

- Brain, W. R. (1928). *Lancet*, **1**, 1295.
 Clark, L. P. (1912). *J. Amer. med. Ass.*, **58**, 838.
 — (1914). *Med. Rec.*, **85**, 232, 371, 553.
 — (1917). *Ibid.*, **92**, 48.
 Finney, J. M. T., and Hughson, W. (1925). *Ann. Surg.*, **81**, 255.
 Frazier, C. H. (1930). *Ibid.*, **91**, 848.
 Halliday, J. L. (1937). *British Medical Journal*, **1**, 213, 264.
 Janet, P. (1907). *Major Symptoms of Hysteria*, New York.
 Meigs, H., and Feindel, E. (1907). *Tics and their Treatment*, London.
 Schaller, W. F. (1913). *J. Amer. med. Ass.*, **60**, 1421.
 Yaskin, J. C. (1935). *J. nerv. ment. Dis.*, **81**, 299.

MULTIPLE EMBOLI TREATED SURGICALLY

BY

J. A. MacFARLANE, F.R.C.S.Ed.

Lieut.-Colonel, R.C.A.M.C.

(From the Department of Surgery, University of Toronto)

The following case is reported because it is felt that some of the principles illustrated in the treatment may well be applicable to the surgery of the vascular system in war.

The patient, a young unmarried woman of 27, was admitted to the surgical wards of the Toronto General Hospital on April 12, 1939. Her physician had seen her a few hours before and thought that she was suffering from acute appendicitis. She had wakened the previous evening with abdominal pains and vomiting. The pain seemed to be localized in the right lower quadrant. When she was seen on admission she was somewhat cyanosed and her heart was fibrillating at 170. It was then learned that she had had chorea at the age of 12 and that her doctor had been giving her digitalis at intervals for three years. It was obvious that she had mitral stenosis, and that this was the cause of the fibrillation. When her abdomen was examined she was found to have definite tenderness in the right iliac fossa, but no rigidity. The rectal examination was negative: there was no blood. The leucocyte count was 18,500. Her urinary examination was negative and her temperature 99.6°. It was felt that she might have a small mesenteric occlusion from an embolus, so she was watched carefully. Her pain and nausea subsided in twenty-four hours and she felt much better. Digitalis therapy was initiated.

On April 17, five days later, when the weekly ward round was in progress, the nurse called attention to this patient, who

a few minutes before had complained of a numb feeling in the left leg, accompanied by fairly sharp pain. In a very short time there was loss of sensation to the knee, pallor, and loss of position sense. Pulsation could not be elicited at any point below the termination of the common femoral artery. This whole picture did not develop immediately. Only after several careful examinations was it definitely decided to operate on the femoral artery in Scarpa's triangle. The procedure was carried out under spinal anaesthesia. An embolus was found riding on the bifurcation of the common femoral and the profunda femoris. A solution of heparin was used to flush out the artery at the operation site both before and after the closure. There was marked spasm of the artery below the line of suture, but on removing the tapes it could be seen to pulsate below the bifurcation. The colour of the foot improved immediately. Sensation was normal by the time the patient had recovered from the anaesthetic. A continuous intravenous solution of heparin, in normal saline, was given under the supervision of Dr. Ross MacKenzie and a careful check was kept on the clotting time. The heparin was discontinued on April 25.

In the morning of April 26 she began to have numbness in her right foot and had difficulty in moving it. It gradually became cold and cyanosed. No pulsation could be felt in the right femoral artery or any vessel below it. She was again subjected to operation under spinal anaesthesia, with a diagnosis of common iliac occlusion. The exposure was through the right rectus behind the peritoneum of the iliac fossa. Again a large embolus was removed and heparin was used locally and intravenously. Again colour and sensation returned.

On May 3 she was feeling quite well and the heparin was discontinued. On May 4 she had sudden lower-abdomen pain, with some cramps and pain in the region of her sacrum and finally pain down both thighs. She showed some shock. Both legs became cold and cyanosed and her pain increased. Her general condition appeared more grave than on the previous occasions. No pulsation could be felt below the bifurcation of the aorta. It was with considerable misgiving and a feeling of discouragement that further operative treatment was contemplated. Again under spinal anaesthesia the abdominal aorta was exposed at its bifurcation as well as the common iliac vessels on both sides. This was done through the old right rectus incision prolonged upwards. The site of the previous exposure of the common iliac bifurcation was examined and the incision in the external iliac artery was well healed. A large embolus was found at the bifurcation of the aorta. It is not easy to prevent part of such an embolus from slipping one way or the other. The major portion of the clot was removed through an incision in the right common iliac artery.

After this operation colour and sensation immediately returned on the right side but were slow in returning to the left side. Heparin was instituted again. On May 7, three days later, she developed all the signs of another embolus in the right femoral artery! One began to wonder whether the whole thing was futile; but after further serious consideration, and with the advice of Dr. Gallie and Dr. Gordon Murray, the artery was again exposed—this time the right femoral bifurcation—and once again a good-sized clot was removed. On this occasion it was felt that a small piece had got away into the femoral artery.

Following this procedure the patient regained colour and sensation in her leg but had pain and weakness in the posterior and lateral groups of leg muscles. I believe this indicated an embolus in the posterior tibial artery. The leg was put in plaster, and the posterior group of muscles regained their power within three weeks. The peronei remained weak. Heparin was discontinued on May 28 and she was allowed up. On June 6 she complained of sudden headache and nausea. On examination she had a right facial weakness, paralysis of the right side of her tongue, and a cold blue left arm. Fortunately all these symptoms began to improve within four hours. By the following day there were no cerebral symptoms and the arm was again warm. Sensation had returned and she could

move it, although there was no return of the radial pulse. She was allowed up again the following day, and within a week was discharged to a convalescent home. Here she rapidly regained her strength and was finally sent home on June 28. She was advised to continue with the digitalis therapy, which had been maintained throughout her stay in hospital.

Seen three months ago she looked and felt well. She was still fibrillating, but had no failure and has had no further vascular accidents.

Comment

This case serves to illustrate the tremendous advantages of heparin in blood-vessel surgery. Dr. Gordon Murray, who delivered the Hunterian Lecture on this subject last June, saw the patient at the time of the second accident and assisted with two of the operations. Dr. Ross MacKenzie, who has been associated with Dr. Murray in this work, supervised the administration of heparin during and following each operation. Before the use of heparin embolectomy was so frequently followed by thrombosis that it came to be regarded as a procedure of questionable value. Now one can carry out surgical procedures on arteries with the assurance that thrombosis will not occur and that normal healing of the artery will take place.

CYANOSIS AND ANOXAEMIA IN GAS-AND-OXYGEN ANAESTHESIA

BY

R. L. WYNNE, M.B., M.R.C.S., D.A.

Anaesthetist, Birkenhead General Hospital

The justifiable objection to cyanosis occurring in the course of anaesthesia by ether and the other lipid-soluble agents has in the past tended to lead to an unjustifiable assumption that there is something about mere blueness in a patient which is wrong, irrespective of its cause; and, since with most gas-and-oxygen anaesthetics cyanosis is an almost invariable concomitant of good anaesthesia in the robust patient, the assumption has in turn led to inefficient anaesthesia and backwardness in the use of a valuable agent, and discouragement of its use by those not perfectly familiar with the distinction between the different causes of cyanosis and between cyanosis itself and anoxaemia.

Cyanosis occurring almost "physiologically" is seen in the congenital heart patient. Here one may be presented with the picture of a person who spends his whole life fairly comfortably carrying out a reduced scale of exertion with a permanently blue complexion, but whose physiological make-up has become so modified that in his own metabolic range he is no more anoxaemic than his perfectly pink relatives with normal hearts.

Cyanosis begins in any person when his circulating blood contains, not any definite *ratio* of reduced to oxygenated haemoglobin, but more than a *fixed amount* of 5 grammes per cent., whatever his total haemoglobin resources may be. Thus, in the congenital cardiac, we may be sure from his blueness that his blood contains considerably more than 5 grammes per cent. of reduced and useless haemoglobin, but we may also be sure from his absence of distress in his permitted range of exertion that besides the 5 grammes per cent. of useless haemoglobin there is side by side with it a sufficient number of grammes per cent. of active haemoglobin to carry all the oxygen he is accustomed to require. In other words, he is cyanosed but not anoxaemic.