BRITISH MEDICAL JOURNAL

LONDON SATURDAY AUGUST 5 1939

THE PRESENT POSITION OF CHEMOTHERAPY BY DRUGS OF THE SULPHANILAMIDE GROUP

THE BACTERIOLOGICAL AND EXPERIMENTAL ASPECT*

BY

C. H. BROWNING, M.D., F.R.S.

(From the Bacteriology and Pathology Department of the University and Western Infirmary, Glasgow)

The cure of experimental streptococcus infections by drugs of the sulphanilamide group marks an epoch in the progress of chemotherapy begun by Ehrlich and Shiga in 1904. It is not surprising that the initiative in this work has always come from the laboratory, since it is only by the use of small animals that the essential desideratum of large numbers of infected individuals showing a relatively uniform and progressively fatal disease can readily be secured. In them the assessment of therapy is conclusive, death or survival being the criterion. A great difficulty, of course, is the transfer of results from one animal species to another or to man, as different hosts may vary either qualitatively or quantitatively in their behaviour both to the pathogenic agents and to the drugs. Nevertheless, the essential basis for the clinical use of compounds of the sulphanilamide class must continue to be their performance under experimental conditions; hence it is fortunate that observations with bacterial infections in mice should have proved so accurate a guide to the successful treatment of corresponding conditions in man. In giving an account of the subject I shall try to select from the voluminous literature what presents a fair general perspective, and shall speak in the light of unpublished work done jointly with Dr. J. V. M. Robb† and Miss M. W. Leckie, B.Sc., during the past three years.

Historical

Morgenroth and Levy in 1911 cured a general infection with pneumococci in mice by means of optoquine, although apparently results were uncertain and the drug was rather toxic for use in man. Before Domagk reported the discovery that the azo "prontosil" dyes of Mietzsch (1938) and Klarer could cure a general infection with streptococci, it was known that by applying suitable antiseptics early enough to the site of inoculation the progress of an infection due to these organisms could be prevented. It must be emphasized, however, that the general infection caused by intraperitoneal inoculation with streptococci was controlled by injecting prontosil subcutaneously. Thus it is not necessary for this drug, as it were, to chase up the organisms: cure occurs when the two meet in the tissues, having come from opposite direc-

tions. The Tréfouëls, with Nitti and Bovet, then simplified remarkably the position as regards the chemical constitution of effective drugs by showing that one moiety of prontosil—namely, the long-known compound *p*-aminobenzenesulphonamide (sulphanilamide)—also has a curative action. This substance, which is colourless, water-soluble, and comparatively little toxic for mammals, contains sulphur in the form of a sulphonamide group (SO_2NH_2). It is not known whether the sulphur atom is indispensable for the chemotherapeutic property, but, so far, no compound showing activity comparable to that of sulphanilamide has been described which does not contain sulphur in some form. Therapeutically active substances exist, however, such as the diphenyl sulphones, which it is difficult to conceive as being converted into sulphanilamide or yielding identical derivatives *in vivo*.

Sulphanilamide and Streptococci in Mice

The test infection is with a culture of streptococci highly virulent for mice: such streptococci are rare; only one was obtained among over fifty strains from human lesions.‡

The intraperitoneal inoculum consists of an overwhelming dose which in untreated animals causes septicaemia and death within twenty-four hours. The drug may either be injected subcutaneously or introduced into the stomach. In order so far as possible to secure uniform and uncomplicated conditions we have adopted the subcutaneous method, and a spacing of doses similar to that originally used by Buttle *et al.* (1936)—an injection at the rate of 0.01 gramme for a 20-gramme mouse being given at the time of inoculation, followed by a second of 0.0066 gramme four to five hours later, and a third dose of the same amount after twenty-four hours; animals of 19 to 25 grammes weight were used.

With an inoculum consisting of about one hundred times the number of organisms certainly fatal within twenty-four hours, this treatment is by no means surely curative. Death often occurs from septicaemia after an interval of three or four days, or longer. Again, some animals which appear to have been cured die after many weeks from streptococcal septicaemia. The tendency to relapse which is so marked in infections with

‡ This strain, from a carbuncle of the neck, also a culture for which we are indebted to Dr. Adolf Feldt, Berlin, have been chiefly used for our experimental work. The former is slightly more virulent for mice; but the therapeutic results are similar with both.

* Read in opening a joint discussion at the Annual Meeting of the British Medical Association, July, 1939.

† Working as a Muirhead Scholar and with the support of the Medical Research Council.

protozoa and spirochaetes has also been noted by others in streptococcus infections treated by sulphanilamide, and it was recorded by Bordet in rabbits infected with streptococci and treated by antiserum. Buttle *et al.* (1936) found that several different serological types of streptococci were influenced.

Similar treatment with sulphanilamide of an infection with pneumococci in mice, say of Type I—of the same severity as that with the streptococcus, so far as can be judged by the time taken to produce a fatal septicaemia after inoculation—leads to a much smaller proportion of cures. We attempted to better the results by continuing administration of 0.0066-gramme doses once daily; but in spite of this the animals succumbed to septicaemia in several days during the treatment. Accordingly infection with the pneumococcus is less amenable than that with haemolytic streptococci: this is general experience.

Drugs More Effective than Sulphanilamide

In the hope of improvement the abundant possibilities presented by synthetic derivatives of sulphanilamide and related compounds have been widely examined by Buttle, Fournau, and others. It has been found with certain of these substances that streptococcus infection in mice is influenced by a smaller amount, being also a smaller fraction of the tolerated dose as compared with sulphanilamide (this dose, however, may be difficult to fix with the relatively insoluble compounds, whose low toxicity may be chiefly due to slow absorption). Further, their action on pneumococcus infections is more marked. Among these may be mentioned *p-p*-diamino-diphenyl sulphone (Buttle *et al.*, 1937b), rodilone (*p-p*-diacetyl-diamino-diphenyl sulphone) tested by Fournau *et al.*, corresponding monoacyl derivatives—for example, the monoacetyl compound*—and Whitby's (1938) "May & Baker No. 693" (M & B 693) (a sulphanilamide derivative introduced on account of its activity against pneumococcus infection). As tested by subcutaneous injection in mice infected with streptococci, we have found monoacetyl *p-p*-diamino-diphenyl sulphone to be the most effective of the last three. It has shown a similar behaviour with a Type I pneumococcus. The mere statement of these results, however, leaves many points unsettled. For instance, do the more efficient compounds simply persist longer in the body and so maintain, as by a depot, a more continuous action than sulphanilamide, which is very rapidly excreted, or, other things being equal, are they (or still more active derivatives formed *in vivo*—although there is no experimental proof of such) actually more potent? At present it is not possible to give an answer, since there are wide gaps in knowledge. Again, the distribution of sulphanilamide in the body and secretions is traced by means of colour reactions given by the amido side chain, but nothing is known of the behaviour of the sulphonamide group.

We examined the effect of varying the time between the injection of the drug and subsequent inoculation with streptococci. Conceivably such an experiment might yield results pointing to an optimum interval, which would indicate that the streptococci were most efficiently dealt with either when they met a maximum concentration of the drug in the tissues (the maximum in the blood occurring in mice one hour after either a subcutaneous injection or administration per os of sulphanilamide—Feinstone *et al.* (1938); Marshall *et al.* (1938)) or when enough time had elapsed for some alteration to occur in the circulating drug or in the tissues. Accordingly a single subcutaneous dose of sulphanilamide, rodilone, or the monoacetyl derivative was given approximately five hours,

three and a half hours, one hour before, or simultaneously with, an intraperitoneal injection of streptococci. Untreated animals infected with one-hundredth of the inoculum used in the therapeutic experiments died from septicaemia in twenty-four hours. Under these conditions sulphanilamide merely delayed death for a day or thereby. With the two other compounds, especially the monoacetyl derivative, a considerable proportion of cures resulted, but no marked difference was observed according to the above-mentioned intervals before inoculation at which the drugs were injected. It should be noted that these two substances, being little soluble, are administered as a fine suspension and are relatively slowly absorbed.

In the case of drugs which act well when injected in a finely divided state, solubilized derivatives in similar doses have been found less effective. Thus, different compounds of the series vary in their therapeutic efficacy, and this depends, in part at least, on their solubility; and other physical properties such as diffusibility probably have an influence. Whether differences in chemical constitution in the series influence therapeutic action in other ways is uncertain, but this seems likely.

Influence of the Individuality of the Host

It must be strongly emphasized, however, that the individual animal is a most important factor in modifying results; this is true even when, as Buttle *et al.* (1936) rightly insist, all are infected at one time with the same culture. Both with sulphanilamide and other members of the group full dosage has failed to effect cure in certain mice, whereas among those which received only a sixth or a tenth some were cured, although all had been similarly inoculated and otherwise treated. Also, animals inoculated with a certain number of streptococci and receiving a given dosage of drug may not respond better than those which have been infected with thirty or a hundred times as many organisms but similarly treated. These unavoidable irregularities have rendered much quantitative work inconclusive.

Effect of Sulphanilamide Drugs on Various Infections in Mice and in Other Species

Successful results have been recorded by various workers in the treatment of other experimental infections in mice—for example, with meningococci (Buttle *et al.*, 1936; Proom, 1937), certain salmonella and coliform bacilli (Buttle *et al.*, 1937a; Mayer, 1938); but not others (Gross *et al.*, 1938b). These organisms tend not to be highly virulent, and the device has been employed of introducing the inoculum along with mucin into the peritoneum. Staphylococcus infection, which for its establishment in the mouse usually requires large inocula along with mucin, has not proved highly susceptible to the sulphanilamide drugs. In controlled experiments, however, with the compounds sulphanilamide, uleron, M & B 693, rodilone, and the monoacetyl derivative, the last three were the most effective against staphylococci† (P. Browning, unpublished). It was also found, similar to the observation of Buttle (1937), that infection may persist in the track of the inoculating needle and lead to local suppuration, with death of the animal from general infection some weeks after treatment. *B. welchii* infection responds well according to Long and Bliss (1937a), provided the inoculum does not contain too much exotoxin; Natvig (1938) met with failure, however. Infection with pasteurella organisms was little influenced (Buttle *et al.*, 1937a); but Schütze obtained favourable results in rats and

* We are indebted to Imperial Chemical Industries, Ltd., for some of these compounds.

† We are indebted to the Wellcome Laboratories for the culture used.

mice infected with *B. pestis*, especially with M & B 693. *Treponema pallidum* in rabbits was not affected (Campbell, 1937).

In other species of hosts also success has attended treatment—for example, in general streptococcus infection of rabbits and guinea-pigs; streptococcal pleurisy in rabbits, by Gay and Clark (1937) (but poor results were obtained by Nicholson (1938), who, however, began treatment relatively late and used small dosage); intradermal infection of rabbits with pneumococci (Kreidler, 1937); and spontaneous “sniffles” pneumonia in rats (Maier, 1938.) Certain virus diseases have responded to the sulphanilamide drugs—experimental meningitis due to the virus of lymphogranuloma inguinale in mice, also distemper in dogs. In the latter it is not clear whether the virus itself is influenced or the associated bacterial infection. In general, virus diseases have proved resistant (McKinley *et al.*, 1939). Reports have been contradictory on the effects in experimental tuberculosis in guinea-pigs (Buttle and Parish, 1938). For the sterilization of brucella infection in guinea-pigs doses of sulphanilamide approaching the toxic limit were needed (Wilson and Maier, 1939). Much remains to be done, however, on dosage and on the response of various infections in different hosts to drugs of the series other than prontosil and sulphanilamide.* There is general agreement that little or no effect is exerted on diseases due to exotoxins—for example, of *B. diphtheriae*, *B. tetani*, *V. septique*, *B. histolyticus*, staphylococcus, etc. (Levaditi and Vaisman, 1938; Gross, Cooper, and Lewis, 1938)—or on the action of such exotoxins as can be demonstrated *in vitro*—for example, streptococcus haemolysin and fibrinolysin (Huntington, 1938; Gross *et al.*, 1938a), or on the formation of streptococcus erythrogenic toxin (Huntington, 1938) or leucocidin (Gay *et al.*, 1939) *in vitro*.

The infections which have proved tractable in most cases are of the acute generalizing or diffuse type, when treatment is begun early after inoculation. Where focal lesions have been a feature, especially when of some duration, treatment has often been less effective—for example, streptococcal lymphadenitis of guinea-pigs (Seastone, 1937; Habs and Bader, 1938) and intradermal streptococcus infection of rabbits (Kolmer *et al.*, 1938); also in naturally occurring mastitis in cows (Hoge *et al.*, 1939). In mice, too, this has been noted: Berger (1937) injected streptococci subcutaneously and began prontosil treatment by mouth, either simultaneously or five hours later; while the occurrence of septicaemia was regularly prevented, local lesions tended to develop or spread. Clinical observations have corresponded—for example, those of Lockwood *et al.* (1938) in surgical conditions and Longcope (1938) in tonsillitis.

Other Factors that may Limit the Efficacy of Treatment

Dosage of the inoculum, other things being equal, is important; with a given virulent culture of streptococci a small inoculum is more easily overcome by the drug than a large one. The statement has been made, however, that infections due to highly virulent streptococci are more tractable than those caused by strains of less virulence. But it is by no means clear that virulence of the causal organism *per se*, as measured by the size of the certainly fatal inoculum, determines the responsiveness of the infection. Further, apart from the suggestive experiment of Maclean *et al.* (1939) with the pneumococcus, it has not been proved that unsuccessful treatment may lead in a

virulent strain of organisms to the development of marked resistance to the therapeutic agent, similar to the drug-fastness of trypanosomes. The evidence is not convincing in favour of specificity, in the sense of one drug acting better on infection with one species of organism while another compound is more effective against another, although Whitby (1937) has recorded observations in this sense with streptococci and meningococci.*

Mode of Action of Sulphanilamide Drugs

Attempts to explain the mode of action of the sulphanilamide drugs have not led to concordant results, but the problem is clearly one of wide importance in view of the variety of infections which are influenced. The chemotherapeutic property of prontosil could not have been discovered except by trial in infected subjects, because the drug lacks antiseptic action in the test-tube. But in the case of sulphanilamide certain observers noted that *in vitro*, provided the number of organisms used as the inoculum was small enough, streptococci were inhibited or even killed by considerable dilutions of the drug in ordinary media, human blood (with leucocytes present or removed), or serum (Colebrook *et al.*, 1936; Buttle, 1937; Hoare, 1938; Britton, 1938). This influence exerted by the size of the inoculum is manifest, however, with various other antiseptics—for example, mercuric chloride or acriflavine (Garrod, 1935; personal observations). The anti-bacterial effect has been held to explain the therapeutic action of the sulphanilamide group. As Levaditi and Vaisman point out, however, far more powerfully antiseptic substances exist which cannot cure a general streptococcal infection. For example, acriflavine, or especially proflavine, is little toxic for the mammalian body while at the same time an enormously more potent antiseptic than sulphanilamide, and its rate of excretion is not significantly different from that of the latter; nevertheless it fails to cure a general infection. Colebrook *et al.* (1936) remark: “It is somewhat surprising that the low bactericidal power of the blood of mice treated with the sulphonamide should be associated with such a notable therapeutic effect.” Therefore it must be concluded that the weak anti-bacterial property demonstrable *in vitro* is not likely to be more than a subsidiary factor contributing to the chemotherapeutic action of the sulphanilamide group. We have not detected any noteworthy diminution in virulence when streptococci were grown in ox serum (heated at 56° C.) plus a concentration of sulphanilamide as high as 1 in 2,000. Further, no opsonic effect of the drug has been found when in the usual phagocytic experiments it was added in concentrations of 1 in 5,000 to 1 in 40,000 to a mixture of heated serum, organisms, and leucocytes, although this has been claimed (Finkelstein and Birkeland, 1938). Levaditi and Vaisman concluded that endotoxins of various organisms, although antagonized *in vivo*, were not destroyed *in vitro* by the addition of sulphanilamide compounds. Gay *et al.* (1939) have made the striking observation that sulphanilamide is not fixed *in vitro* by large masses of streptococci; this contrasts with what occurs in the case of bacteria or protozoa and other compounds which are lethal for the organisms *in vitro* or *in vivo*. Accordingly, test-tube experiments do not, so far, supply data which enable the curative action of sulphanilamide drugs to be assessed.

The problem remains as to whether, when an infection is cured by these drugs, the organisms are acted on or the host. In the first place, sulphanilamide itself does not set up an anti-bacterial condition of any permanence within the body, for when, subsequent to its repeated adminis-

* The promising field for the combined use of chemotherapeutic agents and antisera is not dealt with here.

tration, inoculation with the organisms is made after an interval of some hours following the last dose, the progress of the infection is less influenced than when similar dosage is given at and after inoculation. Thus a necessary condition for therapeutic action is that the organisms and the drug (or perhaps the active derivative of the latter) shall actually meet in the body of the host. According to the observations of McIntosh and Whitby (1939) sulphanilamide drugs do not stimulate leucocytosis, neither do they influence the quality, quantity, or rate of formation of specific antibodies to pneumococci. Long and Bliss (1937b) and Mellon *et al.* (1938), in order to examine the events following treatment, inoculated mice intraperitoneally and, after allowing time for the organisms to proliferate, then administered the drug. A difficulty in this form of experiment is to influence the infection sufficiently. We proceeded as follows:

Mice were saturated so far as possible by adding rodilone in very fine suspension to their food. Animals which ate freely for a week were inoculated intraperitoneally with 0.25 c.cm. of a 1 in 10 dilution of a virulent twenty-four-hours broth culture of streptococci—over a million times the certainly fatal dose; at the same time a subcutaneous injection of sulphanilamide was given. The inoculated mice were examined at intervals, films and cultures being made from the peritoneal contents.

We found in confirmation of Mellon that no dramatic occurrence took place in the peritoneum; certainly phagocytosis, though seen, was not a striking feature.

Films of peritoneal contents showed the usual appearances which in human material characterize an acute inflammatory exudate proceeding to resolution—chiefly polymorphs at first, with the presence of large mononuclears in greater numbers later. But in uninoculated animals similar films, made by pressing a microscope slide against the visceral peritoneum, showed both types of cells; also in the infected animals the exudate was scanty. Accordingly, no very great abnormality was detected in the cells of the peritoneum of infected-and-treated animals. It is noteworthy that after upwards of twenty-four hours a small loopful of the peritoneal exudate of these mice yielded an abundant growth of streptococci, although in the direct films they appeared scanty. Several animals which continued to be fed with the drug for a week after inoculation remained well during this time, but when treatment was then stopped they died about a week later from streptococcal septicaemia.

Although the heavy dose of virulent organisms had been almost completely suppressed, the body was not sterilized,* and any immunity which developed was not sufficient to delay for long the recrudescence of the infection. The culture recovered from one of these animals when tested for virulence was found to be distinctly attenuated as compared with that originally used for inoculation—a result which may be adequately accounted for by the influence of the tissues after the initial action of the drug. In the case of intraperitoneal inoculation with pneumococci, McIntosh and Whitby have demonstrated that successful treatment with M & B 693 does not prevent initial invasion of the blood stream. Their finding that in treated animals the capsules of pneumococci in the peritoneal cavity show changes of the nature of defect is, on the analogy of *B. anthracis*, what would be expected when conditions are unsuited for continued proliferation and the development of pathogenic properties. In fact, the events occurring in the course of cure of various infections by sulphanilamide drugs

resemble strikingly those observed by Preisz (1909) in immunized animals inoculated with virulent anthrax bacilli. Gay *et al.*, in an extensive investigation of possible anti-bacterial properties of sulphanilamide, have failed to obtain results markedly significant of such action.

Accordingly it would appear that while it is under the influence of the drug the body behaves towards particular highly virulent organisms much as the normal body does to those of non-pathogenic type, which are gradually destroyed. Levaditi and Vaisman have reached a similar conclusion from their observation that certain drugs of the sulphanilamide group render mice resistant to various endotoxins—that is, killed cultures of organisms such as gonococci (confirmed by Carpenter *et al.*, 1938), meningococci (questioned by Feinstone *et al.*, 1938; and Gross, Cooper, and Lewis, 1938), *B. aertrycke*, dysentery bacilli, etc. Thus the chief action of the drug would be to alter the reactivity of the host. Such an explanation also fits well the clinical observation in pneumonia that the first and most striking effect of the treatment is reduction of the pyrexia. Of course, it is impossible to exclude some gradual direct action of the drug on the organisms' metabolism taking place in the infected animal, according to the hypothesis of McIntosh and Whitby, who found that successful treatment does not prevent an initial multiplication of the inoculum. However, there is little positive evidence to support this as yet. The failure of the sulphanilamide drugs to influence an infection in mice with *Trypanosoma equiperdum* (Buttle, *et al.*, 1936) or *T. brucei*, which pursues a course similar to that with the streptococcus, or the more chronic *T. congolense* infection, even when treatment is intense and prolonged (P. Browning, unpublished), makes unlikely any universal tissue reaction such as stimulation of the reticulo-endothelial system. Histological evidence also has been said to afford no support for such stimulation. But this does not exclude the participation of the usual cellular defensive mechanisms in conducing to cure by means of those drugs.

Practical Conclusions

1. While cure of certain experimental general infections with bacteria by sulphanilamide compounds is an enormous advance, the efficacy of the present drugs is by no means ideal. So far at least as mice are concerned, sulphanilamide is surpassed by, for example, M & B 693, rodilone, and especially the related monoacetyl *p-p*-diamino-diphenyl sulphone. But with them all the range of effective dosage is not wide, and treatment to be most successful must be begun early.

2. Further work on the absorption and excretion of members of the series, and on the chemical transformations which they may undergo in the body, as well as clinical observation, is required in order to enable them to be used to the best advantage as regards the choice of drug, and the dosage in amount and spacing in time. For instance, in treating infection of the meninges by oral or subcutaneous administration it will probably be found that one which rapidly attains a high concentration in the cerebrospinal fluid, such as sulphanilamide, ought to be given in the first place, while more potent but more slowly absorbed compounds, provided they ultimately reach the spinal fluid in sufficient amount, are used to reinforce and maintain the action.

3. Where conditions favourable to general infection have arisen prophylactic use of the sulphanilamide group should be valuable. Where infection has been established the best therapeutic effects are secured in acute, diffuse condi-

* MacCallum and Findlay (1938; and personal communication from Dr. Findlay), in the case of mice infected with the virus of lymphogranuloma inguinale, found the virus particles still present in the brain after successful treatment with sulphanilamide.

tions without marked local tissue changes. Hence these drugs are not likely to obviate operative procedures where focal lesions such as accumulations of pus or extensive necrosis of bone or other tissues have occurred. Accordingly, powerful antiseptics which act when brought into close contact with the organisms locally, and which are relatively harmless to the tissues, will continue to be required also in the treatment of many septic infections.

REFERENCES

- Berger, E. (1937). *Klin. Wschr.*, **16**, 53.
 Britton, C. J. C. (1938). *Brit. J. exp. Path.*, **19**, 140.
 Buttle, G. A. H. (1937). *Proc. roy. Soc. Med.*, **31**, 154.
 — and Parish, H. J. (1938). *British Medical Journal*, **2**, 776.
 — et al. (1936). *Lancet*, **1**, 1286.
 — (1937a). *Ibid.*, **1**, 681.
 — (1937b). *Ibid.*, **1**, 1331.
 Campbell, A. D. (1937). *Amer. J. Syph.*, **21**, 524.
 Carpenter, C. M., et al. (1938). *Science*, **88**, 530.
 Colebrook, L., et al. (1936). *Lancet*, **2**, 1323.
 Feinstein, W. H., et al. (1938). *Johns Hopk. Hosp. Bull.*, **62**, 565.
 Finkelstein, R., and Birkeland, J. M. (1938). *Science*, **87**, 441.
 Garrod, L. P. (1935). *J. infect. Dis.*, **57**, 247.
 Gay, F. P., et al. (1939). *J. exp. Med.*, **69**, 607.
 — and Clark, A. R. (1937). *Ibid.*, **66**, 535.
 Gross, P., Cooper, F. B., and Lewis, M. (1938). *J. infect. Dis.*, **63**, 245.
 — et al. (1938a). *Proc. Soc. exp. Biol.*, N.Y., **38**, 275.
 — (1938b). *Ibid.*, **39**, 12.
 Habs, v. H., and Bader, R. (1938). *Z. Immunforsch.*, **94**, 66.
 Hoare, E. D. (1938). *Lancet*, **1**, 655.
 Hoge, W. G., et al. (1939). *J. infect. Dis.*, **64**, 27.
 Huntington, R. W. (1938). *Proc. Soc. exp. Biol.*, N.Y., **38**, 328.
 Kolmer, J. A., et al. (1938). *Ibid.*, **39**, 95.
 Kreidler, W. A. (1937). *Ibid.*, **37**, 146.
 Levaditi, C., and Vaisman, A. (1938). *Ann. Inst. Pasteur*, **61**, 635.
 Lockwood, J. S., et al. (1938). *J. Amer. med. Ass.*, **111**, 2259.
 Long, P. H., and Bliss, E. A. (1937a). *Canad. med. Ass. J.*, **37**, 457.
 — (1937b). *J. Amer. med. Ass.*, **108**, 32.
 Longcope, W. T. (1938). *Amer. J. med. Sci.*, **195**, 577.
 MacCallum, F. O., and Findlay, G. M. (1938). *Lancet*, **2**, 136.
 McIntosh, J., and Whitby, L. E. H. (1939). *Ibid.*, **1**, 431.
 McKinley, E. B., et al. (1939). *J. infect. Dis.*, **64**, 36.
 Maclean, I. H., et al. (1939). *Lancet*, **1**, 266.
 Maier, N. R. F. (1938). *Science*, **87**, 439.
 Marshall, E. K., et al. (1938). *Johns Hopk. Hosp. Bull.*, **63**, 318, 328.
 Mayer, R. L. (1938). *C. r. Soc. Biol.*, Paris, **129**, 480.
 Mellon, R. R., et al. (1938). *Sulfanilamide Therapy of Bacterial Infections*, Baltimore.
 Mietzsch, F. (1938). *Ber. dtsh. chem. Gesell.*, **71**, 15.
 Natvig, H. (1938). *Norsk Mag. Laegevidensk.*, **99**, 631.
 Nicholson, W. F. (1938). *British Medical Journal*, **1**, 115.
 Preisz, H. (1909). *Zbl. Bakt.*, 1 Orig., **49**, 341.
 Proom, H. (1937). *Lancet*, **1**, 16.
 Schütze, H. (1939). *Ibid.*, **1**, 266.
 Seastone, C. V. (1937). *J. Immunol.*, **33**, 403.
 Whitby, L. E. H. (1937). *Lancet*, **1**, 1517.
 — (1938). *Ibid.*, **1**, 1210.
 Wilson, G. S., and Maier, I. (1939). *British Medical Journal*, **1**, 8.
- (A full list of references to earlier work will be found in Findlay's *Recent Advances in Chemotherapy*, second edition, London, 1939, and in Mellon, Gross, and Cooper's *Sulfanilamide Therapy of Bacterial Infections*, Baltimore, 1938.)

The International Council of Ophthalmology met in London at the Royal Society of Medicine on April 19, 1939, when there were present Messrs. Nordenson (chairman), Ehlers (honorary secretary), Pflüger (honorary treasurer), Paton, v. Grosz, Cavara, Terrien, MacCallan, Bailliant, Löhlein, Knapp, Tooke, Sinclair. The treasurer reported that the assets of the council amounted to 2,232 Swiss francs. The president announced that the German Government had accepted the responsibilities undertaken by the former Austrian Government with regard to the next congress. Professor J. Meller of Vienna was nominated president of the next congress, and the following topics were chosen for discussion: (1) Plastic operations on the eyeball; (2) The pathology of retinal detachment, including the biology and pathology of the vitreous body. It was decided to lay before the next congress the question of internationalizing the determination of the axes of astigmatism. The next meeting of the council will be held in Paris in 1940 on the day before the opening of the annual meeting of the Société Française d'Ophthalmologie.

PHARMACOLOGY OF THE SULPHANILAMIDE GROUP OF DRUGS*

BY

G. A. H. BUTTLE, M.A., M.R.C.S.

Wellcome Physiological Research Laboratories

The Drugs and their Uses

The drugs now in general use for the chemotherapy of bacterial infections are sulphanilamide and certain of its derivatives, notably 2-sulphanilylaminopyridine (M & B 693). Sulphanilamide consists of a benzene ring to opposite ends of which are attached an amino group and a sulphonamide group (Diagram 1). From the chemical

DIAGRAM 1.—Sulphanilamide

point of view the derivatives of sulphanilamide can be divided into those in which substituents are introduced into the amino group and those in which substituents are introduced into the amide group. The drugs of the first class, where the amino group is substituted, are the original prontosil (rubrum),† prontosil soluble, rubiazol, proseptasine, and soluseptasine (Diagram 2). It is prob-

DIAGRAM 2.—Compounds of Sulphanilamide. Class I: Substituents of Amino Group

able that all these substances owe their activity to their splitting up in the body into sulphanilamide and an inactive substance, although this cannot be regarded as finally proved.

Class 1 Compounds

Fuller (1937) showed that after the administration of prontosil rubrum and prontosil soluble large amounts of sulphanilamide were excreted in the urine. His work

* Read at a joint meeting at the Annual Meeting of the British Medical Association, Aberdeen, 1939.

† Prontosil rubrum was the original material discovered by Domagk. Prontosil album is the name given to one of the brands of sulphanilamide or *p*-aminobenzenesulphonamide.