events; and for various reasons I very much doubt if the same underlying sequence can be postulated for the types of headache I have been considering here. I should much appreciate the views of others on this speculation.

Finally, I am very conscious of deficiencies, distortions, and omissions, and of the fact that this paper is full of the dogmatism and economy of truth which is perhaps inseparable from a brief presentation of such a colossal subject.

Summary

The recognition of headache as evidence of a psychiatric problem is discussed.

Positive points in the diagnosis of such headaches are put forward as a supplement to the principle of exclusion, so apt to be used alone.

REFERENCES

- Brown, F. (1936). *J. ment. Sci.*, **82**, 295.
 Claparède, E. (1924). Quoted by Schilder, *op. cit.*
 Curran, D., and Slater, E. T. O. (1935). *Lancet*, **1**, 69.
 Lewis, A. J. (1934). *J. ment. Sci.*, **80**, 277.
 Mayer-Gross, W. (1938). *British Medical Journal*, **2**, 936.
 Muncie, W. (1932). *Arch. Neurol. Psychiat.*, Chicago, **27**, 352.
 Reilly, T. F. (1926). *Headache: its Cause and Treatment*, Philadelphia.
 Schilder, P. (1933). *J. nerv. ment. Dis.*, **78**, 1, 137.
 Spriggs, E. I. (1935). *Lancet*, **2**, 1, 63.
 Wilson, H. (1938). *Ibid.*, **1**, 367.
 Wolff, H. G. (1937). *Arch. Neurol. Psychiat.*, Chicago, **37**, 895.

RESULTS WITH EPANUTIN IN MENTAL HOSPITAL PATIENTS

BY

SUSANNAH DAVIDSON, M.B., B.S., D.P.M.

AND

JOHN D. SUTHERLAND, B.Sc., Ph.D., M.B.

Assistant Physician, Glasgow Royal Mental Hospital

We have recently treated twelve epileptic patients in the Glasgow Royal Mental Hospital with epanutin (sodium diphenyl hydantoinate) in place of either sodium phenobarbitone or prominal. The drug was administered according to the advice of the makers (Parke, Davis and Co.) by gradual replacement of the drug already in use, and the changes in the frequency of fits or in the mental state were studied. The following notes give the results.

Case Reports

Case 1.—A man aged 30, whose first fit occurred eight years previously, has been three years in hospital. He is well preserved intellectually, with several characteristics of the epileptic personality, especially irritability and egocentricity. Major and minor attacks occur with about equal frequency. As he has ground parole and there is amnesia for the fits it is impossible to keep accurate account of the incidence. Before epanutin was given he was receiving in alternate periods of about six weeks either 1 grain of sodium phenobarbitone or 3 grains of prominal twice daily. These changes were usually followed by a considerable reduction in the number of fits for about two weeks and an associated improvement in his mood. Epanutin, one capsule (0.1 gramme) thrice daily, was given for sixty-four days. During the first month there was a striking diminution in the fits, and he was bright and enthusiastic about the new treatment. In the following five weeks, however, he had more fits than usual, twenty being recorded. He relapsed into his former state and requested that the drug be changed; no toxic effects occurred. On resuming prominal he had only seven recorded fits in the next twenty-five days.

Case 2.—A man aged 31 had his first fit when six years old, and he has been ten years in hospital. He is a deteriorated childish personality, with minor fits almost as frequent as major fits. Usually he is maintained on 3 grains of prominal twice daily. Epanutin, one capsule thrice daily, was given for thirty-six days, and was stopped because of increased irritability and difficult behaviour, though no other toxic symptoms were observed. There was a slight increase in the number of major fits and a diminution in minor attacks.

Case 3.—A man aged 46, whose fits began twenty years ago, has been thirteen years in hospitals. His personality is very deteriorated and his behaviour noisy, aggressive, and impulsive. Before the change to epanutin he was given 3 grains of prominal twice daily. On one capsule of epanutin three times a day the fits were less frequent, but the drug was stopped because of nausea and increased irritability.

Case 4.—A man aged 40, with fits during the past twenty years following a war injury, since when he has been in hospitals, has minor fits at long intervals and an occasional major fit. His personality is deteriorated, and occasionally he has attacks of vomiting with headache, though no signs of raised intracranial pressure are present. Epanutin, one capsule thrice daily, was stopped after thirty-six days when one of his vomiting attacks became unusually severe and prolonged. The vomiting disappeared shortly after the withdrawal of the drug; it is probable that the drug was causing an exacerbation of the usual attack. No change was observed in his mental condition, nor was there any significant change in the number of fits.

Case 5.—A man aged 40 whose fits started at the age of 10, has spent the last twenty-six years in hospitals. His mental condition is very deteriorated, his speech being incoherent. The fits are all major ones, and he is usually given prominal. Epanutin, one capsule thrice daily, had to be stopped after thirty-four days on account of nausea and refusal of food. There was no change in his mental condition, and his fits were more frequent.

Case 6.—A man aged 48, whose first fit occurred when he was 22 years old, has been in hospital five years, during which time he has shown progressive deterioration, with considerable memory defect, some confusion, and occasional outbursts of irritability. He has major fits, mainly during the night, at long intervals while on prominal. Each major fit is followed by about twenty-four hours of marked confusion. Epanutin, one capsule thrice daily, had to be stopped after nineteen days because of increased confusion and ataxia, which made it difficult for him to find his way about the ward and which prevented him from being able to feed himself. He was also much more dull and unresponsive than usual. The toxic manifestations disappeared a few days after withdrawal of the drug.

Case 7.—A man aged 55, whose fits began when he was 14 years old, has been thirty-four years in hospital. He is an extremely egocentric childish personality, vain and with numerous mannerisms, who has slightly more minor than major fits. Epanutin, one capsule thrice daily, was stopped on the thirty-seventh day of its administration when he developed an erythematous rash on his hands and legs which gradually spread over his body and which was accompanied by intense itching. Later, oedema appeared and was succeeded by an exfoliative dermatitis. There was no change in his mental condition, but the fits ceased during the epanutin period. The toxic manifestations were relieved by intravenous sodium thio-sulphate.

Case 8.—A woman aged 51, with onset of fits in childhood, has been nineteen years in hospital and is now a somewhat deteriorated person with great irritability. Her attacks are usually major with occasional minor fits. Epanutin, one capsule twice daily, was given for fifteen days, when she refused the drug because of headache and nausea. There was an increased irritability, but there was no change in the incidence of the fits.

Case 9.—A woman aged 47, with a history of fits since infancy, has been twenty-eight years in hospital. She is fairly

well preserved mentally, with marked religiosity and spells of irritability. Major and minor fits are about equal in frequency, and she is usually given 1 grain of sodium phenobarbitone twice a day. Epanutin, one capsule twice daily, was maintained for sixty-seven days without any change in her mental state or in the frequency of the fits being noted. The drug was stopped when a vesicular rash appeared over the upper half of the right side of the trunk and spread down the right arm.

Case 10.—A woman aged 32 has had fits since the age of 9 years. She has been in hospital for four years and is a very demented personality. Major and minor fits are about equal in frequency. Epanutin, one capsule twice daily, has been given for sixty-eight days with no change in her mental status, but a diminution in the number of fits. No toxic effects developed.

Case 11.—A woman aged 53 has had fits for the past thirty-eight years, twenty years of which have been spent in hospitals. She is a depressed and rather demented bedridden woman whose fits, usually major, are infrequent. On the seventeenth day after epanutin, one capsule twice daily, was begun she asked to have the drug stopped because of headache, dizziness, and general malaise. It was noticed at the same time that a blotchy erythema had appeared on her legs.

Case 12.—A woman aged 49, whose fits began at the age of 13 years, has been in hospital three years and is now a noisy, excitable, and deteriorated personality. She was formerly given 1 grain of sodium phenobarbitone twice a day. As she became noisy at night after epanutin was substituted, she was given 1 grain of sodium phenobarbitone at nights in addition to one capsule of epanutin twice daily. There has been no change in her mental condition or in her fits after seventy-eight days of the epanutin, and no toxic effects developed.

Conclusions

Before drawing any conclusions it must be pointed out that these patients are a rather selected group. Most of them are over 40 years of age and have had a long history of hospitalization and a correspondingly long treatment with drugs. They were almost all admitted for psychotic behaviour as well as fits, and most of them were unsuitable for epileptic colonies, as they were unable to work.

The outstanding result of the treatment is the appearance of toxic effects, eight of the twelve cases showing these within a few weeks. They included headache, giddiness, nausea, loss of appetite, and ataxia. Three cases developed rashes, severe in two, with one being followed by an exfoliative dermatitis. These effects have already been referred to in connexion with this preparation (see references), but their frequent and early appearance obviously restrict the use of the drug, for though some are possibly not serious in themselves they antagonized these patients. One point perhaps worthy of note is that the three youngest patients showed no toxic effects.

In considering the mental effects we had only one patient who showed improvement, but this lasted for only a few weeks, after which he resumed his former state. It has also to be recalled that this patient showed a temporary improvement upon a change from sodium phenobarbitone to prominal or vice versa. The effect of the change to epanutin was rather more marked, however. His relapse followed upon the recurrence of fits, but the appearance of toxic effects in other cases made us reluctant to increase the dose. It would appear that in his case, as in some of the others, epanutin makes a useful addition to the drugs on which patients may be maintained for a few weeks. The absence of sedative effect proved an undesirable feature with several of our patients, as it

released the excessive irritability and, in some cases, impulsive behaviour which was much relieved by the other drugs.

With regard to the anti-convulsant properties, the data are too limited to make critical observations. The table shows the number of fits to be expected during the period that epanutin was given, this being calculated from the average incidence of fits in the preceding six months in each case. As there appeared to be little change in the character of fits in most cases we have combined the major and minor attacks in the term "fit." It will be seen that there is a balance in favour of epanutin.

Table showing Summary of Results

Case No.	Age in Years	Duration of Epilepsy in Years	Change in Mental State during Epanutin Period	Period of Epanutin Administration in Days	No. of Fits in Epanutin Period	No. of Fits expected on Former Drugs	Reasons for stopping Drug
1	30	8	Brighter for 4 weeks	64	Not recorded	4	More fits than usual
2	31	25	Increased irritability	36	4	5	Increased irritability
3	46	20	" "	34	2	4	" " and nausea
4	40	20	No change	36	3	2	Vomiting
5	40	30	"	34	13	10	Nausea; refused food
6	48	26	Dull and unresponsive; increased confusion	19	1	2	Confusion and ataxia
7	55	41	No change	37	0	4	Severe rash followed by exfoliative dermatitis
8	51	45	More irritable	15	3	2.5	Headache and nausea; refused drug
9	47	46	No change	67	7	8	Vesicular rash
10	32	23	"	68	3	5.5	Not stopped
11	53	38	"	17	—	—	Rash, headache, and giddiness
12	49	36	"	78	11	12	Not stopped

In short, our experience suggests that the drug is of use in the younger and better-preserved patients, but on account of its toxic effects it is advisable not to maintain it for more than a few weeks at a time.

We are indebted to Dr. Angus MacNiven, physician-superintendent of the Glasgow Royal Mental Hospital, for permission to give the treatment and to publish the results. We have also to express to Messrs. Parke, Davis and Co. our thanks for a trial supply of the drug and for advice.

REFERENCES

- British Medical Journal*, 1939, 2, 179.
Lancet, 1939, 1, 180.

Muscular Dystrophy and Hypothyroidism.—In the issue of *Revue Neurologique* for May, 1939 (p. 513), P. Mollaret and J. Sigwald contribute a paper on "Generalized Muscular Atrophy of Rapid Onset in an Adult associated with Latent Myxoedema clinically cured by Thyroid Medication." In the same number (p. 548) G. Bourguignon also deals with the improvement effected by thyrid treatment of the condition. A case of generalized muscular hypertrophy without weakness or myotonia is described in a man of 56. The basal metabolic rate was -13 per cent., and there were clinical signs of hypothyroidism. Biopsy showed no definite muscular abnormality. These symptoms, and also alterations in electrical excitability of the muscles, disappeared with thyroid medication. The literature on this condition is reviewed. The muscular changes are thought to be directly related to the hypothyroidism.