oestrol) has come to the fore. This compound can be prepared synthetically, and already its cost is much less than that of oestrone and oestradiol. Moreover, it can be administered orally or parenterally, and the evidence so far available shows that it is as active by mouth as it is by injection. One milligramme administered orally exerts an effect equal to that of the same amount of oestradiol benzoate injected intramuscularly. The only disadvantage is that stilboestrol in large doses frequently produces a sensation of nausea and malaise. This occurs after parenteral as well as oral administration. It is doubtful whether stilboestrol, which, so far as we know, does not occur naturally, should be regarded as a "hormone." I prefer to regard it as a hormone substitute.

Stilboestrol, even when given by mouth, has been shown to produce good results in the treatment of genital hypoplasia, amenorrhoea, menopausal symptoms, senile vaginitis and vulvitis, suppression of lactation, and, in fact, all the conditions for which oestradiol and oestrone have proved their value. Moreover, I have used it with success to induce labour in cases of intra-uterine death of the foetus. The dosage laid down for oestradiol should form the basis of stilboestrol therapy, but when it is administered by mouth it is advisable to give it more frequently and in slightly larger quantities. The same principles of treatment should be followed, and it is rarely necessary to give more than 5 mg. daily.

At the present time stilboestrol therapy is still in the trial stage, but the initial reports go to prove its value. Figs. 1 and 2 illustrate the effect of stilboestrol on the vaginal epithelium. The original smear (Fig. 1) was obtained from a woman suffering from senile vaginitis: 50 mg. of stilboestrol were administered by mouth during a period of five days and no local antiseptics were used. The second smear, that of a healthy vagina, was obtained six days after beginning treatment.

The efficacy of stilboestrol is further evidenced by a recent case of kraurosis vulvae occurring in a woman aged 72. The condition had been present for five years, during which time she had received all manner of treatment, including x-ray therapy on two occasions. No form of treatment had given her more than slight and temporary relief from her intense pruritus. When the patient was first seen by the author the vulva showed a severe degree of kraurosis vulvae with superimposed ulceration and induration—the result of scratching. The lightest touch produced unbearable pain (Fig. 5). Large doses of stilboestrol were given by intramuscular injection and also by mouth. In the course of seventeen days she received 220 mg. of stilboestrol (probably an unnecessarily large dose), but no other treatment except the local application of an analgesic ointment to ensure sleep. At the end of that time the second photograph (Fig. 6) was taken, and this shows the vulva completely healed except for two small ulcers. The patient was then given 3 mg. of stilboestrol daily, and one month later she was free from symptoms and the kraurosis was cured.

Hormone therapy is, no doubt, still in a crude state, but already it has passed beyond the stage of "gland extracts." Oestrogenic substances, both natural and synthetic, are pure chemical compounds dispensed by weight and possessing remarkable properties. It is truly amazing that such small amounts of material can exert such pronounced local and general effects. Stilboestrol is an adequate substitute for the natural oestrogens, and possesses advantages in that it is cheaper and is also equally active when administered by mouth. Treatment must, however, be based on physiological principles, and often needs to be prolonged for some time before its full effect is manifest. Careful selection of cases suitable for treatment is essential, and the empirical use of oestrogenic substances can only lead to disappointment.

GLOMANGIOMA

A FORM OF "PAINFUL SUBCUTANEOUS TUBERCLE"*

BY

ALAN C. LENDRUM, M.A., B.Sc. M.B., Ch.B.

Assistant to the Professor of Pathology, University of Glasgow; Assistant Pathologist, Western Infirmary, Glasgow

AND

W. ARTHUR MACKEY, M.B., Ch.B., F.R.F.P.S., F.R.C.S.Ed.

Assistant to the Professor of Surgery, University of Glasgow; Dispensary Surgeon, Western Infirmary, Glasgow

(WITH SPECIAL PLATE)

Less than three years ago we published a brief paper on an entity then regarded as extremely rare, the tumour of the neuro-myo-arterial glomus (Mackey and Lendrum, 1936). In concluding our paper we quoted the aphorism of Picard (1931), that recognition of this, as of all rarities, demands only that it be borne in mind: the truth of this has been strikingly shown by the fact that in the intervening brief period we have seen a further twelve cases. Since the lesion is one which is usually productive of quite crippling pain in its victim, and since its treatment, too often unreasonably delayed, is of the simplest and most satisfactory nature, it appears desirable to bring this interesting little tumour to more general notice.

It would be surprising had a lesion, with such a clamant syndrome remained quite unobserved, and indeed there can be little doubt that tumours of this nature must have constituted a portion of the group described in 1812 by Wood as painful subcutaneous tubercle (Greig, 1928); but its histological identification had to await the work of Masson (1924), who also suggested its probable origin from the normal arteriovenous anastomoses present in the skin.

Clinical Features

It is characteristic of these cases that symptoms greatly overshadow signs. The patient presents himself because of pain, usually in a limb, and generally paroxysmal. In many cases the pain is related to a "trigger point," sensitive to a minimum of trauma; in others the attacks are elicited by change of temperature or even by change of position; whilst attacks no less severe may seem entirely spontaneous in origin. If the tumour lie in the fatty subcutaneous tissue it may take the form of a bluish nodule of angiomatous appearance, which has frequently been observed to engorge during a painful attack. The patient, however, may fail to observe the nodule or to relate it to his pain. In some cases, especially if the lesion be subungual, no abnormality is to be seen.

Inquiry usually elicits the fact that the symptoms have developed gradually, sometimes over many years, and frequently the onset is related to a single obvious trauma, especially in the subungual tumours. The typical pain is often preceded by vague tinglings and mild discomforts; it is at first moderate in degree, local in distribution, and merely occasional, but it tends to increase in severity, to radiate more and more widely, and to occur with greater frequency. Ultimately the pain is agonizing, and the patient may even demand amputation of an affected finger

* Read in the Section of Pathology and Bacteriology at the Annual Meeting of the British Medical Association, Aberdeen, 1939.

(Case V; and Fingerland, 1938). If aware of the origin of his pain, he often evolves a complicated defence mechanism to protect the trigger area. One of our patients (Case I), a physician, had come to use only those pockets accessible to his left hand because a glomangioma on the back of his right hand made its use intolerable, and for the same reason he would never shake hands in the normal fashion. Another (Case II) slept with his arm over the edge of the bed lest the bedclothes might touch the sensitive area.

The radiation of the pain does not follow the distribution of any skeletal nerve, but tends to be diffuse, to involve an entire limb, to spread to the adjacent part of the trunk, and even to involve another limb. Blumental (1937) describes a woman of 53, who for sixteen years had increasing pain in the left arm with radiation to the precordial area, leading to a false diagnosis of angina pectoris. During the last three years a small blue subungual tumour had developed on the ring finger of the left hand. Surgical removal led to the cure of her pain. Histologically the tumour was a typical glomangioma.

It is often in the milder attacks that the patient is best able to identify the focus of origin of his pain; when the paroxysm is more violent he may feel that a large area is suddenly involved or even that the pain begins elsewhere and spreads toward the focal point. The type of pain described above is similar to that which Leriche (1939) has attributed to disturbance of the sympathetic neurovascular mechanism. In this regard it is noteworthy that many patients have observed the affected limb to be constantly either warmer or colder than its fellow (Case II), or to show disturbance of the sweat mechanism (Lee, 1938). Barré's case (Masson, 1935) actually showed on the affected side a Horner syndrome, which disappeared on removal of a subungual tumour. Stabins *et al.* (1937) have studied the vasomotor reactions associated with glomus tumours, and report two cases in which before removal of the tumour the limbs showed a markedly asymmetrical recovery of temperature after cooling, a disparity which disappeared in a week or two following operation.

Case Histories

CASE I

A man aged 48. Tumour on dorsum of right hand, just proximal to index knuckle. His trouble began at the age of 39; at that time he was operated on for appendicitis, and, following the operation, the hand, which apparently had been placed behind his hip, was painful. This pain disappeared and he had no complaint for three years. Thereafter he became subject to attacks of pain of progressively increasing severity, localized on the dorsum of the right hand in the region of the index metacarpal. These attacks were almost invariably induced by trauma, even of mild degree, and ultimately he devoted great care to the shielding of the hand from the slightest external stimulus (hand-shaking, touch of clothing). The pain was worse when the hand was dependent, and was slightly alleviated by elevation of the limb; it did not radiate, and it was never elicited by temperature changes. There was no local vasomotor upset, but during a severe attack of pain his face was blanched.

The patient, a physician, was mystified by the earlier attacks, for there was no visible local abnormality, but in the course of time he became aware of an extremely sensitive bluish nodule which increased gradually in size, and at the time of consultation (March 23, 1938) was fully three-quarters of an inch in diameter. It was roughly hemispherical, of a bluish colour suggesting a subcutaneous varix, and the patient stated that when painful it was notably engorged. The diagnosis of glomangioma seemed obvious, and operative removal was

carried out under local anaesthesia by one of us (W. A. M.). Histologically this is a typical glomangioma of solid cellular type but showing in addition a pronounced dilatation of one of the blood spaces. There has been no recurrence of pain (May, 1939).

CASE II

A man aged 73. Outer side of right upper arm. In this case symptoms were severe over a period of thirty years. The patient cannot remember the onset, but he has been aware for the period stated of a small bluish lump on the outer side of his right arm. It was agonizingly painful if touched; he was led, therefore, to avoid crowds, and even to sleep with his arm over the edge of the bed to avoid the contact of the sheets. Despite his care the pain gradually became worse, both in severity and in frequency, and some attacks might stop his work for half an hour. He knew no way of relieving it. There was no radiation of the pain. Occasional attacks were spontaneous in origin. During the crises the tumour was turgid and prominent. At all times the affected arm was colder than its fellow; since removal of the tumour by Mr. J. J. Robb, of Dundee (1938), this phenomenon has disappeared (395/38, Dundee Royal Infirmary). This is a well-encapsulated glomangioma of typical fully differentiated type.

CASE III

A man aged 61. Web between fourth and fifth fingers of right hand. In this case the glomangioma appears to have developed as a direct result of trauma. Ten years ago the fifth finger was abducted violently by accident, causing severe pain; thereafter the patient gradually became subject to attacks of pain in the injured web, and concurrently he noted the development there of a small bluish nodule. The pain was often spontaneous, but was readily provoked by touching, so that he would not allow the doctor to handle the part. In some attacks it appeared to shoot down the arm from the shoulder to the hand. He found that he could diminish the pain by immersing the hand in hot water, but in no other way. He had not noted any change in the tumour during the pain, or temperature difference between the two limbs. The tumour was removed by Mr. Walter Campbell, of Dundee (1938). (679/38, Dundee Royal Infirmary.) This is a typical glomangioma of differentiated type.

CASE IV

A man aged 24. Left shin. For fourteen years he had had a bluish lump on the middle of the front of the left shin, for the most part without discomfort. For the last two years, however, there had been increasing pain; he described this as a feeling of stiffness and soreness, situated in the lesion, and radiating laterally to the calf, and up and down the front of the leg and to the knee. It was aggravated by exercise, especially when, at his work, he carried heavy loads. Occasionally pressure over the site caused pain to shoot up the front of the thigh. The lesion appeared to be more sensitive to heat than to cold, and latterly his sleep had thus been much upset. He noted that when painful the lesion became congested, swollen and dark in colour. There was no history of trauma.

Examination (W. A. M.) revealed a mulberry-like mass with a coarsely nodular surface, bluish in colour as if engorged with blood; it was one and a half inches in vertical length by one inch in breadth. The patient made no complaint of tenderness when it was examined. The first impression was of a localized varicosity, but the absence of change in the remainder of the saphenous system, the long history, and the slow growth led to a diagnosis of venous angioma. So thin was the overlying skin that some sacrifice of this at operation seemed probable. Under local anaesthesia (February 26, 1936) a short transverse flap was therefore raised; the lesion was then seen lying almost free, like a bunch of small grapes, in a pocket in the subcutaneous fat. It was dissected out with ease, and found to be united, not to a vein, but merely to a thin fairly firm pedicle, which was clamped and ligated. It seems probable that this included the afferent vessel of the

tumour (162/1936, Western Infirmary, Glasgow.) This, our first case, is fully described in our previous publication. It is histologically the most differentiated glomangioma of the series, a fact which helped the microscopical identification (Fig. 2).

CASE V

A woman aged 44. Tumour subungual, left middle finger. This case has a long history, for symptoms were noted first at the age of 11, following punishment at school with "the strap." Thereafter the left hand was noted to become numb on exposure to cold. The time of actual appearance of the pain was not exactly remembered. In the early stages she had from time to time the sensation of a needle prick in the tip of the left middle finger, and by the age of 20 the syndrome was well established. She was suffering then from attacks of pain in the finger which radiated over an ever-increasing area, ultimately involving the homolateral half of the body. The attacks varied in severity, minor ones always localized in the finger or centred in it, while the more severe were of wide distribution and might appear to start elsewhere—for example, in the toes of the same side. Recent attacks were noted sometimes to be spontaneous, but more usually were precipitated by trauma to the finger or by exposure to heat or cold. Pain caused by trimming of the nail was described as maddening. She knew of no method of aborting the pain, which by 1928 was occurring in attacks lasting one and a half hours and more recently was practically continuous. A noteworthy feature of the case was the fact that the affected finger showed no visible abnormality; nevertheless she had asked repeatedly for its amputation, and so convinced was she of the site of origin of the pain that she cut into the finger-tip with a razor. Before this had healed she consulted Mr. C. A. Ferguson, who in March, 1937, made a presumptive diagnosis of glomangioma, and removed a small portion of subungual tissue (March 30, 1937). (338/1937, Western Infirmary, Glasgow.) The tumour is a glomangioma of solid cellular type, and does not appear to be completely removed.

This patient was seen in the autumn of 1938 and again in April, 1939; she has been entirely free of symptoms since the operation (two years), and there is no sign of recurrence. In view of the work of Stabins *et al.*, mentioned above, and of the possibility of persisting tumour remnant, it was decided to examine the temperature reactions of her hands. Dr. J. B. Gaylor kindly measured (April 25, 1939) the vascular dilatation as induced by the immersion of the legs in hot water, taking the skin temperature as index of blood flow. There was no significant difference between the two sides in the initial temperature of the finger-tips (indices and middle) under resting conditions or after the induction of vascular dilatation. Thus the dilatation in the involved finger and in the normal counterpart on the other side was both equal and normal. There appears to be at present no upset of the vasodilator response as assessed by skin temperature measurement. If, therefore, removal of the tumour has been incomplete, the remnant may well have become cicatrized, for there are neither symptoms nor signs of its presence.

CASE VI

A man aged 53. Tumour on buttock; age at onset, 33 years. The history of this case extends over a period of twenty years, and it begins with a fall in the sitting position. Shortly thereafter the patient became aware of a minute nodule in the buttock which was the centre of a sort of itching discomfort, gradually developing into pain of increasing severity. The pain occurred in attacks, some spontaneous, others precipitated by temperature change or trauma, and it radiated over an increasing area of the gluteal region. The patient noticed that the tumour was turgid during the painful attacks. He was never able to cut short an established attack of pain, but he did all in his power to shield the site. He sought medical aid first in 1931, when the nodule was incised, without benefit. He consulted Mr. W. A. McLennan (September 29, 1938), who found a well-localized nodule, in appearance like a thinly covered convoluted varix. He performed a wide excision, and since then there has been no pain (April, 1939). (1181/38, Western Infirmary, Glasgow.) The

tumour, which is a cavernous type of glomangioma with a high content of glomus cells and a mature differentiated structure, lies partly in the deeper layers of the dermis where there is no real encapsulation, and partly in the subcutaneous fat, into which it extends as a nodule, demarcated by a well-developed collagenous membrane (Fig. 1). The upper portion contains abundance of myelinated nervous tissue. There is no definite histological evidence of the incision performed in 1931.

CASE VII

A man aged 41 years. Lateral aspect of right elbow. The trouble began at the age of 39 in an insidious way with the appearance on the lateral aspect of the left elbow of a small lump which at first had no special colour but gradually took the appearance of a venous angioma and shaded through blue to black. There were no symptoms in the early stages, but presently the patient began to suffer from tingling sensations in the whole arm which he learned could be relieved by rubbing the lump. The discomfort was not severe, and in fact if his interest were held he forgot it. The tingling attacks, however, gradually became more frequent, and he noted that elevation of the arm would often bring on an attack, a causal relation seen most clearly when he held the newspaper for reading. In the ordinary course the tingling would last for about five minutes, and was usually relieved by hanging the arm down.

Latterly the tumour became obviously the trigger point, a blow on which would set up a paroxysm of numbing pain comparable to that experienced after trauma to the ulnar nerve (funny-bone). The pain lasted generally for a few seconds, but on occasion as long as ten minutes. It died away gradually, and he knew no way of relieving it. During the attacks of pain the whole arm became hot. The nodule was excised by Mr. R. S. Kerr (August 8, 1938). (995/38, Western Infirmary, Glasgow.) This is a sharply demarcated glomangioma lying in the deepest layer of the dermis, its structure virtually identical with the deep nodule of the previous case (Fig. 3). There is considerable myelinated nervous tissue in the periphery. The patient was seen again in April, 1939, and was symptom-free.

CASE VIII

A young woman aged 18. Tumour on dorsum of right foot. From the time of her earliest memory the patient had been aware of a bluish nodule on the dorsum of the right foot. This had been neither painful nor tender until a fortnight before consultation, when it became sensitive to pressure and also the seat of spontaneous pain. The pain spread up the front of the ankle region. She obtained some relief when recumbent. She noted that the nodule was sometimes darker in colour when painful, and, furthermore, that the affected foot was often colder than its fellow. Examination revealed a bluish nodular lesion measuring approximately 2 cm. by 1 cm. and projecting slightly above the level of the skin over the first cuneo-metatarsal joint of the right foot. Its appearance was that of a venous angioma or clustered varix. A presumptive diagnosis of glomangioma was made and the nodule excised by one of us (W. A. M., December 20, 1937). (1330/1937, Western Infirmary, Glasgow.) This is a typical cavernous glomangioma with the glomus cells occurring as a single or double layer in the fibrous walls of the irregular blood spaces (Fig. 4). The patient was seen again (May 9, 1939). Some three months after the operation a bluish nodule appeared in the proximal wall of the operation scar and increased slowly in size. It now measures about 1.5 cm. by 1 cm., and reproduces on a slightly smaller scale the features of the original growth. She still feels the foot abnormally cold, but there has been no pain or tenderness since operation. Dr. J. B. Gaylor kindly investigated the thermal reactions of the feet (May 16, 1939). Vasodilatation was induced in the feet by heating the upper extremities. There was no significant difference in surface temperature between comparable areas on the feet, at resting level or during dilatation. It is justifiable, therefore, to conclude that the apparently recrudescence lesion is producing no objectively demonstrable vasomotor anomaly.

CASE IX

A man aged 53. Tumour on left shin. The history of this case is brief and unusually featureless. For about eighteen months the patient was aware of a small spot on his left shin, which he regarded as a pimple and poulticed without result. It increased slowly in size: it was painful if struck, but was never the seat of spontaneous pain. On September 24, 1926, he consulted Mr. G. H. Edington, who found a dark purplish tumour, about half an inch in diameter, over the subcutaneous surface of the left tibia at the junction of the middle and upper thirds. It was mobile over the deep structures, and not sensitive to ordinary pressure. The appearances suggested melanoma, and free excision was carried out. Recovery was uneventful, and the patient has had no further trouble (October, 1938). It was reported as an unusual haemangio-endothelioma, and the tissues were filed for future reference. The slides were discovered accidentally in 1938 by Dr. J. S. F. Niven, who recognized the lesion as glomangioma. This is histologically identical with, but somewhat larger than, the tumour in Case I.

CASE X

A man aged 77. Ulnar border of forearm. In this case symptoms developed gradually over a period of eleven years. In the course of his occupation as a watchmaker the patient was accustomed to rest his left forearm on the edge of the work-bench, and he became aware that a certain point on the ulnar border of his forearm was abnormally sensitive to this pressure, producing a sharp non-radiating pain. In order to protect the sensitive point he wore a pad on his arm. There were no associated disturbances of the sympathetic nervous system. He was seen in 1938 by Mr. N. J. Logie of Aberdeen, who discovered a small tender nodule midway on the ulnar border of the left forearm, and made a clinical diagnosis of glomangioma. The tumour was removed and the patient's discomfort completely relieved. Histologically the tumour is a glomangioma of solid cellular type.

CASE XI

A man aged 31. Skin over left third metacarpo-phalangeal joint. The clinical history, for which we are indebted to Dr. G. P. Middleton of Ballater, begins in 1929 with the appearance of a small bluish nodule over the third left metacarpo-phalangeal joint. The patient has no recollection of any trauma as possibly causative. Attacks of pain were elicited by lifting weights and by trauma, but not by temperature changes. The attacks were never spontaneous and the nodule was not tender. During the painful periods, which never lasted more than five minutes, the tumour was blue and swollen; pressure on it at this time did not affect the pain. The pain radiated to the front of the joint and distally to the first interphalangeal joint of the same finger, a distribution which had not increased in extent, although the intensity of the pain had become greater. During the paroxysms there was generalized sweating. He knew no way of relieving the pain, which passed off gradually. The tumour was incised in 1935 and some brown granular material obtained; it swelled up again a few months later. He was sent to Mr. Fowler of Aberdeen in November, 1937, and the bluish soft fluctuant tumour was removed (16981/37, Royal Infirmary, Aberdeen). Histologically this is a glomangioma of rather differentiated cavernous type; it shows little evidence of the earlier incision. He has had no trouble since (May, 1939), apart from a little throbbing pain for a day or two in the spring of 1938.

CASE XII

A man aged 59. Medial aspect of elbow. For this and the following case we are indebted to Professor Keith Inglis of Sydney; both were included in our previous paper. While this case was under treatment for epithelioma of the lip a pedunculated soft blue mass was noted on the medial aspect of one elbow and was removed for examination. It is not recorded whether the lesion was painful (Sydney Hospital, 74/1920). Histologically the tumour was a glomangioma of mature structure almost identical with Case IV.

CASE XIII

A boy aged 12. Lateral aspect of knee. This lad presented a painless swelling on the lateral aspect of the knee considered to be the result of a blow with a cricket ball seven years before. At operation an unencapsulated tumour of fatty vascular appearance was found, lying antero-lateral to the tendon of the biceps femoris. It was unattached to skin or to any of the deeper structures (Sydney Hospital, 16001/1936). Histologically this is a typical example of the solid cellular type of glomangioma; it is illustrated in our previous paper.

CASE XIV

A woman aged 46. Pulp of left index finger. We owe this case to the courtesy of Dr. T. C. Patterson, pathologist to the Bolingbroke Hospital, London. His patient had complained for two years of pain in the pulp of the left index finger. This pain had an extraordinary periodicity, for it occurred regularly in attacks beginning one week before the onset of menstruation and persisting until the end of the flow. At these times the pulp of the finger was tensely swollen. She had noted the presence in the pulp of the finger of a small spherical tumour, extremely sensitive to trauma, and on several occasions a knock had brought on such pain that she fainted. In general her suffering was greater in wet weather. It is possibly of interest that as a result of an accident twenty years previously the affected finger was the seat of mallet deformity. The tumour is a white spherical elastic nodule, 0.6 cm. in diameter. Histologically it is a glomangioma of small cavernous type, with abundant glomus cells.

CASE XV

A woman aged 50. Palmar aspect of middle phalanx of left index finger. In 1914 she suffered from pain in the left index finger similar in character and site, so far as she can remember, to her recent complaint. The pain disappeared, however, and there was a period of freedom lasting twenty-four years. About one year ago there appeared a small tender swelling close to the skin on the palmar aspect of the second phalanx of the left index finger. The tenderness had lately become extreme, and deep pressure produced a pain of sickening character. The pain tended to be paroxysmal, and was elicited by the lightest touch or by cooling, as when taking the hand out from below the bedclothes; spontaneous attacks also occurred. The pain radiated to the point of the finger, never up the arm. During the attacks, which lasted from one to several hours, the nodule appeared harder. There were no associated vasomotor disturbances. The only known method of relief was to immerse the hand in water as hot as could be borne. She made conscious efforts to protect the sensitive finger, keeping it flexed and holding her arm against the breast. She can recall no injury to the finger. Mr. Fowler of Aberdeen (to whom we are indebted also for the clinical notes) made a diagnosis of glomangioma, and excised the nodule (727/1939, Royal Infirmary, Aberdeen). Histologically this is a small circumscribed glomangioma of moderately differentiated cavernous type.

In the syndrome produced by glomangioma four features appear to be of outstanding importance; these are pain, radiation of pain, associated vasomotor upset, and an aetiological relationship with trauma. Of our fifteen cases, only one was painless. In eight the pain radiated, and in seven there were associated vasomotor disturbances. The validity of a history of trauma is difficult to assess, but in three of our cases the disease was definitely brought on by injury, and in three others there is a strong presumption that a preceding injury was causative.

Pathology

The gross appearance of this tumour would seem to be primarily dependent on its site. If it occurs in the nail-bed, where there is little room for expansion, it has the form of a nodule of rather cellular-looking tissue, not obviously angiomatous. In other sites the tumour is

situated in the deeper layer of the dermis and commonly expands into the adjacent fat as a bluish somewhat lobulated mass, rarely more than 2 cm. in greatest diameter. It may resemble a localized varix. The occurrence of multiple glomangiomas has been recorded (Bergstrand, 1937; Lee, 1938), but this is rare; the reports have unfortunately not always been accompanied by illustrations (Touraine *et al.*, 1936). Weidman and Wise (1937) describe a case with forty-eight subcutaneous lesions which they consider to be telangiectases of glomera, an explanation they suggest is applicable to other reported cases.

In our histological description we follow the classification suggested by Masson (1935) in his study of forty-seven cases. The general structure is that of a tumour of blood spaces characterized by the presence in the walls of large cuboidal cells, the so-called glomus cells to be discussed later. The three main variations on this theme are produced by unequal growth of the constituent tissue elements; when the glomus cells show a predominance then the first impression is of a sheet of epithelial-like cells in which lie narrow clefts lined by endothelium (Masson's paucivascular form). When the connective tissue is predominant and the vascular spaces larger, the appearances are essentially those of a cavernous angioma modified by the presence of glomus cells, which reveal the true nature of the growth. In this form the glomus cells may be scanty, but characteristically lie in a row or rows parallel to the endothelium and separated from it by a well-defined fibrous layer; there is frequently straggling continuity with the glomus cells of adjacent blood spaces. A further element of these tumours is nervous tissue, and rarely this is so proliferated that the appearances are those of neurofibroma (Dupont, 1935). In the majority there is participation of all three elements and an orderly arrangement that has been interpreted as indicating the origin of this tumour in hyperplasia of the normal glomus, rather than in true neoplasia (Masson). In all types the histological structure may be complicated by focal areas of proliferation—fibromatous, neurofibromatous, or fibromyxomatous. Myxomatous degeneration in the stroma or general oedema of the growth occurs not uncommonly.

A detailed description of the typical appearances is given in our earlier publication, and the additional cases give no cause for modification; it may, however, be noted that none of our cases has shown any tendency towards the neuromatous form, despite the ample evidence of pain in the clinical histories, and that nervous elements have not been found in those showing a generalized paucivascular structure. It remains only to describe the glomus cells, since it is on the presence of these that the diagnosis rests. These large cuboidal cells have a striking and peculiarly uniform appearance; they have a circular nucleus with a well-stained reticular chromatin, a pale rim of cytoplasm, and a well-marked limiting membrane as shown by Masson's trichrome stain. There seems no doubt that this is the same type of cell as is seen in the wall of the Sucquet-Hoyer canal of the normal arteriovenous anastomosis of the skin. In places they show evident transition to smooth muscle cells, and some contain minute fuchsinophil twigs and granules which Masson considers to be the precursors of smooth muscle fibres. Only once have we noted a mitotic figure in one of these cells.

For the histologist the differential diagnosis is generally easy, although in some unusual cases with neuromatous overgrowth confirmation may be obtained, as Dupont (1935) points out, only by a careful study of the periphery of the tumour. The paucivascular form of glomangioma

may (Dupont, 1935) show some resemblance to cylindroma (sweat-gland adenoma), but we have not seen a tumour such as Dupont describes. Silver impregnation of the reticulum is a helpful diagnostic method when doubt of this type exists, since the reticulum in glomangioma is delicate where it is in relation to the epithelioid glomus cells and tends to run between them as very fine fibrils. In cylindroma, on the other hand (three cases), the reticulum is much thicker and shows a frankly peri-alveolar membrane with no tendency to fine branching or intercellular spread.

Related Cases

Since our interest was aroused in the glomangiomas we have seen two cases with highly suggestive history and clinical findings, which, however, do not show the full combination of histological features characteristic of glomangioma; both are histologically cutaneous cavernous angiomas. The first (Case XVI) shows a few cellular areas approaching the structure of capillary angioma, and in addition the presence in the fibromatous and partly myxomatous stroma of abundant myelinated nerve fibres and irregular masses of smooth muscle; no glomus cells are seen in the tumour. The other (Case XVII) shows a lesion of rather uniform type with the structure of simple cavernous angioma, situated in the dermis. The endothelium lining the blood spaces is prominent in places and is supported by bands of dense connective tissue of adult type, presenting no evidence of neural or muscular tissue. In short, the lesion shows none of the complexity characteristic of glomangioma.

CASE XVI

A woman aged 46. Medial aspect of second toe of left foot. The history begins eight years ago. The patient had an operation for removal of an ovarian cyst, and within four or five days of this she became aware of a throbbing feeling in the second toe of the left foot. The pain began as occasional short stabs in the medial side of the toe, at which time there was no abnormality to be seen. As time went on the pain came in attacks of progressively longer duration, and latterly they have been as long as two weeks, and never shorter than two days. Sometimes a few months would pass without any pain. For at least six years she has associated the pain with a small bluish nodule on the medial side of the toe, which was always extremely sensitive to trauma. When she was much on her feet or otherwise tired the pain tended to be worse. Latterly the pain has been of the type of an extreme soreness, radiating from the toe to the dorsum of the foot, giving there the sensation of millions of pins and needles. During the attacks the tumour was swollen and there was a tense prominent vessel on the dorsum of the toe. She knew of little to relieve the pain; sometimes it seemed as if elevation of the limb or warmth was of some comfort, but so slight that she begged for excision of the nodule (1932), and would have been willing to sacrifice the foot (1936). She consulted Mr. A. I. L. Maitland in October, 1938, and he removed the nodule on November 3 (1938/39, Western Infirmary, Glasgow). She was seen again on May 11, 1939. Since operation she has been free of pain. It was observed that she had, on the medial aspect of the third toe of the right foot, a small brownish symptomless nodule of rubbery consistency, which she is certain is of the same type as the original painful nodule.

CASE XVII

A young man aged 19. Ulnar border of left forearm. When the patient was 10 years old he observed the presence of a small blue nodule rising above the skin of the ulnar border of his left forearm. At times it seemed to swell, but it was some years before any pain was associated with it. The pain took the form of attacks of throbbing discomfort aggravated by heat. Latterly this local discomfort changed in character to become a feeling rather of numbness in the nodule, associated during the attacks with a tingling pain in

the fourth and fifth fingers of that hand. None of the symptoms was disabling. Attacks were usually spontaneous in origin, but could be precipitated by a sharp blow. The duration of the discomfort was from thirty minutes to a few hours. A tentative diagnosis of glomangioma was made, and the tumour was removed by one of us (W. A. M.) in April, 1937 (643/37, Western Infirmary, Glasgow). Since then the patient has had no further pain, but periodically experiences a feeling of coldness in the fourth and fifth fingers of the left hand.

Treatment

So far as our observations go these tumours are entirely benign, pursuing a slow growth which is self-limited. Treatment by complete excision is usually very easy. Many are so sharply demarcated that they will shell out, but since histological examination shows in some cases part of the tumour embedded in the collagenous dermis, it would seem preferable to remove an ellipse of skin. Despite the highly vascular nature of the growth, its removal is practically bloodless, and if the vascular pedicle be recognized one is surprised by its thinness. Local anaesthesia is perfectly satisfactory, and we have not found it to be attended with any special difficulty. In view of the uniformly good results attending excision, it is extraordinary that in many cases the latter has been withheld for years.

Comment

In view of the experience of Masson (forty-seven cases) and to a lesser extent of our own, there can be little doubt that in glomangioma we are dealing with a definite entity which is far from rare. Its actual frequency may indeed be higher than the published records suggest, since this small tumour will rarely be removed if it is producing no disturbing symptoms. One of our cases (No. XIII) was definitely painless, and similar examples have been published by Picard (1931) and by Schreus (1934).

In each of our cases the tumour was situated on a limb. Freudenthal *et al.* (1937) have cast doubt upon certain of the published cases, and especially upon those occurring elsewhere than on the hands. It is of especial interest that Grauer and Burt (1939) report two cases of glomangioma of the penile skin and submit a series of illustrations which fully establishes their claim. The histological features of all the cases in the present series fully conform with the criteria established by a critical reading of the literature at the time of our previous publication; the illustrations of the present paper are complementary to the earlier series. There is no histological difference between the tumours on the hand and those occurring elsewhere.

As mentioned above, the important constituent of these tumours is the glomus cell, which is probably a modified or specialized form of blood-vascular smooth muscle cell. Freudenthal *et al.* point out the frequent painful nature of cutaneous leiomyoma, and state: "It can hardly be a mere coincidence that the unstriped muscle cell is the mother cell of both the painful tumours." Certainly our histological study leads us to believe that the pain of the tumour is quite independent of its neural content.

In a significantly high proportion of cases the development of glomangioma has followed a single severe trauma. Reuther (1938) has suggested that the persistent pain which may follow injury to the coccyx may be related to the presence of the glomus coccygeum, and in fact may be associated with actual glomangioma formation. So far as we know this is unconfirmed.

Since Masson suggested for this tumour the term "tumour of the neuro-myo-arterial glomus" various other

names have been proposed, the most recent being "tumour of the neuro-vascular body" (Aschoff, 1938). This nomenclature seems unnecessarily cumbersome and therefore inferior to "glomangioma" (Bailey, 1935).

Summary

Fifteen cases of glomangioma are reported.

Their relation to "painful subcutaneous tubercle" is discussed.

A description is given of the striking clinical picture, and a brief discussion of the pathology.

The simplicity and efficacy of complete operative removal are emphasized.

Except where specifically designated, those named in the text are colleagues in Glasgow; to all we are grateful for their generosity and ready help. We are also indebted to Professor D. F. Cappell of Dundee, Professor Keith Inglis of Sydney, and Professor J. S. Young of Aberdeen, for their supply of material, and their friendly interest. The photomicrographs are by Mr. John Kirkpatrick, of the Department of Pathology, University of Glasgow.

REFERENCES

- Aschoff, L. (1938). *Klin Wschr.*, **17**, 1497.
 Bailey, O. T. (1935). *Amer. J. Path.*, **11**, 915.
 Bergstrand, H. (1937). *Amer. J. Cancer*, **29**, 470.
 Blumental, M. (1937). *Zbl. Haut. u. GeschlKr.*, **56**, 436.
 Dupont, A. (1935). *Bull. Soc. franç. Derm. Syph.*, **42**, 1260.
 Fingerland (1938). *Arch. Derm. Syph.*, Vienna, **177**, 255.
 Freudenthal, W., Anderson, R. G., and Weber, F. P. (1937). *Brit. J. Derm. Syph.*, **49**, 151.
 Grauer, R. C., and Burt, J. C. (1939). *J. Amer. med. Ass.*, **112**, 1806.
 Greig, D. M. (1928). *Edinb. med. J.*, **35**, 565.
 Lee, R. C. H. (1938). *Chin. med. J.*, **53**, Supplement 2, 175.
 Leriche, R. (1939). *The Surgery of Pain*, tr. and ed. by A. Young, Baillière, London.
 Mackey, W. A., and Lendrum, A. C. (1936). *Brit. J. Surg.*, **24**, 208.
 Masson, P. (1924). *Lyon chir.*, **21**, 257.
 — (1935). *Bull. Soc. franç. Derm. Syph.*, **42**, 1174.
 Picard, H. (1931). *Zbl. Chir.*, **58**, 2133.
 Reuther, T. F. (1938). Quoted by J. D. Kirshbaum and S. L. Teitelman (1939). *Arch. Pathol.*, **27**, 95.
 Schreus (1934). *Zbl. Haut- u. GeschlKr.*, **49**, 418.
 Stabins, S. J., Thornton, J. J., and Scott, W. J. M. (1937). *J. clin. Invest.*, **16**, 685.
 Touraine, A., Solente, and Renault, P. (1936). *Bull. Soc. franç. Derm. Syph.*, **43**, 736.
 Weidman, F. D., and Wise, F. (1937). *Arch. Derm. Syph.*, Chicago, **35**, 414.

In view of the separation of Burma in 1937 the triennial review of the state of the hospitals and dispensaries during 1935-7 has special interest. Financial stringency during the previous three years had led to a very close supervision of the medical department with a view to economies. Little progress or expansion was then possible, but in the succeeding triennium happier conditions prevailed, and the work of the medical department increased much, irrespective of the additional work arising from separation. The actual number of hospitals and dispensaries in Burma at the end of 1937 was only 294, and work is already proceeding to add considerably to this total. In Burma the Western methods of medical treatment are highly popular. Malaria cases numbered 469,074 in 1937, and it is remarked in the report that "the prevention and treatment of this disease, which is a menace to Burma, constitutes one of the most important as well as the most difficult tasks of the public health and medical departments." Adequate supplies of quinine were maintained in all hospitals and dispensaries. All hospitals in snake-infested areas stock anti-venine. Anti-tetanic serum is used as a prophylactic in all cases of severe injury and contaminated wounds. A tuberculosis and leprosy relief association has been formed under the patronage of the Governor of Burma, and it is hoped that a complete scheme for combating both diseases will come into operation before long. Cases of venereal disease are also common and are treated at all hospitals, some of which provide now special clinics. The establishment of a special venereal diseases block at the Rangoon General Hospital is under consideration.

A. C. LENDRUM AND W. A. MACKEY: PAINFUL SUBCUTANEOUS TUBERCLES—THE GLOMANGIOMA

FIG. 1.— $\times 8$. H. and E. Illustrating topography of tumour from Case VI, the lesion situated principally in the dermis, but extending as a lobulated nodule into underlying fat.

FIG. 2.— $\times 250$. H. and E. Silver impregnation of reticulum (Case IV). Showing abundance of reticulum and its intimate relation to glomus cells.

FIG. 3.— $\times 500$. Masson's trichrome stain (Case VII). Illustrating the typical relation of glomus cells to blood spaces in paucivascular form of glomangioma.

FIG. 4.— $\times 100$. H. and E. Showing typical arrangement of layered glomus cells in wall of cavernous space (Case VIII).

HAMILTON BAILEY AND R. J. McNEILL LOVE: ELECTROSURGICAL OBLITERATION OF THE GALL-BLADDER WITHOUT DRAINAGE (THOREK'S OPERATION)

FIG. 1.—The cystic duct and artery have been divided. The gall-bladder is being split open, and the contents evacuated into Thorek's receptacle. Packs omitted for the sake of clearness.

FIG. 2.—The portion of the gall-bladder which remains attached to the liver is electrocoagulated.

FIG. 3.—The detached round ligament of the liver, or a piece of omentum, is sutured over the gall-bladder bed.