abscess, sinuses or fistulae, foreign bodies and calculi, and neoplasms. Strictures, commonest in the bulb, appear as constrictions of either localized or elongated extent. They are very easily and very accurately defined. A consideration of added importance in the utility of urethrography in these cases is that, with long-standing strictures, complicating lesions often develop behind them. Then owing to the attendant difficulties in performing urethroscopy or cystoscopy, or to the uncritical persistence with no other treatment than that of the perennial passage of sounds, the extent of these complicating lesions may not be recognized. In these cases a urethrogram is often the only method of demonstrating the lesions and indicating the treatment necessary. Peri-urethral abscess cavities which communicate with the canal, and diverticula, sinuses, or fistulae which result therefrom, are readily outlined. Neoplasms such as carcinoma of the urethra are comparatively rare. They appear in a urethrogram as filling defects. Foreign bodies and calculi in the urethra, if opaque, would be demonstrated in the plain radiograph.

Conclusion

While the above indications of the utility of urethrography show the precision of the information which can be obtained by the technique employed, this information must of course be regarded as supplementary to the clinical, urethroscopic, and pathological investigations. Many instances occur, however, in which the data that can be obtained by the urologist by this means prove of the very greatest assistance in directing the lines of correct treatment.

GENERALIZED ACTINOMYCOSIS WITH PREDOMINANT SPINAL SYMPTOMS, INCLUDING COLLAPSE OF A VERTEBRA

BY

G. JOLY DIXON, M.A., M.B., M.R.C.P.

Medical Registrar, Charing Cross Hospital

(WITH SPECIAL PLATE)

For actinomycosis to involve the vertebrae is on the whole rare, though several well-authenticated instances are recorded. On the other hand, Parker (1923), who publishes two cases and collected six others, points out that it is even more rare for symptoms and signs obviously referable to the vertebral column to occur; in fact, in only three of his cases was the diagnosis of involvement of the vertebral column made before death. Tabb and Tucker (1933), in publishing a case, stress that when actinomycosis attacks the spine the disease affects several vertebrae, causing sharply defined areas of destruction in the x-ray picture, and that the individual vertebrae do not collapse. Such a view is also expressed by Holmes and Ruggles (1936).

In the case recorded here spinal symptoms so dominated the picture for the last six weeks that Pott's caries was very seriously considered as a possible diagnosis, and collapse of the twelfth dorsal vertebra was certainly present at the time that radiographs were taken just six weeks before death (Plate, Figs. 1 and 2). Bostroem (1890) publishes a drawing of a human spine attacked by actinomycosis which shows collapse of the eighth thoracic vertebra.

Case Record

The patient was an electrician aged 27. In September, 1938, he had an attack of lower abdominal pain accompanied by vomiting. He was able to remain at work, but the pain persisted, being accompanied by profuse sweating. After one week his doctor ordered him to bed, and found that his temperature swung between 99° F. in the morning and 100° F. at night. The pain was now chiefly in the upper abdomen. After being a week in bed he was admitted to a cottage hospital, where he remained until November 8. During the whole of this period he had an intermittent fever, his temperature swinging between 98° F. and 102° F. The pain gradually subsided. At this time a radiograph showed no abnormality of the chest.

On discharge he was still febrile, but, being free from pain, returned to work, against advice. From the middle of November, 1938, until the end of January, 1939, he remained at work, and appears to have been relatively well except for some lassitude and intermittent pains in the right side of the abdomen and back. He believed he was afebrile.

On February 6 he consulted the casualty officer of another hospital for pain in the region of the right hip and lower portion of the back. He was treated with hot fomentations to the loin for one week, and then on February 14 a mass in the right loin was incised. There was a scanty discharge for a week and the pain became much less; the patient felt generally better and the wound healed, only to break down again on March 6, with a copious discharge. On March 12 the pain suddenly returned. It appeared to radiate from the lump in the right loin up the back into the left side of the chest. The pain gripped him like a vice and seemed to prevent him breathing or moving. During the next four days the pain in the chest became gradually less severe and the breathing easier, but the pain in the back became worse.

The patient was admitted to Charing Cross Hospital on March 16. On examination the picture was dominated by excruciating pain in the back. He was unable to lie down, and enjoyed a moderate degree of relief only when propped up in a sitting position. His lower thoracic and lumbar spine was as rigid as a poker, and when moved he pushed his hands down into the bed as if in an endeavour to take the weight off his body. Over the first, second, and third lumbar spines there was a slight swelling of the subcutaneous tissues with faint reddening of the skin; the centre of this swelling was extremely tender. The patient was thin, although not emaciated; his face was flushed, his tongue very dry. There was some bulging of the left side of the chest, with the physical signs of a large left-sided pleural effusion, but little, if any, displacement of the mediastinum. About one and a half inches above the right iliac crest there was a cold swelling, in the centre of which a sinus discharged a thin serous pus. His blood pressure was 120/75, and no abnormality was demonstrated in the cardiovascular or central nervous system.

While in Charing Cross Hospital his temperature fluctuated daily between 98° F. and 103° F., his pulse ran between 96 and 114, and his respiratory rate was constantly raised slightly above normal. Antero-posterior and lateral radiographs on March 21 showed collapse of the twelfth thoracic vertebra. A few "sulphur granules" were seen from time to time in the pus discharging from the sinus in the loin, and all specimens on microscopical examination revealed the streptothrix of actinomycosis.

The chest was aspirated on April 3, 12, and 18, when thirty, twenty, and ten ounces of pus respectively were removed. On the first aspiration this was thin and serous, with sulphur-yellow granules floating in it; the subsequent specimens were opaque reddish brown in colour, again with numerous sulphur granules. On April 20 it was necessary to incise an abscess to the left of the eighth thoracic spine. Pus and air were exuded from this, and it was thought to communicate with the pleura.

From the fifth day after admission (March 20) his back became progressively less painful and more mobile, though he continued to complain of very considerable pain in the

loin. As time went on, the mobility and loss of rigidity of the spine became a most extraordinary feature. On April 28 the patient was propped up; he fell over sideways, and on being lifted back into the middle of the bed his chest flopped down on his thighs. An examination of his back at this period demonstrated that, although no regional angulation of the vertebral column could be seen, his spine between the region of the tenth thoracic and fifth lumbar vertebra was capable of forming more than five-sixteenths of the circumference of a circle—that is, he was capable of being bent forwards until the whole of the thoracic spine made an angle of 70 degrees with the long axis of the pelvis. This abnormal movement, although unpleasant to observe and not under the patient's control, caused him no pain or discomfort. A neurological examination revealed no damage to the spinal cord. The patient became progressively more emaciated, and died on May 1. It may be of interest to note that sulphapyridine in doses of 3 grammes daily was administered for fifteen days without any effect.

POST-MORTEM EXAMINATION

At necropsy, seven hours later, the appendix and caecum were observed to be bound down in the right iliac fossa by numerous filmy adhesions. At the tip of the appendix these became denser, and this organ was adherent to the right fascia transversalis. The right transversalis muscle was necrotic, and was traversed by several tortuous sinuses which discharged in the swelling in the loin. The other end of these sinuses communicated with the lumen of the gut by a perforation in the tip of the appendix. The deep fascia around the vertebral bodies from about the region of the seventh thoracic to the fifth lumbar was irregularly thickened, shaggy, and necrotic—though its substance could be seen tracking numerous sinuses. This change in the character of the fascia had no definite demarcation; the fascia immediately adjacent to the diseased area was abnormally thickened and tough, with occasional sinuses or abscesses in its substance. The upper part of the capsule of the right kidney was densely bound down to adjacent organs and showed several sinuses, though the renal parenchyma was healthy. The left psoas muscle was necrosed and partially liquified. There was a definite track of hypertrophied deep fascia honeycombed with abscesses, leading to the parietal wall of the left pleura, which latter was thickened and shaggy. In the left pleural sac there was from ten to fifteen ounces of mulberry-red opaque creamy fluid in which floated numerous sulphur-coloured granules. The left lung was collapsed, shrunken, and rather tough to cut, though no obvious disease could be seen in its parenchyma. There was a discharging abscess whose cavity surrounded the broken neck of the eighth left rib. Though the fascia around this was hypertrophied and honeycombed with small sinuses, no obvious communication could be demonstrated with the pleura.

When the diseased fascia was stripped off the vertebral bodies these were seen to be necrosed and eroded in such a manner that it gave them a coarse sandy granular appearance. Apart from the twelfth thoracic, no vertebra had lost shape, and all remained rigid. The twelfth thoracic vertebra was partially collapsed and in the intervertebral disk below was a large cavity about twice the size of a pea (Fig. 3). The excessive mobility of the spine appeared to be due to the laxity of the ligaments and intervertebral substance rather than any change in the vertebra.

On cutting the vertebral column in half, abscess cavities were seen in the body of the twelfth thoracic vertebra, and there was a small one in the second lumbar vertebra (Fig. 4). The proliferative suppurative process had invaded the tissues around the dura, and a mass of necrotic material was pressing upon the spinal cord opposite the body of the twelfth thoracic vertebra.

Summary of Case

September and November, 1938.—Four weeks' lower abdominal pain accompanied by high remittent fever. Partial recovery; returned to work, probably afebrile.

February, 1939: pain over right hip. Cold abscess developed in right loin; incised scanty discharge; nearly healed after three weeks, only to break down again, afebrile.

March 12.—Sudden exacerbation of symptoms; left-sided pleurisy followed by empyema, severe pain and rigidity in thoracic spine, and high remittent fever. Radiograph showed collapse of twelfth thoracic vertebra.

Gradual relief of pain in back with increased mobility of spine; this latter became extremely abnormal. Remittent fever continued; patient wasted away and died on May 1.

At Necropsy.—Left-sided empyema; extensive involvement of fascia around lower thoracic and upper lumbar vertebrae; abscess cavities in twelfth thoracic and second lumbar vertebrae. Collapse of twelfth thoracic; necrosis of right transversis abdominalis, and left psoas muscles. Disease appeared to originate from the appendix.

I have to thank Dr. Gordon Holmes, who had this case under his care, for permission to publish these findings.

REFERENCES

- Bostroem (1890). *Beitr. Path. Anat.*, **9**, 1.
Holmes, G. W., and Ruggles, H. E. (1936). *Roentgen Interpretation*, Kimpton, London.
Parker, C. A. (1923). *J. Bone Jt. Surg.*, **5**, 759.
Tabb, J. L., and Tucker, J. T. (1933). *Amer. J. Roentgen*, **29**, 628.

Clinical Memoranda

A Treatment for Phagedaena (Tropical Ulcer)

Like most medical men in the Tropics I have tried various treatments for the rapidly sloughing ulcer, recognized from near or far by its smell, and called phagedaenic ulcer or *ulcus tropicus*. Any form of excision short of amputation is not only insufficient as a rule to stop the burrowing process locally, but is also liable to be followed by rapid increase of the general septicaemic process and death. Cauterization with neat carbolic acid seemed to have the same results, though I remember getting some satisfaction from one of these processes—done under anaesthesia—when I placed a bag of common salt in the resulting wound to produce an exaggerated flow of serum. The rationale of this must be largely that the causative organism finds it difficult to swim or to multiply against the stream.

For the last dozen years, however, I have used one treatment almost exclusively. It was quoted to me from a now unknown source by a non-medical colleague. So far as I know it has only once appeared in print, and then in a booklet for laymen, my *First Aid Supplies* (U.S.C.L., 1939). I call this treatment P.P.C. (potassium permanganate cautery).

METHOD OF TREATMENT

A 5 per cent. permanganate solution is used. It should be freshly made up, but this creates no difficulties, for 5 per cent. is full concentration, and so long as a few crystals remain undissolved the strength is right. A few crystals are put into a small vessel, such as a bottle cap or gallipot, and covered with a little water. When, a few minutes later, a pledget of cotton-wool is dipped into this, the fact of full concentration is established by finding crystals adhering to it. This pledget is pressed first against a larger piece of wool to remove excess

G. JOLY DIXON: GENERALIZED ACTINOMYCOSIS WITH PREDOMINANT SPINAL SYMPTOMS, INCLUDING COLLAPSE OF A VERTEBRA

FIG. 1.—Antero-posterior view of spine.

FIG. 2.—Lateral view of spine.

FIG. 3.—Spine from eleventh dorsal to second lumbar vertebra.

FIG. 4.—Section of spine from eleventh dorsal to second lumbar vertebra.