BRITISH MEDICAL JOURNAL

LONDON SATURDAY OCTOBER 28 1939

PROPHYLACTIC IMMUNIZATION AGAINST MEASLES, SCARLET FEVER, DIPHTHERIA, WHOOPING-COUGH, AND INFLUENZA *

BY

SIR J. C. G. LEDINGHAM, LL.D., D.Sc., F.R.C.P., F.R.S.

Director, Lister Institute, London; Professor of Bacteriology, University of London

The prevention of infectious disease by active immunization methods is no new thing, and I am going to assume that what is chiefly wanted at the moment is some reassertion and fortification of the scientific background of these procedures and some account of their achievements in the field, with a view to rendering them more acceptable to those responsible for the conservation of child life in this country and thus to facilitate much-desired propaganda on the large scale. Whatever may be the reasons—and I hope to suggest some later on—there can be no doubt, I think, that in the field of prophylaxis of the kind we are considering to-day this country has in the main been a follower, and, I'm afraid, rather a tardy follower; for, though our laboratory workers have contributed very materially to the knowledge of methods, we have left to other countries the task of trying them out seriously and intensively in the field.

The safety of any method to be applied to man must always be a prime consideration. There must be no risk to life, and even the minor inconveniences, local and general, incidental to the introduction of a prophylactic through the skin must also, so far as possible, be reduced to a minimum consonant with the effectiveness of the product. In the course of securing these desiderata—and I refer here particularly to the diphtheria prophylactics—there have been in the early period of their use, though not in this country, some very few unfortunate experiences that could not have been foreseen at the time but which further experiment explained and taught us to avoid in future. With the introduction of the Therapeutic Substances Act, however, such accidents should never recur if the prescribed rules for the preparation and testing of prophylactics for issue are carefully carried out, and also if those responsible for administering them perform their duty intelligently. We have now the definite assurance that the prophylactics we are here considering are both safe and potent, and, when administered with the care and circumspection that proceed from knowledge, are calculated to cause only quite negligible inconvenience to the subject inoculated.

Diphtheria

Of the five infectious diseases with whose prevention by prophylactic inoculation I am dealing, diphtheria undoubtedly merits chief consideration because we now

know quite definitely from field trials in other countries that intelligent and persistent immunization of the child population, starting with the pre-school child one year old, can bring the diphtheria morbidity down to zero. Indeed, as that great pioneer in this field the late Dr. W. H. Park remarked at the International Congress for Microbiology in London in 1936, "The control of diphtheria by specific treatment and specific prophylactic immunization is one of the most gratifying victories of bacteriology." Let me just enumerate some of the landmarks in the history of this matter.

In 1889 Roux and Yersin demonstrated that the diphtheria bacillus elaborated a toxin, and that to this toxin was due the symptom-complex of diphtheria. This led in the first instance to Behring and Kitasato's great discovery of antitoxin in 1890 and to the treatment of diphtheria by passive immunization methods still current and without a rival. The use of diphtheria toxin for preventive purposes was also early foreshadowed, but many years had to pass before the time became ripe for exploring the possibilities of active immunization in diphtheria control schemes. The Schick test, introduced in 1913, gave us most valuable information regarding natural or naturally acquired immunity to diphtheria at various age groups, and now that we have that knowledge we can to a large extent dispense with it, particularly in dealing with young children. It still, however, has its uses as a guide to prophylactic immunization in the older age groups, in which the proportion of Schick-positive persons steadily diminishes owing to increasing opportunities of acquiring immunity through, it is believed, mild or unrecognized infections contracted from their surroundings.

Then came the era of preventive control by partially neutralized mixtures of toxin and antitoxin. Behring introduced these for human use in 1913, and Park and his colleague Zingher exploited them very successfully in New York during the period 1915-23. These partially neutralized mixtures were not ideal immunizing agents, as they were of doubtful stability and their safety was not easily controlled owing to dissociation phenomena under different conditions of storage which I need not go into. They left much to be desired for acceptable propaganda purposes, but nevertheless when administered by good hands and prepared with the necessary care they did excellent service and showed conclusively what could be done in this way to transform a condition of susceptibility into one of more or less solid immunity.

* Read in opening a discussion in the Section of Medicine at the Annual Meeting of the British Medical Association, Aberdeen, 1939.

The Introduction of Formol-toxoid

The next notable advance was the introduction of modified toxin or toxoid for prophylactic use. Glenny and Südmersen in 1921 recorded the fact that toxin which had been converted into an atoxic product by contact with formalin was capable of immunizing guinea-pigs against diphtheria toxin. Ramon of the Pasteur Institute who in 1922 published the first of an important series of papers on the flocculation that took place in neutral mixtures of toxin and antitoxin, and thereby provided immunologists with an *in-vitro* method of titration, extended his observations to formol-toxoid, or, as he called it, "anatoxine," and found this to behave similarly, showing that its power of combining with antitoxin was unimpaired. In 1924, we are told, he inoculated himself with "anatoxine," and in subsequent years he did much to popularize its use in France and other countries, particularly America. In an address on world experience of formol-toxoid at the jubilee celebrations of the Pasteur Institute last March, Ramon stated that more than three millions have now been vaccinated in France, while in Canada, of a population of ten millions, more than two millions have been immunized. The new antigen was immediately exploited by Park in New York, and by 1935 one million had been immunized in that city. In Paris since 1928 some 50 per cent. of Parisian children have been immunized, and deaths from diphtheria fell from 318 in 1928 to seventy-two in 1938. In December, 1931, it became obligatory to immunize the French soldier against diphtheria and enteric by the use of a mixture of "anatoxine" and T.A.B. vaccine, and since that time tetanus toxoid has been added to the mixture. By the end of 1938 some 800,000 troops had been so vaccinated, and Ramon was able to state that clinical diphtheria was disappearing from the French army. He bemoaned the fact, however, that civilian France, which depended so much on the conservation of its child population, was still too apathetic in the matter of diphtheria immunization, and he looked forward to this procedure becoming obligatory in the civil population. This, I understand, is now the case.

The most arresting figures are gleaned from Canada, where efforts to popularize the immunization of children, on whom diphtheria exerts its worst toll, have been most intense and persistent. Some towns and rural districts in Canada have 70 to 90 per cent. of the child population vaccinated. The experience of the city of Hamilton (150,000 population), which was the first city of this size to attempt to control diphtheria with formol-toxoid, is widely quoted, for no deaths from diphtheria have occurred there since 1930 and no case of diphtheria since 1933. In Toronto, with a population of 650,000, some 104,000 infants have been vaccinated since 1926, and the diphtheria incidence fell in the four years 1930-4 from 164 to 3.5 per 100,000. In 1936 there were only forty-eight cases and two deaths. In New York in 1936 some 75 per cent. of the infant population had been vaccinated. Round about 1924 this city had annually some 10,000 cases of diphtheria with 700 deaths, but in 1936 the cases had fallen to 1,143 and the deaths to thirty-five (or about 0.47 per 100,000 inhabitants. In 1938 the reported cases fell to only 163 in this great city.

It is only when the proportion of vaccinated children reaches 70 or 80 per cent. that, as Ramon says, one sees the real fruits of one's work in a great reduction of the diphtheria incidence, perhaps to zero. We have nothing like this to show in Great Britain. In trials hitherto

made in this country the value of prophylactic vaccination against diphtheria can be measured only by the relative immunity of the vaccinated groups as compared with that of the unimmunized. These data are arresting enough, so arresting indeed as to render urgently desirable the immunization of all pre-school children from the age of one year, with a view to the ultimate extinction of this disease from the community. No other method of diphtheria control can offer such prospects. What is the problem in this country?

Our Need of More Intensive Propaganda

The chief medical officers of the Ministry of Health and of the Scottish Board of Health both emphasize the necessity for more intensive propaganda, for the diphtheria incidence remains high and almost stationary. In England and Wales there have been annually some 60,000 to 70,000 cases of diphtheria, with some 3,000 to 4,000 deaths, and in Scotland some 10,000 cases, with 400 to 500 deaths. Sir Arthur MacNalty in his report for the year 1937 informs us that Chester, with 45 per cent. of its children vaccinated, "is probably the best-immunized town in the country." After it come Birmingham, Walsall, Worcester, Leeds, Manchester, and Chatham, "all of which are in the neighbourhood of 35 to 40 per cent." London occupies an unenviable position at the bottom of the list with only some 5.3 per cent. of its child population immunized.

To illustrate what can be done by less intensive or, more correctly perhaps, less effective propaganda so far practised in this country, I may cite briefly from current reports the experiences of the city and county of Aberdeen and of the city of Birmingham. During the period 1935-7 some 16,000 of the 25,000 children on the school rolls in the county of Aberdeen have been immunized; together with some 3,000 to 4,000 pre-school children, the antigen employed being alum-precipitated toxoid (A.P.T.), given by what has come to be known as the one-shot method. It is interesting to note that the parents' consent was obtained for about 80 per cent. of the children attending the county schools, a result most creditable to the intelligence of the Aberdeenshire parent. What has been the effect of this crusade? Among the immunized children there have been since March, 1935, seventy-three notifications of diphtheria, giving an attack rate of 26 per 10,000 children, while among the unimmunized, including some 3,000 known Schick-negatives, the attack rate was 251 per 10,000, or ten times that among the immunized. In the city of Aberdeen active immunization of school children and those of pre-school age began in September, 1936. Unfortunately, parents' consent was difficult to obtain, and only 52.3 per cent. of new school entrants were immunized. During the period September, 1936, to March, 1938, 3,596 children were immunized, of whom fifteen contracted diphtheria, giving an attack rate of 0.4 per cent., whereas among the non-immunized the attack rate was 1.5 per cent., or nearly four times that among the immunized. There were no deaths among the immunized, and the disease was mild.

In the city of Birmingham an immunization crusade was started in 1925, the agent used being toxoid-antitoxin mixture (T.A.M.) in three doses of 1 c.cm. at fortnightly intervals. Those completely inoculated under the age of 15 in 1938 numbered 96,446, or some 42 per cent. of the total population in that group. As for the pre-school group under 5, about 27 per cent. had been immunized by the end of 1938. The total cases of diphtheria in the last five years have varied from 1,019 to 1,359, and the deaths from sixty-three to eighty-four annually. Only

fifty-six cases in all have occurred among the inoculated children, and there have been no deaths.

It is quite clear from these illustrations that it will be necessary to increase greatly the percentage of those immunized before evidence is obtained of any material fall in the general diphtheria incidence.

The Antigens and their Administration

Now let us consider as briefly as possible the problem of antigens and schemes of administration appropriate to each. The Chief Medical Officer of the Ministry of Health remarked in a reference to this subject in his report for the year 1937: "It is unfortunate that the exact technique, preparation of vaccine, and method of administration are still matters of medical controversy. It is probable that the lack of a general standard has militated against the more general adoption of anti-diphtheritic immunization in this country." I would consider this a rather unfortunate statement. Though admittedly there is no general standard, the lack of one has been an advantage rather than otherwise, for it has stimulated valuable comparative field trials, and for several years now we have learnt to know more than one type of antigen which, when administered appropriately, is capable of reducing diphtheria incidence materially. Progress has been impeded by many factors, and not least by what a recent leader writer in the *Journal* calls that "doltish prejudice," not confined to any one social or intellectual stratum, which is fostered by anti-social societies posing as the protagonists of liberty. When liberty becomes anti-social and impedes the application of scientific truth to the improvement of human well-being then I think some form of compulsion is the only remedy, though one might hope it would be merely temporary, pending the wider diffusion of scientific knowledge in the community. I might add that I should have liked to see a far more encouraging lead on the part of health authorities generally, from the Ministry of Health downwards, while giving all credit to those enlightened health authorities who have tried their best.

We have at least four antigens which we know to be potent—namely, F.T., T.A.F., A.P.T., and T.A.M. F.T., or formol-toxoid, has done marvellous service in France, Canada, and the United States, though in the latter country A.P.T. is now largely used, while T.A.F., or toxin-antitoxin floccules, A.P.T. (alum-precipitated toxoid), and T.A.M. (toxoid-antitoxin mixture) have been more popular in this country. The general tolerability of these antigens undoubtedly differs, although possibly some workers with large experience have laid more stress than others on the expediency of guarding against unusual reactions during immunization by performing preliminary detector tests. The Canadian workers who have used F.T. for the most part emphasize the value of Moloney tests, particularly in older children and adults, while in France the matter of unusual reactions with the same antigen has not been greatly stressed.

T.A.F. and T.A.M. are relatively innocuous at all ages, while A.P.T. is well borne in young children. In older children and adults it may give rise to painless nodes, which take some time to disappear. A.P.T., by virtue of the slowness of absorption of the antigen and consequent longer stimulus to the antibody-forming mechanism, can give rise, as Glenny, who introduced it, showed, to a considerably higher antitoxin titre, given in a single dose, than any of the other antigens. For this reason it may well become the antigen of choice for young children, the group we wish above all to see in course of time fully

immunized, and there is now a consensus of opinion that no countenance should be given to the use of this excellent antigen in any one-shot method.

Antigenic Values Compared

Fitzgerald and his colleagues have tested on batches of children the comparative value of three doses of F.T., two doses of F.T., one dose of A.P.T., and two doses of A.P.T., and have concluded from the results of actual antitoxin titrations that three doses of F.T. give a superior, but not a greatly superior, immunity to that given by two doses of A.P.T. Two doses of F.T. or one dose of A.P.T. yielded by comparison very poor results. On the other hand Volk and Bunney (1939) found that two doses of A.P.T. given at three weeks' interval were superior to three doses of F.T. at similar intervals.

There is no doubt that it is highly expedient that comparative trials should still go on, for it is desirable in planning extensive crusades to determine what form of antigen and what scheme of administration are best calculated to yield the most lasting immunity, due consideration being always given to the tolerability of the antigen or antigens in the interest of acceptable propaganda. On this point Volk and Bunney make a very pertinent remark: "The ultimate basis," they say, "for evaluating a particular scheme of immunization is whether it protects against diphtheria or not, and not whether it confers Schick-negativity or raises the antitoxin level to any definite point." Further, they remark that, "It is entirely possible that a procedure which results in a lower level of antitoxin than some other may still be the method of choice from the public health standpoint—that is, it might be less expensive, easier to administer, and confer a high enough percentage of immunity to be the most efficient use of the public health dollar in prevention of diphtheria."

In an important study by Jones (1938) the comparative value of F.T. and A.P.T. was tested on guinea-pigs, which after immunization were bled every four weeks for a period of thirty-two weeks, their sera being then tested for antitoxic value. This study emphasized the importance of lengthening the interval between the first and the second dose in a two-dose system. He found A.P.T. to be an excellent primary immunizing agent, and he suggested that if an adequate interval elapsed between the first and the second dose it mattered little which type of toxoid was used for the second or later injections. Whether F.T. or A.P.T. was used, it was important that the dose should contain at least 15 Lf. The interval between doses should in his opinion be prolonged to two months or even longer. It may be noted that in the immunization of horses against tetanus or diphtheria toxins the importance of a lengthy interval between the primary and secondary stimuli is well recognized.

I may here summarize for purposes of discussion the main points on which there is already fairly general agreement:

(1) A single immunizing dose of any antigen is not effective if high and lasting immunity is desired, and, where still used in certain localities, should be abandoned. (2) A.P.T. is *per se* a very significantly more effective reagent than F.T., and if the scheme adopted involved two doses only, then two doses of A.P.T. would give higher levels of antitoxin than two doses of F.T. (3) Any one of the three methods in common use—namely, three doses of T.A.F., three of F.T., or two of A.P.T.—will greatly reduce the diphtheria incidence among the inoculated. Further, as A.P.T. is the most effective antigen and is easily tolerated in children under 8, it may be recommended for the primary stimulus and followed at

not less than one month's interval by two doses either of T.A.F. or F.T. Whether to stick to the same antigen throughout is a matter for discussion. (4) With the knowledge that any one of these systems can be guaranteed to reduce incidence, local health authorities contemplating crusades can still provide useful comparative information by employing two different schemes, probably in alternate children. (5) There is general agreement that longer intervals between doses, and specially between the first and the second dose, are calculated to yield superior results. (6) At the start of a crusade in any community children from infancy to school-leaving age should be immunized in the largest possible numbers. (7) As time goes on immunization would be confined to the pre-school children of 1 to 2 years, and when such children go to school it may be very desirable to give a single reinforcing injection. (8) In children over 10, T.A.F. is to be preferred to F.T. or A.P.T. unless preliminary detector tests are performed. (9) Not until some 80 per cent. of the child population under 15 have been immunized will a material decrease in the incidence rate of diphtheria be realized.

Whooping-cough

I pass now to whooping-cough, which, with a death rate per 1,000 population of 0.043 (England and Wales, 1937), comes next to diphtheria, with a corresponding rate of 0.072. In England and Wales 1,750 deaths from this disease were recorded in 1937, practically all in children under 15. In Scotland in 1938 some 6,000 cases were reported, with 111 deaths of children under 1 year and 105 in the age group 1 to 5. As a killing disease of infancy, therefore, whooping-cough closely rivals diphtheria, and undoubtedly its prevention or control by active immunization methods is a challenge to us all. The prospects of such control, in so far as we may judge from results of crusades of quite recent years, particularly in the United States, are now certainly bright, and there is every reason why health authorities in this country should promptly organize immunization schemes on sound lines, with a good prospect of reducing the incidence of whooping-cough among vaccinated children definitely exposed to this highly contagious and distressing infection.

The aetiological agent of whooping-cough is recognized to be the *Haemophilus pertussis*, originally discovered by Bordet and Gengou in 1906. In recent years valuable bacteriological work has been accomplished in identifying this organism in the early catarrhal stages, when it may be found in some 80 per cent. of cases by the cough-plate method, and also in diminishing percentages as the weeks pass. It is a fortunate thing that the organism is a single antigenic type, for it makes immunization by vaccine far more practicable; but it is also most important for immunization purposes that only smooth freshly isolated and virulent strains be chosen, as artificial passage of the organism on unsuitable media is accompanied by degradation of the efficient antigen and by loss of virulence. It is highly probable that failure to choose strains of full antigenicity for vaccine purposes accounted for conflicting results in some of the earlier trials in the past decade, and that there has also been a failure to follow up carefully the degrees of exposure to this disease not only of the vaccinated but also of the control groups. I propose, therefore, to sketch one or two only of some quite recently recorded crusades which were embarked upon with great care and carried out in the most thorough and painstaking manner.

Studies on Whooping-cough Immunization

Kendrick and Eldering's study (1939) was made in the city of Grand Rapids, Michigan, with a population of about 168,650.

1,815 children between 8 months and 6 years were fully vaccinated with thoroughly tested strains from cough-plates, seventy-two-hour growths being washed off the medium, filtered through cotton-wool, and then centrifuged. The deposit was suspended in saline containing 1 in 10,000 merthiolate or 0.5 per cent. phenol and stored till sterile. It was standardized to 10,000 millions per c.cm., and the standard scheme was 4 doses at weekly intervals of 1 c.cm., 1.5 c.cm., 1.5 c.cm., and 3 c.cm. Some local soreness and induration resulted, but systemic reactions were slight or absent. The controls numbered 2,397, and observation was made over a period of forty-four months—March 1, 1934, to November 1, 1937. These 4,212 children were at risk for an average period of 13 months per child, 15 months in the vaccinated group and 11.6 months in the control group.

For any number of vaccinated children in a district a like number of controls were chosen in the same area, and as providing evidence of equality of exposure, both groups, we are told, had a very similar experience with such diseases as measles and scarlet fever.

The follow-up entailed 23,961 visits, or 5.7 visits per child at an average interval of 2.3 months. In 69 per cent. of the children in the vaccinated group, cough-plates were made in accepting the diagnosis of pertussis, and also in 67 per cent. of those in the control group. It is of considerable interest that, in those of the vaccinated group who contracted pertussis, cough-plates were positive in 6 per cent. as against 24 per cent. in the control group.

The results were impressive: 400 attacks occurred in all—namely, fifty-two in the vaccinated group and 348 in the control group. As to the type of attack, "severe" cases formed only 3.8 per cent. of the total in the vaccinated group. In the control group they constituted 13.1 per cent.

Statistically investigated in relation to exposure, the results showed that 12.8 per cent. of exposures in the vaccinated group were followed by pertussis, as compared with 68.5 per cent. in the control group. I may add that this investigation entailed the services not only of the Michigan Board of Health with its Bureau of Laboratories but also of the City Health Department of Grand Rapids, with its health officer and school physician. Local doctors helped in the immunization clinics, and assistance was given by nurses from the Bureau of Public Health Nursing of the City Health Department.

As a corollary to this field trial I would mention a most interesting experiment by Lawson (1939), who attempted to evaluate in mice the prophylactic value of various pertussis vaccines prepared from smooth, rough, and intermediate strains, the test of immunity being the capacity of the immunized mice to withstand an intratracheal injection of 0.05 c.cm. of a virulent culture calculated in the controls to produce a pertussis pneumonia. The results showed great differences in the immunity following vaccination with the various types: 92.6 per cent. survivals occurred among those vaccinated with the smooth strains, while those immunized with the intermediate and rough strains had only 30.8 per cent. and 7.1 per cent. survivals respectively.

One final study reported by Miller and Faber (1939) is worthy of note. This study on whooping-cough immunization was made at a baby clinic in California. Every month a number of infants less than 1 year old were vaccinated with pertussis vaccine given in three doses of 20,000 millions, 30,000 millions, and 30,000 millions at weekly intervals, while a similar number were selected by chance as controls. The vaccine was made from smooth Phase I organisms grown on Bordet-Gengou medium with human blood added. In all, 211 children were vaccinated and 182 acted as controls. All exposures

were carefully noted, and throughout an observation period of two and a half years there occurred among the test group twenty-six exposures which were followed by nineteen escapes, and among the controls twenty-five exposures with three escapes. The "communicability" rate among the vaccinated was therefore only 26.9 per cent., as against 88 per cent. among the controls.

Apart from the results achieved in these trials, which I can only sketch here, the mere reading of the records gives one a lively impression of the courage of the organizers and their determination to probe the possibilities of a method of prophylaxis against whooping-cough in spite of early failures by others, and to endeavour to find out to what factors, whether concerned with the vaccine and its administration or with the interpretation of the resulting data, such failures were most probably due. There is no doubt that a correct evaluation of the method can be made by an organization capable of observing as precisely as possible the exposures of the test and the control subjects and noting the proportion of escapes in the two groups.

Another problem arises here. The age at which a pre-school child, say 9 months, may be submitted to immunization against diphtheria is also the age at which this same child might very desirably be immunized against whooping-cough. What is to be done? I am not aware that diphtheria toxoid has so far been administered in combination with a pertussis vaccine, but I see no reason why this should not be attempted if Ramon's claim holds that the potency of diphtheria toxoid in mixture with T.A.B. vaccine is in no way interfered with and indeed appears to be enhanced. Such mixtures would first require to be tested on animals in which, at any rate, the rise in antitoxin titre can be accurately evaluated.

Before, however, we can embark seriously on well-planned crusades to prevent diphtheria and whooping-cough, one great stumbling-block has to be got rid of, and it is not the stumbling-block of general apathy. It is the stumbling-block occasioned by the fact that many large health authorities either cannot afford or are not sufficiently alive to the necessity for trained whole-time bacteriologists on their staffs capable of organizing and co-operating in such crusades. It is a short-sighted policy to which the Ministry of Health might well direct its attention, because, if persisted in, we cannot hope to keep pace in this country with the progress in preventive medicine registered elsewhere.

Scarlet Fever

With regard to scarlet fever, I think it is fair to say that even if a thoroughly dependable scheme of vaccination with the appropriate streptococcal toxin, the strength of which is measured in skin-test doses, could be recommended and put in practice, the mass immunization of children would not appear to be particularly urgent in view of the benign character of the malady in these days. In the United States the position is similar, and no active immunization against scarlet fever seems to be taking place, though both here and in that country much has been and still can be successfully done to protect nursing staffs and others in institutions. This limited application of the method of active prophylaxis has undoubtedly resulted in a reduced incidence of scarlet fever. A few months ago the Surgeon-General reported to the Office Internationale in Paris that active immunization against scarlet fever had made little progress in the United States, and he gave as reasons: (1) the benign character of the malady; (2) indecision as to the practical value of the

method; and (3) the limitations imposed on production of the antigen by holders of the Dick patent. The decision of the Ministry of Health not to give its blessing to the practice of prophylactic immunization against scarlet fever, at least under present circumstances, would seem to be justified, and I do not propose to say anything more about it.

Measles

The possibility of preventing measles or of mitigating its attack by passive immunization of contacts with convalescent measles serum or, it may be, of adult serum, has been recognized for over twenty years, and the method has been practised extensively, but less perhaps with the object of preventing attack than of moderating possible attacks on weakly children who happen to be contacts. A considerable amount of success has been achieved in both directions, but there has been no widespread effort to secure and store the human sera necessary for more than a small proportion of contacts, so that, as a measure calculated to reduce epidemic incidence, it plays only a very minor part. It is unnecessary for me to enlarge on the technique and practice of the method, and I shall therefore content myself with referring briefly to certain difficulties and problems that have arisen in connexion with large-scale experience. In this country the London County Council has during the past decade devoted much attention to measles prophylaxis, and has published elaborate accounts and analyses of its experience both with convalescent and with adult sera. Gunn (1938) in a recent account of experiences during the London measles epidemic (1935-6) draws attention to the fact that there is no method of assaying beforehand the potency of a particular batch of human serum. It is probable, as he says, and as we know for certain in the case of convalescent poliomyelitis serum, that potency may vary greatly and may account for disappointing and anomalous results. Measles prophylaxis by the passive method cannot in his view be greatly extended in scope until immune serum is readily available at all times when required and until deliberate exposure is practised so that non-immunes may have the advantage of attenuation. This, as he says, can be justified only if the efficacy of the serum is proved beforehand and parents or guardians give their consent.

Dr. Gunn raises the question whether the possibilities of active prophylaxis, or of combined active and passive prophylaxis, may not reasonably be exploited further. To use for purposes of active prophylaxis infective material from measles in the early catarrhal stage, as some have on occasion done, would, I am afraid, be a dubious and impracticable expedient. We must wait for further knowledge of the infective agent of measles, and it is slow in coming. The stumbling-block in experimental research on measles has been the difficulty of transmitting the disease with certainty to any laboratory animal and carrying it on in passage. If this could be achieved we would be in a position not only to assay batches of convalescent or adult serum but also to explore the possibilities of active prophylaxis.

Influenza

Influenza, and by that I mean more specifically influenza as we know it in epidemic times, is now recognized to be a virus disease communicable to more than one laboratory animal and capable of being passaged with regularity in such, as well as in the embryo chick membranes. We are therefore in an excellent position to explore in man the possibilities of active prophylaxis, and

extensive research to this end is in progress in this and other countries. The position at the moment has been clearly stated by Andrewes (1939) in a recent paper. We now know that active or "living" virus when introduced subcutaneously or intraperitoneally in large doses can protect ferrets from contracting infection by contact and also inhibit the development of lung lesions, but that it does not usually protect them against a test dose of virus administered by the nasal route. When the inactivated virus is introduced subcutaneously in ferrets and mice it will immunize, though less effectively than the "living" virus. Fairbrother and Martin (1939) have shown that elementary-body suspensions obtained from pneumonic lungs of mice, when heated at 57° C., are highly satisfactory immunizing agents for mice when tested for immunity by the nasal route. Both "living" virus and inactivated virus, given subcutaneously to man, induce the development of specific virus-neutralizing antibodies in the blood serum.

One finding, however, which is likely to complicate the problem of active immunization as applied to man is the multiplicity of antigenic types among the strains recovered from different outbreaks of human influenza. These types do not differ so sharply from each other as do the pneumococcus types, but there would appear to be at least four main types, together with others, or master-strains, which contain in themselves most of the major antigens and react, therefore, with the immune sera evoked by all strains isolated from human influenza. The problem raised by this complication, and its bearing on immunization of man as a practical issue, has been dealt with by Dr. Stuart-Harris, who has had especial experience in this sphere of work. Fairbrother and his colleagues claim that these antigenic differences, which are brought to light by the use of the virus-neutralization test, cannot be detected when the complement-fixation test is used, and they suggest that the basic antigen may possibly be identical in all strains. Doubtless research on this problem will be continued with the intensity its importance demands, and we may reasonably hope that some really useful addition to our armamentarium against influenza may emerge.

Conclusion

To conclude, I have written almost entirely from the standpoint of one who has had little opportunity of observing at first hand the difficulties inevitably encountered in attempting to apply effectively results of laboratory research that can be guaranteed to reduce the incidence and mortality of certain of the diseases here discussed. I can appreciate these difficulties, however, and can only regret that what may be called the social impact of medical discovery in the field of prevention takes so long to fructify. In the field of treatment discoveries get across quickly enough, perhaps too quickly. When I reflect that, owing to ignorance, vested interest, or complacency in high places, a municipality is still prevented from ordering the pasteurization of all milk that comes into it, I sometimes despair of getting preventive science across. If a move was made to make diphtheria immunization compulsory, as it is now in Guernsey, we would be told, I suppose, that the Government cannot move in advance of public opinion. In reply I would say that the present time of national stress is rather a favourable one in which to change that policy, and to impose on people what is good for them if they are unable to realize what is good for themselves and their children. It may sound a totalitarian policy, if you like, but in essence it seems to me quite democratic, for

it involves a negligible sacrifice on the part of the individual for his own and the common good.

Summary

A survey is made of the position with regard to control of the five infectious diseases by preventive inoculation.

The various diphtheria antigens are reviewed, together with the methods of administration appropriate to each.

The necessity for more intensive propaganda in favour of immunization against diphtheria is urged.

The subject of active immunization against whooping-cough is discussed, and studies of recent crusades abroad are given.

Prophylactic immunization against measles, scarlet fever, and influenza is briefly dealt with.

BIBLIOGRAPHY

- Andrewes, C. H. (1939). *Lancet*, **1**, 589.
 Fairbrother, R. W., and Martin, A. E. (1939). *Arch. ges. Virusforsch.*, **1**, 114.
 FitzGerald, J. G., Fraser, D. T., McKinnon, N. E., and Ross, Mary A. (1938). *Lancet*, **1**, 391.
 Glenny, A. T., and Südmersen, H. J. (1921-2). *J. Hyg.*, **20**, 176.
 — Pope, C. G., Waddington, H., and Wallace, U. (1926). *J. Path. Bact.*, **29**, 31, Note 23.
 Gunn, W. (1937-8). *Proc. roy. Soc. Med.*, **31**, 828.
 Jones, F. G. (1938). *Amer. J. Hyg.*, **28**, 359.
 Kendrick, P., and Eldering, G. (1939). *Ibid.*, Sect. B., **29**, 133.
 Lawson, G. McL. (1939). *Ibid.*, Sect. B., **29**, 119.
 Miller, J. J., jun., and Faber, H. K. (1939). *J. Amer. med. Ass.*, **112**, 1145.
 Ramon, G. (1922). *C. r. Soc. Biol.*, **86**, 661, 813.
 — (1924). *Ann. Inst. Pasteur*, **38**, 1.
 — (1939). *Int. Clin.*, **1**, Ser. 2, 241.
 — (1939). *Presse méd.*, **47**, 405.
 Volk, V. K., and Bunney, W. E. (1939). *Amer. J. publ. Hlth.*, **29**, 197.

OPHTHALMIC SERVICES TO THE CIVIL POPULATION IN NATIONAL EMERGENCY*

BY

Colonel A. IRVINE-FORTESCUE, D.S.O., M.B.

Special Certificate Civilian Anti-gas School, Easingwold

The subject of ophthalmic service in national emergency is a very wide one, and I propose to restrict my paper to an attempt to describe the conditions under which ophthalmologists will have to work in such an emergency. We have all heard many lectures and read numerous articles on conditions in Britain during the next war. Much that has been spoken and written on the difficulties we shall have to face in the event of hostilities strikes one as academic and unreal because of the lack of concrete examples and of descriptions of actual events.

The civil war which broke out in Spain in July, 1936, has afforded most interesting and instructive lessons in the effects of air raids on large cities. Since over 80 per cent. of the population of Britain is urban and approximately one-fifth of our people inhabit Greater London, it is obvious that the problem of defence against enemy attack from the air is a matter which chiefly concerns the town dweller. The city of Barcelona, with a population approximately equal to that of Glasgow, was frequently bombed from the air during 1937 and 1938. The

* Read in opening a discussion in the Section of Ophthalmology at the Annual Meeting of the British Medical Association, Aberdeen, July, 1939.