INGUINAL HERNIA

A NEW OPERATIVE REINFORCEMENT

BY

A. SIMPSON-SMITH, M.Ch., F.R.C.S.

Surgeon to Out-patients, West London Hospital

Max Page (1934) stated that the recurrence rate in inguinal hernia was far greater than previously admitted, and in healthy metropolitan policemen, whose average age was 32, the rate amounted to 20.2 per cent. in indirect and 25.8 per cent. in direct varieties. Since this paper several carefully examined series of cases supporting Page's view have been published. Grace and Johnson (1937) record 1,032 cases, all over 50, with a recurrence rate of 25.8 per cent., Birkenfeld (1933) gives 28.49 per cent., Lyle (1928) 10.3 per cent., Fallis (1936) 8.25 per cent., and Block (1933), in 642 cases personally followed for two years, 34.8 per cent.

Fascia sutures have been of great service in repairing wide hernias, but even this method is not without defect, for Burdick *et al.* (1937) admit 29.1 per cent. of failures in a series of 975 fascial repairs. In addition, it is tedious and time-consuming to use this material. Ogilvie and Maingot (Maingot, 1937) have employed heavy floss silk loosely woven across the posterior wall of the canal to form a lattice. They claim that the minute fibres constituting the silk cords are readily absorbed, provoke little reaction, and have no tendency to encourage sepsis. Graham (1938), however, states that it is necessary to use fine silk, heavy braided silk being harmful.

Injection treatment alone, while of value in preventing hospitalization, is open to the serious disadvantage that twelve or more appropriate fluid quantities must be placed exactly in the right position. The difficulty of controlling the hernia throughout the treatment and the danger to testis, cord, or bowel by "blind" injections have compelled many surgeons to discard this procedure. What emerges from this method of treatment is the fact that experimentally and clinically selected sclerosing solutions can create a firm area of fibrous tissue.

The general disinclination of the public to accept herniorrhaphy is shown by the number of patients who are prepared to put up with the discomfort of a truss for the rest of their lives: one instrument firm in London manufactures no fewer than 3,000 spring trusses a week.

The present method aims at reinforcing a satisfactory operative repair with the adhesive barrier of the injection treatment.

Technique

The hernial sac is isolated and ligatured as high up as possible, a split salad-spoon facilitating this high excision. A Bassini repair operation is done, approximating the conjoined tendon to Poupart's ligament, either with two strips of external oblique, as in lacing up a boot and originally described by McArthur in 1901, or by the use of non-absorbable sutures. Salmon-gut, because of its lack of capillary attraction and its inability to absorb organisms, is probably safer to use than silk. Orthopaedic surgeons have preferred this material for years. A lozenge consisting of sodium psyllate in gelatin is then placed on the anterior wall superficial to the conjoined tendon but deep to the cord. The external oblique and skin are then closed in the usual manner.

Results

The first case was operated upon in March, 1938. At this time the sac was carefully isolated and, after division between double ligatures, the distal portion was filled with sclerosing fluid. The ball that resulted tended to slip about and was difficult to keep behind the cord. In the next case a slab of gelatin was used, and this proved to be much more convenient.

In all, twenty-three cases have been submitted to this operative reinforcement. Various strengths of sodium psyllate in different conveying materials have been tried. With moderate amounts of sodium psyllate a mild swelling of the wound has occurred, and has subsided in fourteen days. The distinct impression is gained that there is a plate of firm fibrous tissue around the operative site. With too strong solutions intense swelling and bruising of the skin are likely to follow, but even extensive reactions settle down within two or three weeks. Such reactions have not occurred in the later cases.

Conclusions

This is a preliminary report of a method which has not yet stood the test of time. Only the cold candour of a long series of cases will assess its utility. The small number already treated show a firm scar and encourage a further trial. A suitable conveying material that will liberate larger amounts of sclerosing fluid in minute doses over a long period of time is in preparation; this will increase the desirable fibrosis.

I am grateful to Dr. O'Neill and Dr. Shore, medical superintendents of Napsbury and Shenley Hospitals, for allowing me to select cases, and to Mr. Alex Rae, chief pharmacist of the Hospital for Sick Children, for his advice and for the preparation of the psyllate slabs.

REFERENCES

- Birkenfeld, W. (1933). *Arch. klin. Chir.*, **174**, 525.
 Block, W. (1933). *Ibid.*, **175**, 607.
 Burdick *et al.* (1937). *Ann. Surg.*, **106**, 333.
 Fallis, L. S. (1936). *Ibid.*, **104**, 403.
 Grace, R. V., and Johnson, V. S. (1937). *Ibid.*, **106**, 347.
 Graham (1938). *Yearbook of Surgery*, p. 483.
 Lyle, H. H. M. (1928). *Ann. Surg.*, **88**, 870.
 McArthur (1901). *J. Amer. med. Ass.*, **37**, 1162.
 Maingot, R. (1937). *Post-Graduate Surgery*, **3**, 3644.
 Page, C. M. (1934). *British Medical Journal*, **2**, 896.

The Economic Intelligence Service of the League of Nations has just published a study entitled *Urban and Rural Housing*, containing a comprehensive account of the methods employed in a number of countries for improving housing conditions, with special reference to the cost involved and the results obtained. It is of particular interest at the present time, since it deals largely with housing problems raised after the war of 1914-18 and the attempts made to solve those problems in various countries up to the outbreak of hostilities in 1939. It should therefore appeal to all interested in reconstruction schemes at present or in the future. The countries dealt with are Belgium, the United Kingdom, Canada, Denmark, Finland, France, the Netherlands, Norway, Sweden, and the United States of America. The author was able to visit most of these countries and to investigate on the spot the practical difficulties encountered by the housing authorities; in addition he has co-ordinated the material published by or obtained from Governmental and municipal authorities and other sources. A chapter is devoted to each country, and the information arranged in such a way as to permit the reader to compare housing problems and the efforts being made to meet them in different circumstances. A special paragraph is devoted in the case of European countries to the problem of rural housing and the measures taken to improve the housing conditions of agricultural workers or other groups of the rural population. The report is published at 3s. 6d. by George Allen and Unwin, Ltd., 40, Museum Street, W.C.1.