in relieving itching and preventing aggravation of the affection by rubbing and scratching. The local treatment is the same as for ordinary constitutional eczema, simple pastes of the modified Lassar type being most valuable, especially with the addition of prepared coal tar.

Less irritating than these nervous eczematous cases, and generally very much more resistant and widespread, are the seborrhoeic eczemas dependent upon factors of malnutrition. Blood counts, test meals, and vitamin tests will generally indicate fairly marked degrees of malnutrition in these patients, and the case which I illustrate here is a very characteristic example. This patient, I may say, worked eight to nine hours a day for six days a week for a salary of fourteen shillings, and I think such a case will indicate to you the great importance of the general outlook upon this problem rather than the more limited conception of seborrhoeic disorders as of external infective origin. Dietetic treatment, the administration of iron and vitamins by mouth, general ultra-violet-light irradiation of the whole body, and even the use of cod-liver oil as a local application to the skin are measures particularly indicated in this group. (Plate, Figs. 1 and 2.)

From this we pass to the last group of seborrhoeic eczemas, often associated with seborrhoeic sycosis, which are so distressing and intractable. They are characterized by sticky, oozing, pustular, scabbed, and foul eczematous dermatitis in all the sites I have previously mentioned, but particularly affecting the head and about the ears, the eyebrows, eyelids, and beard area. Local treatment, even including x-ray therapy, is often without avail in these cases. They can be cleaned up with starch and boric poultices and fomentations, only to leave a fragile, red, glazed, inflamed skin which within the course of another hour or two becomes covered with small subcuticular pustules which, on the beard particularly, affect the hair follicles. All these cases show some degree of malnutrition, though it may not be of an extreme degree. What is, however, most important is chronic focal sepsis, generally affecting nose, throat, or mouth, and this is hardly ever absent. In the case of a little girl with chronic infected seborrhoeic eczema of the head and face, ears, and flexures, the antra, ethmoids, and sphenoidal sinus contained pus, the gums were infected, and the teeth were degenerate. More often the finding is one of a chronic thickened, hypertrophied, infected, and polypoidal mucous membrane, in which simple drainage of the antra is of little avail. A complete radical operation alone will clear the infected antrum, and unfortunately the majority of cases have infection of this most difficult type. I am convinced, however, that unless the provocative source of the trouble is satisfactorily dealt with these cases remain incurable. It is probable that any operative measures undertaken should be followed by anti-infective measures by mouth and general ultra-violet-light irradiation of the whole body, and any similar therapy likely to increase the patient's resistance to chronic infection.

So far as local application to the skin is concerned in this last group, fomentations, starch, and boric poultices and the use of a brilliant-green or a gentian-violet paint are helpful. Some cases respond to mercury and tar in some such application as the following:

R.	Hydrarg. ammon.	...	...	...	gr. x
	Pic. carb. praep.	...	...	...	gr. xx
	Zinc. oxid.	...	...	...	gr. cxx
	Paraff. moll.	...	...	ad	3 j

Another useful ointment is ung. quinolor co. (Squibb).

Conclusion

These points in treatment of some of the more common of the seborrhoeic disorders illustrate the general principles which I have tried to stress in the earlier part of my lecture. They also remind you, I hope, that the skin is an essential part of the whole organism and reflects most valuably general physiological and pathological changes in that organism. This group of skin disorders illustrates the importance of dermatology in the training of the medical student and the significance of skin disorders to the practising physician.

Summary

It is suggested that the problems of seborrhoeic disorders are related to and dependent upon the seborrhoeic diathesis and not to any great extent upon seborrhoeic or other organisms.

The disturbances of physiological function characteristic of the diathesis are manifest particularly in the membranes of the upper respiratory and gastro-intestinal tracts as well as the skin. As a consequence seborrhoeic patients are subject to chronic infection of the upper respiratory tract and to dyspepsia and malnutrition.

These conditions, together with the underlying nervous and temperamental instability, have an important bearing on aetiology and treatment.

ANTHRA-SILICOSIS AND BRONCHIAL CARCINOMA WITH QUIESCENT TUBERCULOSIS

BY

J. E. LOVELOCK, B.M., B.Ch.

Late House-physician, Brompton Hospital; late House-physician, St. Mary's Hospital

(WITH SPECIAL PLATE)

The increasing incidence of bronchial carcinoma is a widely recognized fact, and statistical surveys in recent years (Duguid, 1927; Simpson, 1928; Kennaway and Kennaway, 1936) suggest that this increase is real, and not merely an apparent one due to improved facilities for observation and more accurate diagnosis. On the other hand, Hadfield and Garrod (1938), who quote Passey and Holmes (1935), against the majority of available evidence, consider that the matter is still *sub judice*. Study and investigation of the aetiological factors of the disease are therefore of particular importance, and with the growing tendency to regard cancer as the product of prolonged mechanical or chemical irritation of an organ with "constitutional and inherited inferiority" in an individual with an intrinsic susceptibility to neoplastic formation (Hurst, 1939), it is inevitable that attention in bronchial carcinoma should be directed towards inhaled irritants. In a condition such as anthra-silicosis, in which chronic irritation of the bronchi and lung by both coal and silica is the outstanding feature, a relatively high incidence of primary bronchial carcinoma might be anticipated. A number of such cases of cancer associated with silicosis have been reported from all parts of the world during recent years, including the studies by Schmorl (1928) on Schneeberg cancer, in which some factor in addition to mechanical irritation would seem to be most active, and by Pancoast and Pendergrass (1933), Dible

(1934), Fine and Jaso (1935), Charr (1937), Anderson and Dible (1938), Turner and Grace (1938), in addition to the Kennaways (1936). At present no definite conclusion has been reached, but that the Silicosis Board in this country is taking serious cognizance of the facts is known.

The following case is recorded because it presented the association of bronchial carcinoma and anthra-silicosis with quiescent tuberculosis, because the difficulty of establishing the diagnosis during life was considerable, and because several features of unusual interest became apparent during the clinical course, at the post-mortem examination, and during subsequent legal proceedings; also it indicates the need for more thorough investigation of such cases, and may "call attention to the need for publishing the post-mortem changes in a large series of cases of silicosis . . . as the recorded instances are as yet too few to be significant" (*British Medical Journal*, 1938).

Case Report

A labourer aged 38 attended a London general hospital in May, 1938, complaining of anorexia, loss of weight, and lassitude of five months' duration, and, more recently, cough and right lumbar pain. Until two years previously he had been a miner of "steam coal" in South Wales for twenty-two years, working mostly at the coal face. The history of this occupation, together with the symptoms and signs which were found, led to the diagnosis of tuberculo-silicosis. Tubercle bacilli were not found in his sputum, and a radiograph of his chest showed "infiltrating inflammatory changes in the left upper, right middle, and lower zones, and possibly left apex—probably phthisis, but not definitely so."

He was referred to the Brompton Hospital towards the end of May. He then complained of lassitude, sweating, loss of weight and strength, cough productive of a small amount of muco-purulent sputum, dyspnoea, and pleuritic pain low down on the right side of the chest and back. He was manifestly ill, wasted, and febrile. An impaired note to percussion was found at both apices, while at the right base there was an area of dullness, poor air entry, and numerous moist rales. His sputum again contained no tubercle bacilli. Radiological examination of the chest showed a mediastinum normal in position, slight restriction of the movements of the right side of the diaphragm, and an area of partial consolidation at the right base, while throughout both lungs changes consisting of diffuse stippled shadowing, compatible with pneumoconiosis, were shown. At the apices the appearances were those of added infiltration due to pulmonary tuberculosis (Plate, Fig. 1).

The patient was transferred to St. Mary's Hospital, where the above history and chest signs were confirmed; but in addition routine examination showed generalized abdominal tenderness, with resistance most marked in the right upper rectus. This was thought to be due to diaphragmatic irritation. There was no clubbing of the fingers, and no enlarged glands were palpable. Other systems were clinically normal. Radiological examination of the chest showed changes consistent with the early nodular phase of silicosis, with superimposed tuberculosis at both apices and in the right lower lobe, where early cavitation seemed to have occurred. Relatively slight basal pleural involvement was shown. The blood sedimentation rate was 63 mm. in one hour (Westergren). Repeated sputum examinations again failed to show tubercle bacilli, and no doubly refractile bodies could be seen under polarized light. His condition deteriorated in spite of absolute rest and symptomatic treatment for cough and pain. A month after admission he developed transient glycosuria for two days, but a blood sugar curve two days later was normal. He next developed symptoms and signs suggestive of involvement of the central nervous system. These were mainly on the left side of the body, and included temporary and varying loss of power, pains and cramps in the left arm and leg, with sensory loss which varied from day to day.

Late in June, in the course of routine re-examination, the abdominal resistance was found to have decreased, permitting the recognition of an enlarged tender nodular liver, which extended nearly to the umbilicus. This finding cast doubt on the original diagnosis, and suggested either that the case was one of carcinomatosis or that a carcinoma with hepatic secondaries had occurred in a case of tuberculo-silicosis. Immediate search was made for other secondaries. Radiological examination of the skeleton showed three doubtful metastases in the right femur and in the third and fourth lumbar vertebrae. A blood count showed 4,500,000 red cells per c.mm., with 98 per cent. haemoglobin and a colour index of 1.1, and 27,000 white cells, of which 86 per cent. were polymorphs and 14 per cent. lymphocytes. Two myelocytes, six nucleated reds, and numerous Howell-Jolly bodies per 100 white cells were seen. An early leuco-erythroblastic anaemia was therefore present, which strongly suggested carcinomatosis of bone marrow. Multiple metastases in the brain were suspected as the cause of the somewhat vague symptoms and signs in the nervous system, while the hepatic condition, with probable pancreatic metastases, might have been responsible for the transient glycosuria.

Though pain and cough did not increase, the patient's general condition continued to deteriorate; he wasted rapidly, became mentally confused and incontinent, and early in July died of a terminal bronchopneumonia.

Post-Mortem Examination

Necropsy showed the double condition of anthra-silicosis of the lungs, most marked in the upper lobes and apices, and carcinoma of the right lower lobe bronchus; there was slight terminal bronchopneumonia (Fig. 2). The right pleura was thickened, and through it at the base grew the branches of an aspergillus. There were extensive anthracotic deposits throughout both lungs, but no microscopical evidence of any tuberculous involvement or caseation of the whorled, nodular, fibrous, silicotic areas, where the compact centre was surrounded first by a zone of collagenous fibrous tissue, then by a zone of compressed lung tissue heavily laden with black anthracotic pigment, which faded out into comparatively normal lung (Fig. 3). However, a guinea-pig injected with a saline suspension of the lung died of generalized tuberculosis in early April, 1939, nine months later. The carcinoma was a typical small oat-cell growth, surrounding and compressing the bronchi, with their cartilage and mucous glands (Fig. 4). Metastases were found in the regional glands, in the liver—which weighed approximately 6 kg. and was studded with growth—and in the skeleton (long bones and skull); and a few were seen in the pancreas.

Chemical analysis of the lungs as compared with a normal control showed a greatly increased silica content—between eight and nine times normal:

	Present Case	Control
Silica in fresh lung ..	2.83%	0.31%
Silica in dry lung..	12.43%	1.56%

Causes of Death

These were certified as bronchopneumonia, carcinomatosis from bronchial carcinoma, with silicosis as an associated condition. The mention of silicosis—an industrial disease—necessitated a coroner's inquest. Here the jury, on the direction of the coroner, after hearing the medical and pathological evidence, returned a verdict of death from misadventure—death in their opinion resulting from "bronchopneumonia and carcinomatosis, following carcinoma of the bronchus caused by silicosis due to his occupation as a coal-miner." As a consequence of this verdict the dependants claimed compensation, and in due course the case came before the Silicosis Board. During its consideration a representative of the board pointed out that in law silicosis and carcinoma of the lung had as yet no proved aetiological relationship,

whatever medical evidence might be adduced to the contrary. At the moment of writing, the dependants inform us that the question of compensation has not yet been settled, but indicate that if their legal advisers consider the verdict of the jury sufficient to justify it, the Miners' Federation, who have taken over the case, will pursue the claim further.

This case may most profitably be discussed under the heading of clinical, radiological, pathological, and medico-legal, of which the first and last are of most concern to the general practitioner, while the second and third interest rather the hospital consultant, radiologist, and pathologist.

Clinical Points

Clinically three points arise. First is the importance of the treble diagnosis, which was for so long overlooked by three large hospitals with all the facilities for full investigation. Diagnostic difficulties during the patient's life were, however, considerable. The occupational history, together with the radiological appearances, suggested that silicosis was present, while the advent of toxæmia, anorexia, sweating, loss of weight, and a productive cough made it seem probable that pulmonary tuberculosis had supervened. Alternatively the possibility of chronic pulmonary tuberculosis complicated by a sub-acute miliary spread was considered, but was regarded on radiological grounds as less likely than the above.

The second point is the fact that this delay in recognition was due to the difficulty of differentiating the clinical symptoms and signs of phthisis from those of carcinoma—cough, pain, dyspnoea, wasting, and sputum. The differential diagnosis in the presence of silicosis is rendered doubly difficult because the association of tuberculosis with silicosis is an accepted medical and legal one, whereas that of a carcinoma with silicosis is a less frequently occurring and less generally recognized condition. In this case, as shown by the long-delayed death of the injected guinea-pig, tuberculosis was present, but it should be classified as quiescent rather than as active. The toxæmic symptoms are much more likely to have been due to the carcinoma than to the tuberculosis, but the emphasis on the latter led to the overlooking of the former. As Charr (1937) points out, the persistently negative tests to tubercle bacilli in sputum from a known or suspected silicosis with such toxic symptoms should excite a suspicion of malignant lung growth.

The third point is the importance of considering the possibility of carcinoma in a middle-aged man in the presence of any chest symptoms. This matter was recently fully discussed at the Royal Society of Medicine by Roberts, Davidson, Mason, and Lakin (*British Medical Journal*, 1939), who emphasized the urgency for full investigation at the very first onset of chest symptoms, so that the condition might be recognized at a sufficiently early stage for something more than palliative treatment to be given. The significance of persistent pain over the base of the right lung was not in this case sufficiently emphasized.

Radiological Appearances

Radiologically the case illustrates the difficulty in diagnosis, even in the presence of full clinical findings, between an inflammatory and a neoplastic pulmonary lesion. Pendergrass (1938) discusses this fully in relation to the pneumoconioses, and he also considers the associated presence of tuberculosis and carcinoma; while in occupational diseases in general the British Institute of Radiology

(1938) concluded that the diagnosis of such cases must be largely based on radiological appearances, for the proper interpretation of which the occupational history, with a knowledge of the minutiae of each occupation, is essential. In the present case we were perhaps a little blinded by the emphasis on occupational history.

Pathological Discussion

The association between chronic irritation due to inhaled dust particles containing silica and the development of carcinoma is one which has recently attracted attention, and to which reference has already been made. Though Valade (1939) questions the validity of the application of animal experiments to man, pathological studies such as those of Willis (1928), Willis and Brutsaert (1928), and Campbell (1937) are significant and have shown that the epithelium of the bronchial mucosa undergoes a hyperplasia as a result of chronic irritation by inhaled dust of this type, and that tar-free dust containing 44 per cent. silicon dioxide increases the incidence of primary lung carcinoma in mice by 45 per cent., but not to so great an extent (74 per cent.) as when tar also is present. Analogous combined chemical and physical irritation arising in industrial life may therefore well be responsible for cancer in man, and in the particular case cited above, the hydrocarbons of coal, which some observers have suggested may to a certain extent protect against tuberculosis, may be the relevant chemical factor. Dible (1934) points out that the rarity of tuberculosis in his cases of silicosis with malignant disease tends to suggest that a degree of resistance to this disease, perhaps unusually well developed, may have permitted an exceptionally long exposure of the tissues to the irritant agent. Similarly Turner and Grace (1938), commenting on the figures of Rennie (1933) on the mortality of grinders from tuberculosis, rightly point out that most of this respiratory mortality operates before the cancer age, therefore depleting the ranks of the potential carcinoma victims. In spite of this depletion they find a great increase of both silicosis and cancer in the Sheffield grinders. In this connexion one is impressed with the fact that throughout an extensive study of the literature carcinoma seems, with the exception of one case of Dible's, to be associated, not with *active* tuberculo-silicosis, but almost entirely with anthra-silicosis. Though this may be a fortuitous association, yet in view of Campbell's work one cannot help wondering how large a part in the carcinoma causation is played by the chemical action of the carbons of the anthracosis superadded to the physical and chemical action of the silica in silicosis production.

In spite of the accumulating mass of evidence, and the conclusion of Anderson and Dible that "as a group of cases of pulmonary carcinoma exists in which the organs contain an excess of silica and show histological evidence of silicotic fibrosis, the role of silicosis is definitely aetiological," I cannot but agree with Gloyne (1938) that the data we possess at the present time are not sufficient to justify any dogmatic statement, and that we cannot do more than regard the association of silicosis and carcinoma (and of asbestosis and carcinoma) with suspicion. There is, however, room for more work and for reports of more cases, together with full post-mortem findings.

The Medico-Legal Aspect

This aspect is one that may be brought to the notice of any practitioner who signs a death certificate in which carcinoma and silicosis are mentioned together. That both existed in this case was proved pathologically and

chemically, for the normal silica content of dried lung does not exceed 0.2 per cent., and Sladden (1933) has reported that a silica content of more than 1 per cent. is usually accompanied by an important and radiologically recognizable degree of fibrosis, while a content of over 1.6 per cent. occurs only in lungs with severe fibrosis sufficient in itself to cause death. Belt, Irwin, and King (1936) consider that reliance must not be placed on chemical analysis alone in establishing the existence of silicosis; in this they are supported by Middleton (1936), who emphasizes that since silicosis is caused by particles less than 10μ in diameter, and since much larger particles may be included in the chemical analysis, much more silica will be demonstrated by this means than is necessarily exerting a pathological fibrotic effect. Therefore, in order to establish a diagnosis beyond doubt the combination of clinical, radiological, microscopical, and chemical findings is desirable. In this particular case the presence of both silicosis and carcinoma was accepted by the representative of the Silicosis Board, but the aetiological connexion between the irritation and the fatal carcinoma was not, in the present legal position, admitted. At the time of writing the dependants await the ultimate legal decision with interest. Whatever that may be, as the evidence and literature accumulate, the compensation laws of silicosis, summarized recently by Davie (1937), may shortly have to be altered to fit in with accepted medical facts. That the law lags behind medicine in many respects is agreed, as, for instance, in the association between tuberculosis and silicosis, which was not recognized till the 1925 Scheme, following the House of Commons' acceptance in 1924 that the presence of the former made the latter a more dangerous disease. In a recent review of the medico-legal aspects of coal-mining, Fisher (1938), in discussing silicosis, makes no mention of carcinoma and very little of tuberculosis, leaving the impression that from the medical standpoint, even in respect of pure uncomplicated silicosis, the present law is not entirely satisfactory. The whole question of chronic pulmonary disease of a disabling nature which does not come within the accepted definition of silicosis is being studied by the Medical Research Council (1937-8) at the request of the Home Office Mines Department, and some change in the present legal position seems likely. In this case an acceptance in law of the importance of silicosis in predisposing to carcinoma would greatly benefit the dependants.

Summary

A case of combined anthra-silicosis and bronchial carcinoma with quiescent tuberculosis in a Welsh coal-miner resident in London is recorded, with full post-mortem findings. The difficulties of diagnosis during life are discussed, and the implications of the association between chronic physical and chemical irritation and the development of neoplasm are considered from the clinical, radiological, pathological, chemical, and medico-legal viewpoints, with reference to some of the modern literature.

It is a pleasure to record my thanks for much advice and assistance on the various aspects of this case; especially to Dr. W. D. W. Brooks for permission to report it, to Professor W. D. Newcomb for the pathological work and the photomicrography, to Dr. Roche Lynch and Dr. R. H. Slater for the chemical analyses, to Dr. E. Rohan Williams for assistance with the radiology, and to Drs. M. Davidson and G. T. Hebert for their kindly criticisms.

REFERENCES

- Anderson, C. S., and Dible, J. H. (1938). *J. Hyg., Camb.*, **38**, 185.
 Belt, T. H., Irwin, D., and King, E. J. (1936). *Canad. med. Ass. J.*, **34**, 125.
 Brit. J. Radiol., 1938, **11**, 345.
 British Medical Journal, 1938, **2**, 411.
 — 1939, **1**, 348.
 Campbell, J. A. (1937). *Brit. J. exp. Path.*, **18**, 215.
 Charr, R. (1937). *Amer. J. med. Sci.*, **194**, 535.
 Davie, P. C. (1937). *Med.-leg. and Criminol. Rev.*, **5**, 154.
 Dible, J. H. (1934). *Lancet*, **2**, 982.
 Duguid, J. B. (1927). *Ibid.*, **2**, 111.
 Fine, M. J., and Jaso, J. V. (1935). *J. Amer. med. Ass.*, **104**, 40.
 Fisher, S. W. (1938). *Med.-leg. and criminol. Rev.*, **6**, 245.
 Gloyne, S. R. (1938). *Silicosis and Asbestosis*, edited by Lanza, A. J., **4**, 198, Oxford University Press, London.
 Hadfield, G., and Garrod, L. P. (1938). *Recent Advances in Pathology*, London.
 Hurst, A. (1939). *Lancet*, **1**, 553.
 Kennaway, N. M., and Kennaway, E. L. (1936). *J. Hyg., Camb.*, **36**, 236.
 Medical Research Council Ann. Report, 1937-8, p. 18, London.
 Middleton, E. L. (1936). *Lancet*, **2**, 1.
 Pancoast, H. K., and Pendergrass, E. P. (1933). *J. Amer. med. Ass.*, **101**, 587.
 Passey, R. D., and Holmes, J. M. (1935). *Quart. J. Med.*, n.s. **4**, 321.
 Pendergrass, E. P. (1938). *Silicosis and Asbestosis*, edited by Lanza, A. J., **3**, 66, Oxford University Press, London.
 Rennie, J. (1933). *Ann. Rep. M.O.H. Sheffield*.
 Schmorl, G. (1928). *Rep. Internat. Conf. on Cancer*, London, p. 272, John Wright and Sons, Bristol.
 Simpson, S. L. (1928). *Quart. J. Med.*, **22**, 413.
 Sladden, A. F. (1933). *Lancet*, **2**, 123.
 Turner, H. M., and Grace, H. G. (1938). *J. Hyg., Camb.*, **38**, 90.
 Valade, P. (1939). *Bull. Acad. Méd., Paris*, **121**, 115.
 Willis, H. S. (1928). *Amer. Rev. Tuberc.*, **17**, 253.
 — and Brutsaert, P. (1928). *Ibid.*, **17**, 268.

THE ORAL APPLICATION OF METHYL TESTOSTERONE AND ITS SIMPLIFICATION OF ANDROGEN THERAPY*

BY

GEORGE L. FOSS, M.A., M.B., B.Ch.

Colston Research Fellow, 1935-7; Clinical Assistant in the Gynaecological Out-patient Department, Bristol Royal Infirmary

(WITH SPECIAL PLATE)

Recently I described the substitution effect and dosage of testosterone (tablets of 2.5 mg.) given to a post-puberal eunuch (Foss, 1938b, 1939). It was found that 100 mg. daily produced very mild potency, and a ratio of 20:1 was suggested as the comparison of this oral effect with that by subcutaneous injection. Although the oral route is to be preferred to the percutaneous inunction (Foss, 1938a; Loeser, 1938a, 1938b) or injection, especially where long-continued substitution therapy is necessary, it is wasteful, expensive, and rather inefficient.

Miescher and Tschopp (1938) have demonstrated that methyl testosterone is considerably more active by mouth than testosterone, androstenediol, androstenedione, testosterone propionate, and androsterone in castrated male rats as shown by the increase in weight of the seminal vesicles and prostate. They suggested that possibly greater protection of the active hydroxyl group from fermentative influences in the alimentary canal was acquired from the adjacent methyl group.

Case Reports

The following report describes, from both the subjective and the objective points of view, the clinical trials of methyl testosterone which I have conducted since February 1 of this year.

Case 1.—Methyl testosterone was given to the post-puberal eunuch, already described (Foss, 1937, 1938b, 1939), after fourteen days' rest from previous treatment, when complete impotence had returned. A dose of 100 mg. was given daily, divided into ten tablets of 10 mg. This dose was taken for

* This paper was received from the author on May 2, 1939.

J. E. LOVELOCK: ANTHRA-SILICOSIS AND BRONCHIAL CARCINOMA, WITH QUIESCENT TUBERCULOSIS

FIG. 1.—Radiograph of chest showing infiltration at both apices, consolidation in right lower lobe, and mottled shadows in both lung fields.

FIG. 2.—Right lung in sagittal section, showing dark anthra-silicosis, most marked in the upper lobe, and white neoplastic extension from the lower lobe bronchus with glandular involvement and small lung metastases.

FIG. 3.—Edge of anthra-silicotic nodule, $\times 100$: A, relatively normal lung; B, compressed lung tissue laden with pigment; C, collagenous fibrous tissue; D, compact centre.

FIG. 4.—Edge of bronchial carcinoma, small oat-cell type, $\times 100$: D, lung tissue; C, carcinoma; B, bronchial mucous glands; A, bronchial cartilage.