onlooker—for in this great field I have been no more—I have reached the conviction that recent years have seen cancer research enter on yet another new phase, one in which experimental studies are going deeper than before into the nature of the phenomena involved and yielding results of unforeseen promise.

I will now before concluding turn once more to the activities of the Imperial Fund, wishing to pay tribute to the wisdom of the policy of its executive and another tribute to the high qualities of the work of its staff to-day. These qualities recall those of its early pioneer days to which I have referred, though displayed on a field so much wider in scope.

Freedom, Leisure, and the Side-paths

I would like to make a few remarks here which I hope will not be thought intrusive. It has often been said, not without justification, that workers in a centre of experimental activity devoted wholly to one limited end are in danger, owing to their special preoccupations, of ignoring too much the advances won in other branches of scientific endeavour. These may often contain suggestions and even inspiration which can contribute greatly to the progress of their special task. Frontal attacks alone are proverbially not always the most successful method of dealing with an entrenched enemy. The case of cancer research is one, I think, in which this is especially true. Many of its problems are to be found in aspects of living nature of so fundamental a kind that inspiration may come unexpectedly from discoveries made in many branches of pure science.

The ideal for a centre of cancer research would seem to be that its workers, though with their ultimate goal always in mind, should have liberty and leisure to pursue paths which, though apparently remote from what seems to be the direct path, may, in the final issue, prove to be short cuts towards that goal. Such freedom is of the highest value in research. If I mistake not, all desirable forms of scientific freedom are to be found and will continue to be found in this laboratory. The Imperial Fund, I think, remains unique among the many endowed centres of cancer research now in this country. The distinguished personnel concerned with the general direction of its policy, its independent finance, its scientific tradition based on nearly forty years of work, the composition of its laboratory staff so widely organized as a working unit—all these distinguish it and guarantee a future of continued and eminent success. Those who have the privilege of knowing its present director and the senior members of the staff have a further reason for being well assured of this.

Grounds for a Hopeful View

Allow me to conclude with an attempt to appraise briefly the present position of the cancer problem, even if I speak with perhaps too little authority. If the position be reviewed calmly and without prejudice it surely shows reasons for much satisfaction. Progress towards an understanding of the actual nature of the disorder is increasing remarkably fast, and when a full understanding of this comes a happy solution of the whole problem will not be too far distant. New constellations of illuminating facts continue to appear above the horizon of knowledge, and it is not too much to hope that at any moment some pregnant item of new knowledge may appear among them which, like a pole star, will point directly to the right path for effective action. Such a consummation may come soon or in a more distant

future. Meantime the patient efforts of investigators must continue.

With the assurance that fundamental knowledge concerning the actual nature of cancer is continuously increasing, together with the realization that success in treatment grows with the clinician's increasing skill and equipment, there is surely justification to-day for an optimistic outlook. It is most desirable that the lay public should be persuaded to share in such optimism.

THE SEBORRHOEIC DIATHESIS*

BY

JOHN T. INGRAM, M.D., F.R.C.P.

Physician in Charge of the Skin Department, the General Infirmary at Leeds; Lecturer on Diseases of the Skin, University of Leeds

(WITH SPECIAL PLATE)

I understand from my medical dictionary that diathesis "is a term at present but little used." I know, too, that it is a little bold and bad of me to journey to Scotland and talk of seborrhoea. At the same time I believe we cannot too often stress the importance of the constitutional side of medical disorders in this age of scientific progress, and the evidences of those constitutional disturbances are constantly written upon and manifest in the skin for those who will regard them. The modern conception of seborrhoeic dermatoses is that they are infective, arising from the activity of organisms which normally inhabit the skin but which abound on a greasy one. To that view I cannot subscribe, and I shall endeavour to express a conception of these many disorders which in my view links them together and justifies the retention of the term "seborrhoeic," which many would abandon.

Literally the seborrhoeic is one whose skin is running with sebum, but this feature may not be very pronounced. It is a fact that such skins also tend to run with sweat, so that seborrhoea and hyperidrosis commonly go hand in hand. In talking, therefore, of seborrhoeic affections one must include hyperidrotic affections such as pompholyx of hands and feet. A second point to be borne in mind is the profound change which takes place in the skin and is essentially centred round glandular secretion at the time of puberty. Some of these changes are reflected in the increased growth of organisms which normally inhabit the skin, and are reflected still more strangely in the disappearance of the ordinary small spore ringworm of scalp (*Microsporon audouini*) and by the appearance of other infections such as the tinea versicolor infection (*Microsporon furfur*). One last point to which I would draw your attention before proceeding to my subject is that these seborrhoeic changes, though normally occurring in the skin at the time of puberty, may upon rare occasions arise as the result of lesions in the central nervous system, as in encephalitis lethargica, and may arise from disease or disorder of endocrine glands at times other than puberty.

Seborrhoeic Diathesis

In seborrhoea there is an increase of sebaceous secretion, which is altered and is more acid; and with this increase there is an overgrowth of the sebaceous glands and an enlargement of the pores. This phase tends to

* British Medical Association Lecture to the Fife Branch, read at Kirkcaldy on March 2, 1939.

produce some thickening of the surface layers of the skin, particularly about the enlarged pores, a condition described as *kérose*. In addition there is increased activity of the sweat glands, which I have previously mentioned. There is a tendency to vasomotor instability, often *perniosis* and chilblain development, and that non-pitting chronic oedema which gives the characteristic podgy clammy hands and stocky ankles and legs of puberty. There is an increased sensitiveness of the skin to ordinary external irritants, as a result of which seborrheic are regarded as bad industrial risks in trades that carry with them dermatological hazards. There is similarly an increased susceptibility to infections, particularly ordinary pyogenic infections such as boils, pustules, and carbuncles, *perionychia*, fissures, and similar conditions. It is difficult to imagine that such considerable physiological changes in the skin can emanate from any but a central source, and certainly the channel along which these influences are exerted, if not the original source, is a nervous one.

The next important point to consider, in my opinion, is the inevitability of similar changes in the mucous membranes; this is borne out abundantly by clinical evidence. In the upper respiratory tract the seborrheic shows a catarrhal mucous membrane with increased mucous secretion and boggy membranes. The membranes, like the skin, are sensitive to external irritants and to infection, with the result that these patients tend to suffer from chronic catarrh and hypertrophy of the mucous membranes, from spasmodic rhinorrhoea and hay fever, from thickened turbinates, infected antra, ethmoids, and other sinuses. Often there is a chronic polypoidal thickening of the mucosa of these sinuses. They have bad teeth and gums, pyorrhoea and gingivitis, caries and sepsis—a condition which the dentists describe as “soft teeth.” In hospital practice it is almost the hall-mark of a seborrheic that he has a complete set of false dentures or that he should have such a set in his early twenties. The importance of this aspect of the seborrheic is that he tends to harbour foci of chronic infection about the upper respiratory tract which are difficult to clear and which have a chronic debilitating effect upon his health.

Similar catarrhal changes, often with a tendency to bronchitis and sometimes to asthmatic disorders, characterize the respiratory tract, but more important from our point of view is that the “seborrheic influences” extend to the gastro-intestinal tract, where they are frequently associated with an early dyspepsia of the functional flatulent type. Here there may be an excess of mucus with masking of hydrochloric acid and, what is probably more important, some degree of hypotonia. Apart from this there is commonly chronic constipation, and these two influences are often responsible for malnutrition of greater or less degree. The dyspeptic tends to adapt his diet to his dyspepsia, which is not helpful. This and impaired assimilation lead to an iron lack with secondary anaemia and to lack of vitamins, both of which changes may be demonstrable in the laboratory, though they are of significance, it may be assumed, before they reach these proportions. The diet best fitted to the seborrheic, and probably to those suffering from any catarrhal state, is the anti-retentional type of dietary, rich in proteins, restricted in carbohydrates, fluids, and salt, and rich in fresh foodstuffs. It is, of course, essential to avoid irritants and highly seasoned foods, fried foods, pork, and probably cheese. It is often necessary to give iron by mouth and vitamin preparations.

There are various minor variations from the normal, such as the secretion of a hyperacid urine, the exact

significance and importance of which is not clear, but attention must be given to temperamental instability. This may vary from the sullen, sulky indolence of some of the heavier forms of acne to the garrulous, palpitating, highly strung overactivity of the rosacea patient. These nervous influences are probably of much greater significance, especially in many of the puberty affections, than we have previously imagined.

It will be seen that the greasy skin of the seborrheic is but one small indication or reflection of a constitutional state or diathesis which is of considerable importance, and the eruptions which we dub seborrheic are in fact reflections of disturbances of the constitutional state rather than infective conditions peculiar to the skin. While circumstances of health, employment, and environment generally may do much to promote the seborrheic state, this diathesis is probably an inherited one in the vast majority of cases. States of poverty and malnutrition; states of anxiety, stress, and strain; employment in hot, dusty atmospheres, as mining; and absence of light and fresh air are conditions which the seborrheic will not readily tolerate.

Treatment

The importance of these considerations lies in the guidance they must give one in the treatment of seborrheic affections. Before proceeding to any remarks upon particular seborrheic affections I would like to indicate those general points in the approach to and treatment of the seborrheic which in my opinion are essential to a proper conception of any and every one of the seborrheic ills. You will understand that it is not necessary for a seborrheic to suffer from any of these disturbances which I have mentioned so long as he is healthy in body and mind. It is also true that in one or two trivial seborrheic affections reasonably satisfactory results can be obtained by attention to local toilet measures alone. What I am particularly anxious to stress is that we should not regard scurf of the scalp as the beginning and end of all seborrheic affections, or the treatment of such scurf, if present, as the essential part in the cure.

Whether you are treating a simple affection like scurf of the scalp or a serious chronic seborrheic eczema and sycosis, it is important, first, that your patient should have any dyspepsia or constipation corrected and that he should be put on to a diet similar to that indicated above. It is important, secondly, that any infection in the nose, throat, and mouth should be eradicated, and in this connexion a careful history, particularly as regards influenza, a direct examination, and a simple transillumination of the antra will generally indicate the presence or absence of infection. Thirdly, the nervous and psychological make-up of the patient must be given some consideration, particularly in relation to his environment and to any changes occurring at or before the onset of the skin affection. Finally, it is desirable for the majority of seborrheics to adopt a strict toilet routine, including a weekly shampoo, a daily bath, the use of a dusting powder, and the wearing of cotton, preferably cellular cotton, undergarments. This may sound a somewhat elaborate approach to what is no more than a rash, but a little thought will convince you that it comprises no more than the simple clinical overhaul which is called for in any constitutional disturbance.

The disorders we have under consideration may be grouped under three headings: first, the simple, infective disorders; secondly, physiological disorders; and, thirdly, the major constitutional seborrheic disturbances.

Infective Seborrhoeic Disorders

The first group comprises merely scurf of the scalp and scurf of the body—that is, pityriasis capitis and pityriasis corporis which arises *in situ* and not as the result of scurf falling on to the body from the scalp. To effect control of these disorders it is sufficient to give attention to local treatment. As you are well aware, the scalp condition can generally be controlled by the regular use once a week of a soap-spirit shampoo, and the use of a weak mercurial or a sulphur and salicylic acid ointment for the severer cases. The milder cases are more comfortably treated with a lotion containing perchloride of mercury and resorcin, which is applied daily:

R̄	Hydrarg. perchlor. ...	...	...	gr. ij
	Resorcin or euresol ...	...	...	3 j
	Industrial spirit ...	...	...	3 ij
	Aq. lavand. ...	...	...	q.s.
	Aq. ...	...	...	ad 3 vj

If you meet with a very obstinate case of scurf of the scalp which completely resists your usual attack, consider whether it may not be a psoriasis. Even though you cannot satisfy yourself that it is psoriasis of the scalp, the use of treatment suited to psoriasis may prove effective. In young babies seborrhoeic infections of the scalp may be ringed like seborrhoeic infections of the body, and may suggest a ringworm infection.

Pityriasis corporis generally clears with baths and an ointment such as the following:

R̄	Ung. sulphur. ...	...	...	3 j
	Ung. zinci (1914) ...	...	...	ad 3 j

These patients should be warned of the necessity for regular toilet baths and the wearing of cotton garments next the skin.

Physiological Seborrhoeic Disorders

The second group consists of acne vulgaris and rosacea. The use of the old term "acne rosacea" should be abandoned for fear of confusing one condition with the other. In acne excessive secretion, in rosacea excessive flushing, are responsible for the reactions. In acne, which affects patients at puberty, there is generally a pallid, coarse, greasy skin and always blackheads, which, blocking the sebaceous glands, are responsible for retention of sebum. Secondary infection may now give rise to the characteristic papular and pustular inflammatory lesions related to the pilo-sebaceous follicles. These skins need to be washed vigorously and thoroughly to counteract the greasiness. In rosacea the skin is flushed and extremely sensitive, so that soap and water is best avoided, and whatever treatment is employed it should be of a gentle type and not of the vigorous peeling character used for acne vulgaris. Rosacea most commonly occurs in females at the menopause, though it may arise at other times, and may, of course, affect males. The papules in rosacea are not related to the glands.

Local treatment is of great importance in acne, and two measures may be stressed. In the first place x-ray therapy in small dosage, unfiltered, at proper intervals, is extremely valuable, since it diminishes the activity of the glands, which consequently get smaller. The "soil" is then no longer suitable for the development of acne lesions. The second measure is the use of sulphur in a gradually increasing strength in a heavy calamine lotion, which should be "whitewashed" on the skin with a brush last thing at night before the patient retires to bed and

immediately after washing thoroughly with soap and hot water:

R̄	Sulphur. praecip. ...	...	...	gr. xx
	Calamin. ...	...	...	gr. xxx
	Zinc. oxid. ...	...	...	gr. xx
	Glycerin. ...	...	...	℥ xxx
	Aq. calcis ...	...	...	ad 3 j

While these local treatments are of great importance, in a very large number of cases other considerations have to be taken into account. It is always wise to put these patients on an anti-retentional type of diet, and it is important that the nose, throat, and mouth should be in good order. In addition to this, however, quite small doses of thyroid are sometimes indicated, especially in the sluggish type of patient with a slow pulse, a chilblain type of circulation, and possibly chilblains themselves. One of the most difficult influences in acne is the nervous and psychological instability associated with puberty and the following years. Quite often it is helpful to give these patients a quarter of a grain of luminal at night over long periods; it makes them less anxious and disturbed over their mental stresses and strains and conflicts, and does not impair their vitality.

Luminal 1/4 grain at night is almost an essential in the treatment of rosacea, which is always an evidence of hypersensitiveness. The most effective local treatment, however, is the use of quite small doses of x rays at infrequent intervals combined with a 2 per cent. sulphur and salicylic acid ointment at night and the use of a simple soothing calamine cream by day. This treatment, which was suggested by MacCormac, is remarkably effective, and though rosacea patients must generally expect relapses from time to time, at long intervals, especially over the phase of the menopause, the rapid relief given is a considerable help psychologically. I will not dwell here upon attention to diet and pelvic troubles.

Constitutional Seborrhoeic Disturbance

The third group of affections consists of those which are essentially constitutional seborrhoeic manifestations. They include, first, eczematous symptoms. These may be localized or widespread and of the acute, subacute, or chronic type. Common examples are the eczemas about and behind the ears; eczemas in flexures, especially the axillae, groins, and perineum; eczemas of the pompholyx type affecting hands and feet; and eczema of the nape of the neck. Any of these affections may become more extensive and widespread, and they may be dependent upon any of those important constitutional considerations which I stressed earlier in this lecture; but by far the most common underlying factors in such cases are psychological and nervous considerations. These may be no more than general nervous debility from wear and tear or the reaction from anxiety or worry, from shock or prolonged nervous strain. It is important to discuss the aetiology with the patient and to get his co-operation, both in elucidating and in treating this aspect of the problem. It will, however, help considerably in the relief of symptoms and in re-establishing the patient's confidence if he is put on sedative- tonic measures internally, such as luminal 1/4 grain or the following mixture:

R̄	Pot. brom. ...	...	...	gr. x
	Sp. ammon. arom. ...	...	...	℥ xv
	Tinct. nucis vom. ...	...	...	℥ iv
	Infus. gent. ...	...	...	ad 3 ss

Here again, if the affection is not too widespread, the use of x-ray therapy may be most valuable, particularly

in relieving itching and preventing aggravation of the affection by rubbing and scratching. The local treatment is the same as for ordinary constitutional eczema, simple pastes of the modified Lassar type being most valuable, especially with the addition of prepared coal tar.

Less irritating than these nervous eczematous cases, and generally very much more resistant and widespread, are the seborrhoeic eczemas dependent upon factors of malnutrition. Blood counts, test meals, and vitamin tests will generally indicate fairly marked degrees of malnutrition in these patients, and the case which I illustrate here is a very characteristic example. This patient, I may say, worked eight to nine hours a day for six days a week for a salary of fourteen shillings, and I think such a case will indicate to you the great importance of the general outlook upon this problem rather than the more limited conception of seborrhoeic disorders as of external infective origin. Dietetic treatment, the administration of iron and vitamins by mouth, general ultra-violet-light irradiation of the whole body, and even the use of cod-liver oil as a local application to the skin are measures particularly indicated in this group. (Plate, Figs. 1 and 2.)

From this we pass to the last group of seborrhoeic eczemas, often associated with seborrhoeic sycosis, which are so distressing and intractable. They are characterized by sticky, oozing, pustular, scabbed, and foul eczematous dermatitis in all the sites I have previously mentioned, but particularly affecting the head and about the ears, the eyebrows, eyelids, and beard area. Local treatment, even including x-ray therapy, is often without avail in these cases. They can be cleaned up with starch and boric poultices and fomentations, only to leave a fragile, red, glazed, inflamed skin which within the course of another hour or two becomes covered with small subcuticular pustules which, on the beard particularly, affect the hair follicles. All these cases show some degree of malnutrition, though it may not be of an extreme degree. What is, however, most important is chronic focal sepsis, generally affecting nose, throat, or mouth, and this is hardly ever absent. In the case of a little girl with chronic infected seborrhoeic eczema of the head and face, ears, and flexures, the antra, ethmoids, and sphenoidal sinus contained pus, the gums were infected, and the teeth were degenerate. More often the finding is one of a chronic thickened, hypertrophied, infected, and polypoidal mucous membrane, in which simple drainage of the antra is of little avail. A complete radical operation alone will clear the infected antrum, and unfortunately the majority of cases have infection of this most difficult type. I am convinced, however, that unless the provocative source of the trouble is satisfactorily dealt with these cases remain incurable. It is probable that any operative measures undertaken should be followed by anti-infective measures by mouth and general ultra-violet-light irradiation of the whole body, and any similar therapy likely to increase the patient's resistance to chronic infection.

So far as local application to the skin is concerned in this last group, fomentations, starch, and boric poultices and the use of a brilliant-green or a gentian-violet paint are helpful. Some cases respond to mercury and tar in some such application as the following:

R.	Hydrarg. ammon.	...	...	...	gr. x
	Pic. carb. praep.	...	...	...	gr. xx
	Zinc. oxid.	...	...	...	gr. cxx
	Paraff. moll.	...	...	ad	3j

Another useful ointment is ung. quinolor co. (Squibb).

Conclusion

These points in treatment of some of the more common of the seborrhoeic disorders illustrate the general principles which I have tried to stress in the earlier part of my lecture. They also remind you, I hope, that the skin is an essential part of the whole organism and reflects most valuably general physiological and pathological changes in that organism. This group of skin disorders illustrates the importance of dermatology in the training of the medical student and the significance of skin disorders to the practising physician.

Summary

It is suggested that the problems of seborrhoeic disorders are related to and dependent upon the seborrhoeic diathesis and not to any great extent upon seborrhoeic or other organisms.

The disturbances of physiological function characteristic of the diathesis are manifest particularly in the membranes of the upper respiratory and gastro-intestinal tracts as well as the skin. As a consequence seborrhoeic patients are subject to chronic infection of the upper respiratory tract and to dyspepsia and malnutrition.

These conditions, together with the underlying nervous and temperamental instability, have an important bearing on aetiology and treatment.

ANTHRA-SILICOSIS AND BRONCHIAL CARCINOMA WITH QUIESCENT TUBERCULOSIS

BY

J. E. LOVELOCK, B.M., B.Ch.

Late House-physician, Brompton Hospital; late House-physician, St. Mary's Hospital

(WITH SPECIAL PLATE)

The increasing incidence of bronchial carcinoma is a widely recognized fact, and statistical surveys in recent years (Duguid, 1927; Simpson, 1928; Kennaway and Kennaway, 1936) suggest that this increase is real, and not merely an apparent one due to improved facilities for observation and more accurate diagnosis. On the other hand, Hadfield and Garrod (1938), who quote Passey and Holmes (1935), against the majority of available evidence, consider that the matter is still *sub judice*. Study and investigation of the aetiological factors of the disease are therefore of particular importance, and with the growing tendency to regard cancer as the product of prolonged mechanical or chemical irritation of an organ with "constitutional and inherited inferiority" in an individual with an intrinsic susceptibility to neoplastic formation (Hurst, 1939), it is inevitable that attention in bronchial carcinoma should be directed towards inhaled irritants. In a condition such as anthra-silicosis, in which chronic irritation of the bronchi and lung by both coal and silica is the outstanding feature, a relatively high incidence of primary bronchial carcinoma might be anticipated. A number of such cases of cancer associated with silicosis have been reported from all parts of the world during recent years, including the studies by Schmorl (1928) on Schneeberg cancer, in which some factor in addition to mechanical irritation would seem to be most active, and by Pancoast and Pendergrass (1933), Dible

W. HUNTER AND HARRY W. DRYERRE: DUODENAL
ULCER IN THE NEWBORN

FIG. 1.—Artery showing bleeding point in floor of ulcer. ($\times 250$.)

FIG. 2.—Ulcer in posterior wall of duodenum; two vessels in base and normal epithelium at edges. ($\times 15$.)

FIG. 3.—Artery in floor of ulcer showing digested state of wall. ($\times 250$.)

JOHN T. INGRAM: SEBORRHOEIC DIATHESIS

FIG. 1.—Case of Seborrheic eczema of metabolic origin.

FIG. 2.—The same case as above.