M & B 693 RASHES**WITH PARTICULAR REFERENCE TO ACUTE
EXANTHEMATA**

BY

A. R. THOMPSON, M.D., B.S., D.P.H.*Senior Assistant Medical Officer, South-Eastern Fever
Hospital, S.E.14*

(WITH SPECIAL PLATE)

Of recent years the encouraging results attending the introduction of the chemotherapeutic group of drugs have led to their widespread employment in many different types of infection. These new preparations are of special importance in fever hospital practice, not only from the point of view of therapy, but also because they may be associated with eruptions closely simulating the rashes of the common infectious diseases. This applies in particular to 2-sulphanilaminopyridine (M & B 693), now widely prescribed for children with acute respiratory and other infections, and may lead to difficulties in differential diagnosis during future epidemics of the acute exanthemata. Sulphanilamide rashes, which tend for the most part to be morbilliform in type, have been described in detail, in association with the related blood changes, by Schlesinger and Mitchell (1938), but in my experience such eruptions occur less commonly than do those following M & B 693. Recent publications on therapeutic trials with this drug contain but scant reference to the appearance of drug rashes, which are variously referred to as blotchy, erythematous, papular, or morbilliform, at times exhibiting a photosensitive distribution. The usual time of onset of these eruptions appears to be between the eighth and the twelfth days of administration, and although they may occur after shorter or longer periods, it should be noted that the average interval falls within the incubation periods of the common infectious fevers. Three examples in patients recently admitted to the South-Eastern Hospital, each bearing a close resemblance to one of the more common eruptive fevers, are described below.

M & B 693 Rash resembling Measles

A girl 9 years of age was admitted in December, 1938, with acute follicular tonsillitis and right otitis media. She was given M & B 693, two tablets (that is, 1 gramme) every eight hours, and the routine ear toilet was carried out. The throat rapidly became clear, and five days later the ear discharge ceased, although the perforation, which was sealed by a flake of lymph on the ear drum, was not yet healed. Eight days after first taking the drug the child developed a blotchy macular eruption on the face, neck, wrists, and forearms, suggesting photosensitivity, while the trunk showed a crop of fine macules. The features were puffy, but there was no catarrh, Koplik's spots, or lymphadenitis (the child gave a past history of measles). The dosage of M & B 693 was halved, and on the following day the rash became generalized and very heavy, the character and distribution being highly suggestive of measles. Coarse blotchy terra-cotta-coloured macules, confluent on the face, gave a bloated appearance, but there were no catarrhal signs (Plate, Figs. 1 and 2). The temperature rose 2° F., the patient was lethargic and complained of headache but no cough, while the buccal mucosa and fauces were normal. A blood count performed on the following day showed haemoglobin 72 per cent., red cells 3,800,000, colour index 0.94, leucocytes 17,000, with 70 per cent. polymorphonuclear neutrophils, 4 per cent. eosinophils, 23 per cent. lymphocytes, and 3 per cent. monocytes.

On discontinuing the drug the rash gradually faded over a period of eight days, leaving a staining indistinguishable from that which follows a heavy measles rash. Ten days later the

otorrhoea recurred, and a test dose of 1 gramme M & B 693 was administered. This was followed in one and a half hours by headache, vomiting, rise in temperature, and the rapid development of the morbilliform eruption, as described, which became generalized within a period of three hours.

It would appear that this child had developed a marked sensitivity towards the drug, and similar cases have been observed after sulphanilamide (Schlesinger and Mitchell, 1938; Schonberg, 1937). It is not at present known how long this sensitivity may last, but it is possible that administration of this drug to a previously sensitized patient on some future occasion may be attended with alarming symptoms. Recently I observed a pronounced idiosyncrasy in a patient with pertussis bronchopneumonia who developed a rash similar to the one described above, eighteen hours after receiving 1.5 grammes of the drug. Severe stomatitis, cheilitis, and a vesicular eruption on the face were additional features in this case, which resembled the one recently described by Davis (1939) (Plate, Fig. 3). Agranulocytosis was excluded after a blood count which showed red cells, 4,930,000; haemoglobin, 84 per cent.; colour index, 0.85; leucocytes, 31,000, with 86 per cent. polymorphonuclear neutrophils, 8 per cent. lymphocytes, and 6 per cent. monocytes; while a stained film was normal. Smears and cultures from the mouth and fauces were also negative for Vincent's organisms, streptococci, and diphtheria bacilli. The rash was followed by heavy and persistent staining and a generalized desquamation of the "pin-hole" type. The child ultimately recovered.

M & B 693 Rash resembling Rubella

This rash is typical of the more common type of toxic eruption following M & B 693.

A girl aged 4 was admitted on March 13, 1939, suffering from pertussis in a moderately severe form. There were marked signs of pulmonary catarrh, and she was given M & B 693, 0.5 gramme eight-hourly. Ten days later the temperature rose to 100° F., and a well-marked finely macular roseolar eruption developed, the distribution being general, with a tendency to become confluent on the arms (Plate, Fig. 4). The tongue and fauces were normal, but the superficial lymph glands generally were palpable. There was a previous history of measles. On discontinuing the drug the patient became afebrile and the rash completely faded in forty-eight hours. On March 28 a test dose of 1 gramme of M & B 693 was administered, and this was followed the same evening by a rise in temperature to 103° F. and a profuse recurrence of the previous type of rash. She also became cyanosed and complained of cramps in the legs and feet. The following morning the temperature fell to normal, the rash had become scarlatiniform in type on the trunk, and there were tingling sensations in the feet. Twenty-four hours later the rash and paraesthesiae had vanished, and her subsequent progress was uneventful.

With regard to these drug eruptions in whooping-cough patients, the writer had on two occasions recently observed the rash to be petechial on the face but not elsewhere, and in only one of these cases was the attack severe.

**M & B 693 Rash resembling Scarlet Fever and followed
by Acute Exfoliative Dermatitis**

The patient, a man of 21, had an appendix abscess drained at another hospital on January 19, and between January 28 and February 16 he was given M & B 693 in 1-gramme doses thrice daily. He had also been having ultra-violet light treatment three times a week, the last exposure being given on February 20. On February 15 the patient developed a generalized morbilliform rash, which subsided and was followed by a rise in temperature and a generalized scarlatiniform eruption of an irritating character. He was admitted to the South-Eastern Hospital with a diagnosis of

scarlet fever on February 23, when his temperature was 102° F. An intense confluent erythema was present on the trunk and a maculo-papular rash on the distal portions of the extremities. There was slight generalized subcutaneous oedema with puffiness of the face, eyelids, and pinnae, and some branny desquamation of the face and scalp. Submucous haemorrhages were seen in the buccal mucosa, but the tongue and fauces were not suggestive of scarlet fever. The Dick and Schultz-Charlton reactions were negative, and repeated swabs from the throat, nose, and appendix wound were also negative for haemolytic streptococci. The urine was normal throughout. Two days after admission an irritating eruption of miliary vesicles appeared on the trunk, and the patient remained febrile for five days. The cutaneous oedema, erythema, and branny desquamation persisted, developing into a generalized exfoliative dermatitis which later involved the palms and soles, with much dryness and irritation of the skin, made worse by exposure to direct sunlight. Three weeks later he was discharged quite fit apart from the skin condition, which persisted in a modified form. There was no history of asthma or urticaria.

The morbilliform-scarlatiniform sequence in the character and development of eruptions due to M & B 693 has been described by others, though it is probable that in this case exposure to ultra-violet light had aggravated the condition. Photosensitivity in association with the sulphanilamide group of drugs has been repeatedly noted, and severe rashes of a similar nature (scarlatiniform) have been recorded by Newman and Sharlit (1937) after sun-bathing. Recently Hallam (1939) has described a severe cutaneous and general reaction following exposure to ultra-violet light in a patient who had been receiving M & B 693, and he lays emphasis on the dangers attendant on such a procedure.

Summary

Attention is drawn to the commoner types of drug rash following treatment with M & B 693 and their possible confusion with the eruptive fevers, especially measles, rubella, and scarlet fever.

It is suggested that, during epidemic prevalence of these diseases in the future, a careful history with regard to the administration of this drug or similar compounds would be of great assistance in differential diagnosis.

Reference is made to the acute reactions, accompanied by a rash, which may occur in patients previously sensitized to these drugs, and also to the harmful effects of actinotherapy in such cases.

I am indebted to Dr. G. W. Ronaldson, medical superintendent of the South-Eastern Hospital, for his helpful advice, and to Sir Frederick Menzies, medical officer of health to the County of London, for permission to publish this paper.

REFERENCES

- Davis, E. (1939). *Lancet*, 1, 1042.
Hallam, R. (1939). *British Medical Journal*, 1, 559.
Newman, B. A., and Sharlit, H. (1937). *J. Amer. med. Ass.*, 109, 1036.
Schlesinger, E. R., and Mitchell, W. L. (1938). *Amer. J. Dis. Child.*, 56, 1256.
Schonberg, I. L. (1937). *J. Amer. med. Ass.*, 109, 1035.

The committee appointed to plan the distribution of the fund, amounting to 5,000,000 kronor (£258,000), which King Gustav V of Sweden received as an eightieth birthday gift from the Swedish people, and which at the King's directions was to be applied for the investigation and combating of disabling diseases—principally infantile paralysis and rheumatic ailments—has now submitted its proposals. It is proposed that the money be applied to three funds: a research fund of 2½ million kronor, an institute fund of 2 millions, and the remainder to constitute a reserve fund. Research into the origin and modes of development of the diseases in question is to be promoted above all, and the money should not be to any great extent used for treatment of sufferers.

A RARE TUMOUR OF THE TONGUE

A MANIFESTATION OF POST-KALA-AZAR DERMAL LEISHMANIASIS

BY

S. K. GHOSH DASTIDAR, M.B., D.T.M.,
M.R.C.P.Lond. & Ed.

Lecturer in Medicine, Prince of Wales Medical College,
Patna, India

(WITH SPECIAL PLATE)

Brahmachari (1922) reported a case with nodular skin lesions simulating leprosy as a sequel of kala-azar. He demonstrated *Leishmania donovani* both in direct smears and in culture, and described the condition as "dermal leishmanoid." Acton and Napier (1927) preferred the term "post-kala-azar dermal leishmaniasis," which, though somewhat unwieldy, is self-explanatory; nevertheless it is often referred to in the literature and in general practice as "dermal leishmaniasis." They described three distinct clinical types: an early depigmented stage; a later nodular stage; and a xanthoma type. Brahmachari and Banerjee (1928) described another type in which the disease showed itself in the form of erythematous patches with nodules before the appearance of depigmentation. Napier and Das Gupta (1930) in their extensive study of this condition recognized the following clinical types: depigmented areas, erythema rash, nodules, verrucose type, papillomatous type, hypertrophic type, and xanthoma type. In the hypertrophic lesion they demonstrated *Leishmania donovani* in the smears obtained from the scrapings of mucous membrane inside the lip and nose, where it was continuous with the skin. Subsequently, in their further clinical observations, Napier and Das Gupta (1934) described four more clinical types in addition to those mentioned before: mucous membrane lesion (on the palate, etc.); small ulcerated lesion; extensive erythema; and perionychial induration. Smith and Halder (1935) added three more types—namely, fibroid, leprosy, and ulcerative; the latter two were really variants of the nodular type.

Though the typical lesions are so characteristic and variable as to be bewildering, I do not think we have as yet reached the journey's end. The greater our experience of the condition the more numerous the clinical types become. During the last four years I have made or collected notes on eleven cases of post-kala-azar dermal leishmaniasis; these eleven cases include a new type—nodular swelling of the tongue as a rare manifestation.

Case Report

A Mohammedan man aged 30, who was a local deck-hand by occupation, sought advice for nodular swellings in the tongue, and was admitted into the Patna Medical College Hospital on April 26, 1938. The nodules appeared in the tongue about four months earlier. He gave a history of suffering from a long-continued fever about two years previously, with enlargement of the spleen and liver, lasting for nearly three months, and was treated with a few injections of neostibosan. He stopped the injections as soon as the temperature came down and the general condition improved. The total quantity of neostibosan injected was approximately 1 gramme. About six months before admission he noticed a few small nodules appearing first on the face. Two months later he felt a sense of heaviness in the movement of the tongue, along with a slight swelling which appeared on the left side of the dorsum (of the tongue) close to the medial line. The lingual nodules (see Plate) were rather big and close together; they were painless and soft, had no ulceration on the surface, and showed

A. R. THOMPSON: M & B 693 RASHES, WITH PARTICULAR REFERENCE TO ACUTE EXANTHEMATA

FIG. 1.—Morbilliform rash due to M & B 693 appearing on the eighth day of administration and photographed two days later. Note general distribution and coarse macular character of the eruption, which tends to be confluent on the arms and other areas exposed to sunlight.

FIG. 2.—Same patient as Fig. 1. Note bloated features but absence of catarrh.

FIG. 3.—Severe reaction after M & B 693 in a sensitive patient. Heavy semi-confluent morbilliform rash becoming vesicular on the face, with severe stomatitis and cheilitis.

FIG. 4.—Rubella-like eruption appearing ten days after first giving M & B 693. The macules are finer and less deep in colour than in Figs. 1 and 2. There is some suffusion of the conjunctivae and a tendency to confluence of the rash on the extremities normally exposed to sunlight. The following day the eruption became scarlatiniform on the trunk.