SLEEP CHARTS

BY

G. de M. RUDOLF, M.R.C.P., D.P.M.

Although the necessity for an adequate quantity of sleep is generally recognized, charts for recording the amount do not appear to exist. A standard quantity of sleep cannot be assigned for any person or age, and tables

the patient is essential. The nurse is therefore instructed to make a slight sound on approaching the bed and the patient is asked to open the eyes, raise a hand, or move the head, to show that he is awake. If the patient fails to co-operate, observation should be stopped. Consequently the chart is unsuitable for semi-conscious patients or for those who are markedly defective or psychotic.

The charts are printed for me by Messrs. John Wright and Sons Ltd. of Bristol. They were drawn up in con-

SLEEP CHART

NAME		SEX		DATE OF ADMISSION		AGE ON ADMISSION																					
DIAGNOSIS										Notes, Medicines, Apparent Causes of Sleeplessness, etc.																	
DATE	TIME	12.30	1.0	1.30	2.0	2.30	3.0	3.30	4.0		4.30	5.0	5.30	6.0	6.30	7.0	7.30	8.0	8.30	9.0	9.30	10.0	10.30	11.0	11.30	12.0	Total
	A.M.																										
	P.M.																										
	A.M.																										
	P.M.																										
	A.M.																										
	P.M.																										
	A.M.																										
	P.M.																										
	A.M.																										
	P.M.																										
	A.M.																										
	P.M.																										
	A.M.																										
	P.M.																										

INSTRUCTIONS

1. The patient should be instructed to open the eyes, raise a hand, or move the head on the approach of a nurse.
2. Observations should be made by approaching the bed and making a very slight sound, such as rustling the apron. The patient should not be touched or addressed.
3. If the patient appears to fail to co-operate when awake, observations should be discontinued.
4. When the patient is asleep a dot should be placed in the appropriate space; when awake, a diagonal line thus . If the patient is both awake and asleep after a half-hourly observation, a dot and half a diagonal line should be placed in the space of the next observation, the second sign corresponding to the state of the patient at the second half-hourly observation.

setting out the amount required at each age appear to be fallacious: means are all that can be relied upon. Thus Laird found, in 509 men, that the hours of sleep obtained varied from five to ten hours a night, with only slight differences between age groups 25 to 75 years. The hours at which sleep takes place may be of more importance. Stoeckmann claims that sleep before midnight is more restorative than sleep after, thus confirming the old-standing belief. Depth of sleep, as measured by auditory stimuli required to waken the sleeper, varies during the sleep period, being greatest within the first and less during the last part.

In practice the quantity of sleep and the hours during which sleep is obtained vary with each individual, some requiring less, others more, but hitherto there has been no means of readily recording how long an individual sleeps. For instance, nurses report that a patient has had a good night. This may mean that, (a) as compared with the nurse's own idea of her own sleep, the patient has slept as much or more; (b) the patient has slept more than is usual for him; (c) he has slept longer than other patients; or (d) he has lain quietly awake with his eyes closed and the nurse has been mistaken in assuming he was asleep.

In order to attempt to obtain reasonably accurate records of sleep a chart has been drawn up divided into half-hourly observations. The observer must place a dot or a stroke in each square. Each record is thus an individual note, so that the observer cannot make one stroke for several observations. The night is broken up into two lines, so that the observer's train of thought is also broken. As, especially in darkened wards, a patient may lie awake but appear to be asleep, the co-operation of

nexion with a Fellowship of the Colston Research Society, to which my thanks are due.

The following medical promotions in and appointments to the Venerable Order of the Hospital of St. John of Jerusalem are announced in the *London Gazette* of June 23. As Knight: Urban Marks, M.R.C.S., L.R.C.P. As Commanders: Lieutenant-Colonel F. Romer, M.R.C.S., L.R.C.P., Sir Henry J. Gauvain, M.D., F. C. Bottomley, O.B.E., M.D. As Officers: H. G. Ramsbottom, L.R.C.P. and S.Ed., N. H. Kettlewell, L.M.S.S.A., E. J. Burnett, M.B.E., M.B., D. H. Griffiths, M.R.C.S., L.R.C.P., G. L. Pierce, L.R.C.P. and S.Ed., R. J. Clarke, M.B., Lieutenant-Colonel E. S. Goss, M.C., M.R.C.S., F. J. Bach, M.D., J. C. Glen, L.R.C.P. and S.Ed., Major E. S. B. Hamilton, M.C., M.B., E. Hoerman, M.B., A. Croll, M.D., F.R.C.S.Ed., E. F. White, F.R.C.S., Major C. Wilson, M.B., C. P. Gaboury, M.D., C. J. A. McKillop, M.D., Major W. E. L. Sparks, M.D., Lieutenant-Colonel R. H. Candy, C.I.E., Lieutenant-Colonel N. Briggs, Jane P. H. G. Brown, M.B. As Associate Officers: Captain M. Hafizuddin, L.R.C.P., M.R.C.S., Lieutenant-Colonel A. N. Bose, O.B.E., M.D. As Serving Brothers: F. L. Richard, M.B., J. Wells, M.R.C.S., L.R.C.P., J. Lamberton, M.B., A. H. Penistan, M.B., Lieutenant-Colonel W. George, M.C., T.D., M.B., Major C. M. Forbes, M.B., J. W. Watterson, M.B., Major N. W. Hammer, M.R.C.S., L.R.C.P., W. D. Jenkins, M.R.C.S., L.R.C.P., E. C. Malden, M.B., E. R. Weaver-Adams, M.B., J. Macfie, M.B., W. C. Lyons, M.B., I. Way, M.B., H. W. Ward, M.B., A. R. Home, M.B., A. U. Sheth, M.B. As Associate Serving Brothers: J. Lovius, M.B., Major D. N. Gupta, L.M.S., P. C. Datta, F.R.C.S., P. N. Gokhale, F.R.C.S., M. C. Madhok, M.B. As Serving Sister: Sara L. Westlake, M.B.