following the introduction of the immunization, while eight cases with two deaths were recorded among the small proportion of non-inoculated workers (Grasset, Lewin, and van der Merwe, 1937).

With a view to the possibility of reinforcing the immunity so produced investigations were undertaken with an alum-precipitated endotoxoid prepared on lines similar to the principle elaborated by Glenny for diphtheria immunization. Such precipitated endotoxoid confers upon mice a protection against *Vi B. typhosus* at least as strong as that conferred by unmodified endotoxoid; in man, without causing undue reaction, it is followed by an agglutinin formation to titres such as 1/800 of H and 1/200 of O during the month following the inoculation.

Recent investigations on the nature of typhoid endotoxoid carried out in collaboration with H. D. Barnes—to be incorporated in a separate publication—have shown that by means of alcoholic precipitation a purified antigen is obtained. On being redissolved in saline the precipitated material confers on the mice and rabbits a degree of immunity and an agglutinin titre equal to that afforded by the untreated endotoxoid. These results are in agreement with those obtained by Topley, and Boivin in France, with the lipide-polysaccharide complex, isolated by various methods from the *B. typhosus*. Immunization of man has been undertaken with this purified endotoxoid, and this procedure also results in the formation of H and O agglutinins to considerable titre.

With regard to the duration of the immunity produced by endotoxoid, serological investigations show that a very appreciable H and O agglutinin titre is still present at the end of the first year in the blood of the persons inoculated. Epidemiological data indicate that there is present in the second year a degree of protection practically equivalent to that observed during the first year.

The introduction of typhoid endotoxoid is too recent for an actual limit of protection to be fixed. That information can only be accumulated over a period of years by epidemiological data and by a close study of the greatest possible number of subjects who have been immunized by this new form of vaccine.

REFERENCES

- De Villiers, J. J. (1937). *S. Afr. med. J.*, **11**, 221.
 Felix, A., and Bhatnagar, S. S. (1935). *Brit. J. exp. Path.*, **16**, 422.
 — and Pitt, R. M. (1934a). *J. Path. Bact.*, **38**, 409.
 — (1934b). *Lancet*, **2**, 186.
 — (1935). *J. Hyg., Camb.*, **35**, 428.
 Grasset, E. (1933). *C. r. Soc. Biol., Paris*, **113**, 20.
 — (1935). *S. Afr. med. J.*, **9**, 208.
 — (1937). *Int. Ass. prev. Pédiat.*, 6th Conference, Rome, 69.
 — and Gory, M. (1927). *C. r. Soc. Biol., Paris*, **97**, 1211.
 — and Lewin, W. (1936). *Brit. J. exp. Path.*, **17**, 179.
 — (1937). *C. r. Soc. Biol., Paris*, **124**, 331.
 — and van der Merwe, T. (1937). *S. Afr. med. J.*, **11**, 660.
 Lister, F. S. (1917). *Publ. S. Afr. Inst. med. Res.*, **1** (No. 10), 303.
 Lister, Sir Spencer, and Ordman, D. (1935). *Ibid.*, **7** (No. 37), 1.
 Meyerstein, R. (1938). *S. Afr. med. J.*, **12**, 324.
 Orenstein, A. J. (1937). *Ibid.*, **11**, 401.
 Perry, H. M., Findlay, H. T., and Bensted, H. J. (1933a). *J. R.A.M.C.*, **60**, 241.
 — (1933b). *Ibid.*, **61**, 81.
 — (1934). *Ibid.*, **62**, 161.
 Topley, W. W. C., et al. (1937). *Lancet*, **1**, 252.
 Transvaal Mine Medical Officers' Association. Report of Meeting (1937). *Proc. Transv. Mine med. Off. Ass.*, **17**, 44.
 Van Blommestein, J. H. G. (1936). *Ibid.*, **16**, 9.

Dr. Nathan Bristol Van Etten, medical director of New York City's Morrisania Hospital, has been chosen president-elect of the American Medical Association, and will be inducted into office in June, 1940, when the Association meets in New York. At the recent annual meeting in St. Louis the attendance was 7,412.

SUDDEN BLINDNESS

BY

JOSEPH MINTON, F.R.C.S.

Ophthalmic Surgeon to Out-patients, Hampstead General Hospital; Ophthalmic Surgeon, Queen's Hospital for Children; Assistant Ophthalmic Surgeon, West End Hospital for Nervous Diseases; late Research Scholar, Royal Eye Hospital, London

Of sudden loss of sight of one or both eyes vascular lesions are the most common cause. Von Graefe reported the first cases of embolism of the central artery of the retina in 1859, and for many years this was accepted as the only possible cause of obstruction of the central retinal artery. The great advance in the study of ophthalmoscopic appearances of the retinal arteries and veins has enabled us to understand more fully the pathological changes which underlie the occlusion or obstruction of the retinal artery and the sudden loss of sight that ensues. George Coats in 1906, as a result of pathological research, came to the conclusion that obliterative endarteritis, with or without thrombosis, can produce a sudden obstruction of the central artery or of any of its branches, and that this form of obstruction is a much more likely event than embolism.

Spasm of the retinal arteries was first observed in 1874 by Raynaud in a patient suffering from the disease which now bears Raynaud's name. Since this date angiospasm of the retinal arteries with associated temporary blindness has been studied by many observers and proved to occur in lead and quinine poisoning, epilepsy, and migraine. In recent years Mylius has given an account of ophthalmoscopic observations in eclampsia and pre-eclampsia, in which conditions temporary visual disturbances due to spasmodic contractions of the retinal arteries are fairly common and may be one of the earliest diagnostic features. The observations of Mylius and Wagener, that the retinal arterioles may show both localized and generalized spasm in the toxæmia of pregnancy, suggest that these are probably general throughout the arteriolar system and are closely related to hypertension. The investigations of Wagener and Keith confirm Allbutt's original conception that hypertension is primarily due to spastic constriction of the peripheral arterioles, resulting in a dynamic increase of resistance.

Hypertension

The work of Dumas, Wagener, and Keith has led to the adoption of a subdivision of hypertension into three phases. Sudden blindness due to the occlusion of the retinal arterioles has been observed in all stages of hypertension.

THE FIRST PHASE

In the first or benign phase of hypertension the patients may be in good general health and have very few symptoms. The retinal arterioles are usually narrowed and may show irregularity of lumen as a result of tonic contraction with superadded localized spasm. The patients may complain of transitory blindness affecting one or both eyes, this blindness lasting a few minutes to a few hours. The sight is usually regained, but in many cases such attacks are followed by permanent blindness of the affected eye. These transitory attacks are due to a sudden spasmodic contraction of the retinal arterioles, which may subside quickly, having caused a temporary anaemia of the retina. Prolonged attacks lead to a post-spastic thrombosis of the affected vessel and permanent damage

of the retina. The ophthalmoscopic appearance of the fundus in the case of thrombosis of the central artery of the retina is the same as the one known to occur in embolism of the central retinal artery: a cherry-red spot at the macula surrounded by a whitish haze, pale disks, general pallor of the fundus, with threadlike retinal arteries. On many occasions one branch of the retinal artery only may be involved, and then one sees an occlusion of the affected branch and pallor of the surrounding retina. The superior temporal is the branch most frequently affected. In the case of occlusion of a branch of the retinal artery the patient may notice the loss of a portion of the field of vision in the affected eye, and on perimetric examination a sector-shaped defect will be discovered.

THE SECOND PHASE

In the second, more advanced, stage of hypertension the patient may have symptoms suggesting coronary, cerebral, or renal arteriosclerosis. The retinal arterioles show the well-known features of arteriosclerosis: crushing and kinking of veins at arteriovenous crossings, tortuous and narrowed arteries with irregular lumen, and occasional retinal haemorrhages and exudates. In this stage of hypertension sudden transitory or permanent blindness is most common. The sudden obstruction of the retinal artery or its branches is due to a localized thrombosis resulting from endarteritis obliterans and the spastic state of the diseased retinal arterioles. In this stage of hypertension, thrombosis of the central vein of the retina or its tributaries is also most common. In the latter condition the fundus presents the well-known picture of flame-shaped haemorrhages and exudates around the disk or, in the case of thrombosis of a tributary, the haemorrhages are confined to the part of the retina drained by the vein. In thrombosis of the central vein the patient becomes aware of a sudden loss of sight in the affected eye. Thrombosis of the tributaries of the central retinal vein is more common than that of the main trunk, and the loss of sight depends on the position of the tributary. In the case of thrombosis of the nasal tributaries the patient may not be aware of any loss of vision. On the other hand, thrombosis of the superior temporal or inferior temporal tributaries draining the central area of the retina leads to an appreciable loss of central vision, which the patient notices and for which he seeks advice.

THE THIRD PHASE

The third stage of hypertension was described by Wagener and Keith in 1928 and called by them the malignant hypertension syndrome. The patient usually complains of dyspnoea on exertion, asthenia, and visual disturbances. The fundi show, in addition to the above-described changes of the arteries and veins, multiple haemorrhages and exudates, oedema of the retina, and papilloedema. The arterioles reveal marked spasm and narrowing of the lumen resulting from advanced endarteritis obliterans. These fundal appearances are typical of severe angiospastic retinitis. Wagener and Keith followed up eighty-five patients who were diagnosed as suffering from malignant hypertension, and found that 78 per cent. of them had died within a year of the diagnosis. The vision of patients suffering from malignant hypertension is usually much affected as a result of haemorrhages and exudates of the central retinal areas. Attacks of transitory blindness are common in this group.

In 1937 I reported on my clinical study of fifty-four cases of occlusion of the central artery of the retina. In the course of this investigation I found that the condi-

tion occurred most frequently above the age of 40: of the fifty-four patients only five were under that age, and of these four had valvular disease of the heart. The majority of those examined showed evidence of arteriosclerosis of the non-affected eye and general hypertension. A number suffered only from temporary blindness, which had resulted from a temporary anaemia of the retina following an active spasm of the retinal arterioles, the sight returning on the disappearance of the arteriolar spasm. In the above-described group of fifty-four patients only three developed bilateral blindness as a result of bilateral occlusion of the central artery of the retina; in the other fifty-one cases the condition was unilateral. In the four young patients with valvular disease of the heart the sudden obstruction of the retinal artery was due to an embolus blocking the central retinal artery.

The other young patient, a woman aged 23, lost the sight of the right eye when getting out of bed on the tenth day after confinement with her second child. She was feeling quite well at the time. On examination no evidence of heart disease was found, and it might be assumed that a thrombus from the pelvic veins was carried to the right ventricle and through a patent foramen ovale to the left side of the heart and from there to the retinal artery.

Sudden loss of sight has been described after severe gastric or rectal haemorrhages, presumably due to a retinal anaemia. One of the patients in this series permanently lost the sight of the right eye on the day following a severe haematemesis. He did not seek advice until six weeks after the attack of blindness. Two patients lost the sight of one eye as a result of a spasm of the retinal arteries.

The first of these, a man aged 52 suffering from typical Raynaud's disease, suddenly lost the sight of both eyes. The left eye recovered its vision, but the right one became blind as a result of post-spastic thrombosis of the retinal artery. The other patient, who suffered from severe migraine, suddenly lost her sight while walking in the street. For a few minutes she could not see with either eye. After a short while she recovered the sight of the left eye, but the right eye became permanently blind.

Results of the Investigation

In this clinical investigation I found that thrombosis of the retinal vein or its tributaries is four times as common as occlusion of the retinal artery. The investigation was carried out at the Royal Eye Hospital, London, and during the period 1931-6, in the out-patient department there, 205 patients were diagnosed as suffering from thrombosis of the retinal vein or its tributaries, while during the same period only fifty-four were seen with occlusion of the retinal artery or its branches.

On reviewing the results of this clinical study one is driven to the conclusion that obstruction of the central retinal artery, which is so often described in textbooks as embolism of the central artery of the retina and said to be associated with valvular disease of the heart and infective endocarditis, is only rarely due to blockage by an embolus. In the vast majority of cases the obstruction is caused by a localized thrombosis in spastic and thickened arteries, occurring in patients suffering from hypertension. A number of these patients remained in good health for many years, with no evidence of disease of the coronary or cerebral arteries. In a follow-up of this group it was found that one-half of them were alive several years after the onset of the sudden blindness. One can therefore conclude that in hypertension the arteriolar sclerosis may follow a patchy and irregular distribution and that the involvement of the retinal arteries does not in all cases

indicate disease of cerebral or coronary vessels, which may or may not be affected later in the long course of the disease.

Treatment

Antispasmodic treatment should be administered to patients seen a short time after the onset of retinal artery obstruction. Among the measures used the following have been found the most successful: subconjunctival injections of acetylcholine, combined with hypodermic injections of the same drug; amyl nitrite inhalations and nitroglycerin tablets may also be useful. In addition, the eye should be massaged at frequent intervals to reduce intraocular pressure. In grave cases not responding to the above treatment, paracentesis of the eye is carried out with a view to reducing intraocular pressure, and thus allowing a dilatation of the retinal vessels. Reports of success following these measures—spasm of the artery subsiding and the vision returning—are often found in ophthalmic literature. The following recent case is instructive.

A man aged 54 suddenly noticed the loss of vision of the right eye when getting out of bed. He applied for treatment six hours later. He had had in the past several attacks of temporary blindness, which never lasted longer than five to ten minutes. On examination the following was found: Vision of the right eye was shadows only. Fundus: disk pale, constricted retinal arteries, and pallor of the retina. With the left eye vision was 6/9, and the fundus presented a typical picture of advanced retinal arteriosclerosis. The patient's blood pressure was 180/100.

Treatment.—Amyl nitrite inhalations were ordered, and the eye was massaged. This had no effect. A subconjunctival injection of acetylcholine (5 minims) was given; this was followed within half an hour by a gradual return of sight. The patient began to discern objects, and after three hours read 6/36. A second subconjunctival injection of acetylcholine (5 minims) was then given. On the following day the patient read 6/18, but had an inferior nasal sector-shaped defect of the field of vision, which became permanent. In this case the prolonged anaemia of the retina had caused partial damage to the central region and the superior temporal quadrant of the retina, supplied by the superior temporal branch of the retinal artery. The patient later received general treatment for arteriosclerosis.

Conclusions

Attacks of transient blindness should be regarded as a warning of vascular disease. In the diagnosis and prognosis of this disease the study and correct interpretation of the accompanying retinal changes are essential.

BIBLIOGRAPHY

- Allbutt, T. C. (1896). *Trans. Hunterian Soc.*, 38-57.
 Bonamour, G. (1936). *Étude clinique du fond d'œil dans l'hypertension artérielle*, Cohendet Frères, Lyons.
 Coats, G. (1906). *Roy. London ophthal. Hosp. Rep.*, 16, 262.
 Dumas, A. (1931). *Presse méd.*, 39, 129.
 Friedenwald, H. (1930). *Trans. ophthal. Soc. U.K.*, 50, 452.
 von Graefe (1859). *Arch. f. Ophthal.*, 5, Abt. 1, 136.
 Gunn, R. M. (1892). *Trans. ophthal. Soc. U.K.*, 12, 124.
 Keith, N. M., Wagener, H. P., and Kernohan, J. W. (1928). *Arch. intern. Med.*, 41, 141.
 Minton, J. (1937). *Proc. roy. Soc. Med.*, 30, 285.
 Moore, R. Foster (1916). *Trans. ophthal. Soc. U.K.*, 36, 319.
 — (1924). *Retinal Venous Thrombosis*, London.
 Mylius, K. (1928). *Ophthal. Gesellsch., Heidelberg*, 47, 379.
 Orr, H. C., and Young, J. H. (1935). *British Medical Journal*, 1, 1119.
 Redslob, E. (1926). *Ann Oculist.*, Paris, 163, 169.
 Wagener, H. P., and Gipner, J. F. (1927). *Amer. J. Ophthal.*, 10, 650.
 — and Keith, N. M. (1937). "Diffuse Arteriolar Disease and Hypertension," 15th Int. Ophthal. Congress, 1937.

NEUROSES AMONG COMBATANT TROOPS IN THE GREAT WAR*

BY

FREDERICK DILLON, M.D.

Medical Superintendent, Northumberland House Mental Hospital; Chief Assistant in the Department of Psychological Medicine, University College Hospital

Recent events would make it appear that we may be called upon to use the experience obtained during the great war. It seems advisable, therefore, to refresh our knowledge of the disorders associated with warfare and of the methods of dealing with them. A series of six lectures on the war neuroses has recently been delivered at the Tavistock Clinic, and the following account, which has not before been published, of experience in the treatment of these conditions in France from 1916 onwards may, I hope, prove also to be helpful.

Functional Nervous Disorders

The functional nervous disorders comprised a considerable proportion of the total casualties. From the statistics given in the *Official Medical History of the Great War*,¹ which are admittedly incomplete, it is not possible to obtain an accurate idea of the comparative wastage from this source. But an estimate was made shortly after the war had ended that one-third of the unwounded and one-seventh of those discharged from the British Army were permanently unfit on account of functional neuroses and mental disorders. For the first two years of the war all cases of this kind were evacuated to England and treated there. This was partly due to the erroneous opinion, widely accepted at that time, that these conditions, to which the misleading name "shell-shock" had been applied, were largely or wholly brought about by damage to the nervous system as the result of shell explosions and similar agents. It was commonly held that a shell exploding near an individual must inevitably produce injury to the nervous system, from the liberation of the tremendous forces of compression and decompression. Cases were described of death following an explosion in which no external marks were visible on the body but in which punctate haemorrhages and evidences of violence to the nervous system were found on post-mortem examination. It was only gradually realized that these were exceptional cases and that every day, in and near the trenches, men were blown up and buried who showed no sign of any injury to the nervous system and who were able to continue on duty without being sent to hospital. There was, further, the increasing evidence that conditions precisely similar to "shell-shock" occurred in men who had never been subject to the action of an explosive. Results of treatment in England had proved very disappointing; the tendency to fixation of the symptoms and the development of an attitude of chronicity had been only too apparent, and a great many of these men had been discharged from the Army and pensioned.

It was in the attempt to stem the loss from this source that the advanced centres in France were established towards the end of 1916. The experiment had previously been tried in the French Army and had met with encouraging success. I was appointed "shell-shock" specialist to the 3rd Army in December, 1916, and sent from an infantry regiment to a stationary hospital on the lines of communication. There I found three or four cases with functional nervous symptoms in an ordinary medical ward. There was no clinical room in the hut which formed the ward, and no place where the patients could be examined

* The facts on which this article is based are taken from a thesis presented for the M.D. at Edinburgh University. Permission has been obtained for publication.

¹ *Official History of the Great War: Medical Services*, 1931, Final Volume, p. 116.