TREATMENT OF HAEMORRHOIDS*

BY

W. B. GABRIEL, M.S., F.R.C.S.

Surgeon, Royal Northern and St. Mark's Hospitals

(WITH SPECIAL PLATE)

The choice of treatment of haemorrhoids depends so much on examination and an accurate estimation of the condition presented in each particular case that I should like to say a few words on these preliminaries.

Distinction between Internal and External Haemorrhoids

The distinction between internal and external haemorrhoids is of the first importance. Internal haemorrhoids arise from the columns of Morgagni as the result of dilatation and engorgement of the internal haemorrhoidal plexus, and are covered by mucous membrane which is at first of the normal bright pink colour. This colour may be changed by superficial inflammation, ulceration, or sloughing. Later, as the result of repeated prolapse and surface irritation, a large internal pile often shows an overgrowth of squamous epithelium over its lower pole. It may be noted that we see exactly the same change quite commonly over the mucous membrane of the appendix when exposed as an appendicostomy. An internal pile may thus become considerably covered by squamous epithelium, and therefore when prolapsed it somewhat resembles on superficial examination an external skin tag, but the groove which originally marked the muco-cutaneous junction, due to the attachment of the terminal fibrous expansion of the longitudinal muscle, persists in most cases and thus separates it quite clearly from an external pile.

As opposed to this an external pile is covered by skin *ab initio*, whether it is in the stage of an acute thrombosed external pile (anal haematoma) or a soft anal skin tag. Under certain conditions, however, an external pile must be carefully inspected to discover its true nature. Thus an acute haematoma may be seen at the stage when the skin has become ulcerated so that blood and clot are being discharged, and it has on cursory inspection a superficial resemblance to a prolapsed internal pile. Again, if a large mass of external thrombosis and oedema is present careful examination is often needed to detect the lower pole of a prolapsed and thrombosed internal pile in the same quadrant, which is really the "villain of the piece." The anal margins must be separated and perhaps some traction in a lateral direction applied to the haematoma, when the lower pole of the thrombosed internal pile, if present, will at once be seen.

Treatment of External Piles.—Excision of a solitary tense anal haematoma under local anaesthesia is the most common, and a very satisfactory, anal operation. A large conglomerate type of anal haematoma is best treated on expectant lines. Skin tags at the anal margin may subside with lead lotion compresses, but if large should be excised under local anaesthesia. When associated with prolapsing internal piles (intero-external type) a formal haemorrhoidectomy is the best treatment.

Degree of Internal Haemorrhoids

The treatment of internal haemorrhoids is intimately linked up with a correct assessment of the degree of prolapse. Ernest Miles (1919) divided piles into three stages.

Primary, being the earliest stage, in which bleeding takes place at defaecation: the pile may just become grasped by the anal sphincters during the act of defaecation, but no prolapse outside the anal margin occurs.

Intermediate, when the pile protrudes through the anal orifice at defaecation and then becomes reduced spontaneously when the expulsive effort has ceased. Some bleeding still occurs, but may be less than before owing to some fibrosis taking place in the substance of the pile, together with some thickening of the mucous membrane.

The final stage is reached when the prolapse is still more pronounced and does not become spontaneously reduced at the end of defaecation; the pile remains in the prolapsed and engorged state, and digital reduction is needed after each act of defaecation. Sometimes such a pile is irreducible by reason of mere bulk, massive thrombosis, or sphincter relaxation. Even after being properly reduced, recurrence of prolapse takes place subsequently on slight provocation, such as coughing, sneezing, any lifting effort, walking or standing for any length of time, or even passing flatus.

These three stages are now more commonly known as the first, second, and third degrees. It is essential to recognize the symptoms associated with piles of the third degree: mucoid discharge and bleeding into the clothing during the day are common complaints in addition to the necessity for digital replacement.

Routine Diagnostic Methods

Inspection, digital examination, and proctoscopy are required in all cases. The patient should be asked to strain down as the proctoscope is withdrawn in order that the full degree of prolapse may be seen, and very often a bleeding point is revealed as the piles become engorged.

The presence of fibrosis in an internal pile, so that it is palpable to a finger, is evidence of recent or old thrombosis. Large prolapsing piles can almost invariably be rolled under the examining finger, and if the process has advanced to the stage of formation of a fibrous polyp this can readily be felt and can usually be hooked down until it prolapses outside the anus. A pulsating branch of the superior haemorrhoidal artery can sometimes be felt in relation to one or more of the primary internal haemorrhoids, and is a clear indication of the need for a ligature operation.

Other associated ano-rectal conditions must be accurately diagnosed. Fissure in ano is one of the most important, and if present may have considerable influence on the choice of treatment. Of the many other conditions which may coexist with haemorrhoids it is necessary to mention the not infrequent association of carcinoma of the rectum: hence a high digital examination and sigmoidoscopy with this in mind should be done in all cases.

The choice of treatment is necessarily much influenced by finding a pelvic tumour of uterine or ovarian origin, and in men prostatic enlargement often introduces a difficulty in deciding on the right line of treatment. Abdominal conditions such as cirrhosis will affect treatment. In many cases I have seen a troublesome degree of piles in patients who were taking intensive alkali treatment for duodenal ulcer, and this combination is a difficult one to treat; so also is any degree of haemorrhoids in patients with chronic cardiac or pulmonary disease. The general state of patients must always be carefully gone into, particularly if surgical treatment is proposed.

As a statement of the practice of St. Mark's Hospital I will now describe three treatments for internal haemorrhoids, each of which has a useful and in fact an essential place when applied to the right type of case.

* Read in opening a discussion in the Section of Surgery at the Annual Meeting of the British Medical Association, Aberdeen, 1939.

Palliative Treatment

Small internal piles associated with a generalized congestion of the lower rectum are not infrequently seen, particularly in stout subjects who may be liable to hepatic congestion and constipation. Medical treatment directed specially towards limiting the intake of meat and alcohol is of great value; the free drinking of water and barley-water will help constipation and enable aperients to be drastically cut down. I am anxious to stress the common tendency among the general public to take aperients unwisely and to excess: drugs such as salts, calomel, phenolphthalein, and some herbal teas undoubtedly produce irritating watery stools, and when these are voided there is a considerable amount of straining and rectal congestion. Women are very liable to these troubles, which often start during pregnancy.

Palliative treatment is necessary for an attack of prolapse with thrombosis. The more the local sepsis, with ulceration or sloughing, the greater the need for palliative treatment. I am strongly opposed to surgical treatment at this stage, and although this is a matter upon which some controversy exists I shall hope to convince you that surgery in the stage of septic thrombosed internal piles is unnecessary; it is apt to result in delayed healing and chronic post-operative ulceration, which may take months or years to clear up.

The most essential item in palliation is for the patient to be in bed: the foot of the bed may be raised upon blocks, or the buttocks be slightly elevated on a pillow. An attempt should be made to reduce the prolapsed pile or piles, and in most cases this can be done with gentle but firm pressure with the forefinger, aided by some dry gauze or cotton-wool. A firm pad of gauze moistened in lead lotion is applied and kept in position by a T-bandage. If the piles cannot be reduced, as sometimes happens when there is a massive degree of thrombosis with much external oedema, the anus is shaved, the patient confined to bed, and compresses of 1 in 2,000 perchloride, eusol, lead lotion, or dettol lotion are applied every three or four hours. This treatment can be carried out effectively in the patient's home; and in the hospital class I have been struck by the excellent results obtained provided the patient realizes the importance of staying in bed. On these lines the thrombosis quickly resolves, ulceration heals, and external oedema disappears; in fact, after a week's efficiently applied palliative treatment the majority of patients are able to begin getting up and about again. Only in the worst cases, with smelly sloughing piles, is longer treatment up to two or three weeks required in bed. A perianal abscess forms occasionally, and for such surgical drainage must be given.

When thrombosis has resolved, digital and proctoscopic examination will show if further treatment is required. In many cases nothing further has to be done. For some, injections are advisable, and in a few it is clear that large prolapsing piles remain and a ligature operation is necessary for cure: in no case should this be done within six weeks after an attack of prolapse and thrombosis.

Injection Treatment

The objects of injection treatment are twofold: (1) to produce a chemical thrombosis in the internal haemorrhoidal plexus, and, following this, (2) a sclerosis in the submucous layer which will fix loose redundant mucous membrane to the inner muscle layer and by fibrosis draw up the pile so that it no longer prolapses or becomes grasped by the sphincters.

The solution which is used at all the out-patient clinics at St. Mark's Hospital is 5 per cent. phenol in almond oil, with 2 grains of menthol to 1 oz. (Albright's solution). This is injected into the submucous layer by a technique which is now so well known as not to need detailed description. The following points might be mentioned.

SELECTION OF CASES

The ideal cases for injection are first-degree piles: a single small vascular pile can almost invariably be cured by a single injection. Second-degree piles are less favourable, for although bleeding can usually be stopped and prolapse improved so that the patient becomes symptom-free, recurrence after a variable time is not infrequent. Third-degree piles, particularly those which show an overgrowth of squamous epithelium at the lower pole and those with a marked external skin tag in relation (inter-external piles), are not suitable for injection, and as a general rule a ligature operation should be performed in these cases. Associated conditions such as anal fissure, a fibrous polyp, proctitis, or any inflammatory condition such as abscess or fistula are contraindications to injection treatment.

AGE AND OCCUPATION

Patients are commonly seen with piles of the second degree verging upon the third degree, and the choice of treatment may reasonably be influenced by age and occupation. In subjects under 30 years of age it is impossible to guarantee permanent cure by any method, including operation, and for this reason injection treatment is the method of choice in young subjects, with the object of postponing the need for operation until middle age. Late second-degree piles have a better chance of responding well to injections when the patient is engaged in some professional or sedentary occupation. Men doing hard manual work are often better treated by operation with a view to permanent cure rather than be subjected to disturbance of their work by repeated attendances for injections which in the end may prove disappointing.

TECHNIQUE

It is essential to use a sufficiently long proctoscope and inject the phenol in oil into the submucous layer at the upper pole of the internal haemorrhoids. No pain should be caused by the injection, and if a patient complains of pain during or soon after injection it is certain that either the puncture has been too low down or a complicating fissure is present. There is no set routine for distributing the injections, but the constant arrangement of the three primary piles (left lateral, right anterior, right posterior) must be recognized. If one internal pile is more prominent than the others, particularly the right anterior, I find it is best to inject this alone at the first session and deal with the remainder at subsequent sessions. If one pile is seen to be bleeding, this should be chosen for the first injection.

The amount injected is usually 2 or 3 c.cm., and sufficient should be used to produce a smooth rounded swelling in the submucous layer, slightly pale but not white. In the case of a pile with a good deal of mucosal laxity it is often advisable to inject rather more, say 3 c.cm. into the upper end, and a further 2 c.cm. through another puncture slightly lower down.

When a pile is particularly vascular and bleeds freely from its lower pole, an injection of a stronger solution of phenol is valuable: for instance, 6 to 8 minims of 20 per cent. phenol in glycerin and water can be injected into the substance of the pile, using Graeme Anderson's syringe, either at the same time as the submucous injection

high up (double-injection method) or at a later session. Subsequent injections are given at weekly intervals, and when a firm submucous induration has been produced in the three primary internal haemorrhoidal areas (as judged by digital examination and by direct touch with a probe through a proctoscope) the treatments should be discontinued. Re-examination is reasonably made after an interval of four to eight weeks, and then if the condition remains satisfactory the patient may be discharged. He should be advised to report at any time in the future if a recurrence of bleeding or prolapse should be noted.

RESULTS

The results of injection treatment, when done efficiently with proper doses at the correct level, depend upon the stage reached by the haemorrhoids at the beginning of treatment. For first-degree piles there is a high percentage of cures; in the second degree the results are less favourable, there being fairly commonly a recurrence of prolapse within three to five years; in the third degree there is at the best some amelioration in prolapse for perhaps three to twelve months, but any sort of permanent cure is unlikely to be achieved. A single third-degree pile, particularly when situated in the right anterior quadrant, sometimes does remarkably well, and by a series of repeat injections every year or so I have been able to keep many such patients very comfortable over a course of years.

Statistics in regard to the proportion of cures in the three degrees of haemorrhoids are very difficult to arrive at, and I believe my colleague, Mr. E. T. C. Milligan, has some interesting figures to present in this connexion. To mention some mass statistics, Kilbourne (1934) reviewed more than 25,000 collected cases, and estimated that recurrence took place in at least 15 per cent. within three years: it is not unfair to assume that this 15 per cent. was probably constituted by third-degree piles which were tentatively treated by injection.

To conclude this section I should like to summarize these few important points:

1. Cases for injection must be carefully selected, chiefly with the object of excluding third-degree piles.
2. Cases must be uncomplicated by other ano-rectal conditions.
3. The technique of injection is not easy, and much care must be taken to give adequate doses of the phenol-in-oil solution at the correct level—that is, at the upper pole of the pile and slightly above the level of the ano-rectal ring.
4. Complications are due to errors in technique; pain is due to the injection being placed too low down; chemical necrosis is due to a too superficial injection, overdosage, or injection into sclerosed tissue. Abscess formation is extremely rare, and is due to a combination of the above faults.

Operative Treatment

Operative treatment is the method of choice for the permanent cure of internal piles of the third degree, particularly those of the intero-external variety—that is, a prolapsing pile with a related skin tag in the same quadrant. Operation is indicated also when piles are complicated by a fibrous polyp, a fissure, a fistula, or any other rectal lesion which also requires surgical treatment for cure.

CHOICE OF ANAESTHETIC

In my opinion the ideal anaesthetic is a local infiltration of the external sphincter with 1 per cent. procaine by Hirschman's technique, supplemented by a subcutaneous injection of the solution at the anal margin in relation to the three primary piles. This form of anaesthesia gives

perfect relaxation and is very rarely followed by any urinary difficulty; for instance, I have operated on a man aged 86 and a woman of 87 by this method; neither patient had to be catheterized after operation and both made a perfect recovery.

The next best form of anaesthesia is a low spinal injection, which also gives a perfect anaesthesia for the operation, but is followed in 5 to 10 per cent. of cases by a troublesome headache, and retention of urine is more common. General anaesthesia is seldom used, and its avoidance has contributed largely to post-operative comfort, for straining and vomiting have been practically eliminated by the use of local or spinal anaesthesia.

CHOICE OF OPERATION

It would appear that the operation for cure of haemorrhoids has now become more standardized, and I submit that Whitehead's operation, the clamp and cautery method, excision and suture over a clamp, and many others should all be relegated to the past. Even Salmon's ligature operation, which was practised by Salmon, Allingham, and their successors at St. Mark's Hospital for nearly a century, has now been superseded.

In 1919 Ernest Miles described his standard method of haemorrhoidectomy: a V-shaped flap of skin at the anal margin was raised at the base of each primary pile, the base of the V being at the muco-cutaneous junction, and a stout silk ligature was tied round the internal pile and the tissue thus dissected up. The anatomical difference between this operation and Salmon's operation—in which, it may be recalled, the scissor cut is made at the muco-cutaneous junction—remained unexplained until Milligan and Morgan's (1934) classical paper on the "Surgical Anatomy of the Anal Canal," which shows clearly that when this wide incision is made the longitudinal muscle is included in the ligature. This has now formed the most important feature of their later paper (Milligan, Morgan, Jones, and Officer, 1937), and a precise and ideal operative technique is now available. I speak with much enthusiasm regarding this operation, which enables the three primary piles to be removed completely; the ligatures are tied round the pedicles of the piles and are fixed by the inclusion of the longitudinal muscle, so that they cannot slip upwards and expose raw surfaces in the anal canal. The best ligature material is a matter of opinion: I prefer No. 3 chromic catgut on a round-bodied needle, tied by transfixion, with the knot on the lumen. The related portions of the external venous plexus are removed, together with the overlying skin, and post-operative tags are noticeably absent after this operation. Other benefits conferred by this operation are quicker healing, greatly reduced and almost absent risk of post-operative stricture, and total abolition of secondary haemorrhage, due, I presume, to the better fixation of the ligatures.

RESULTS OF OPERATION

When the operation is correctly performed for prolapsing haemorrhoids in a patient of middle age or over a permanent cure can be confidently promised in all cases. If the operation has to be done for young subjects under 30 the outlook is not so consistently good; I agree with Miles that some reserve must be exercised in promising cure in these cases, seeing that the primary piles may not all be fully developed at the time of operation. Apart from this, recurrences after operation are only rarely seen, and are due to faulty technique—that is, inadequate exposure of the piles and ligation of their lower extremities only, or omission to secure all three primary piles.

Conclusion

I should like to take this opportunity of stressing the safety of a haemorrhoid operation when done under modern conditions and the certainty of cure of a distressing and disabling complaint. The common notion among the public that haemorrhoidectomy is a very painful operation is one that should be strenuously opposed. The modern operation under local anaesthesia, with a carefully considered and well-ried after-treatment, is far from being a serious ordeal: the patient is able to walk to a bath on the third post-operative day, and the anal wounds are usually almost healed in two weeks after operation.

Summary

Essential points in the preliminary examination of a case of haemorrhoids are detailed: how to recognize the difference between internal and external haemorrhoids, the three degrees of internal haemorrhoids, and other associated anal conditions.

The treatment of external piles is described.

The three main lines of treatment for internal piles in routine use at St. Mark's are given: palliative, injection, and operative.

The safety of a haemorrhoid operation done under modern conditions, and the certainty of cure, are stressed.

REFERENCES

- Kilbourne, N. J. (1934). *Ann. Surg.*, **99**, 600.
Miles, W. E. (1919). *Surg. Gynec. Obstet.*, **29**, 497.
Milligan, E. T. C., and Morgan, C. N. (1934). *Lancet*, **2**, 1150.
— — — Jones, L. E., and Officer, R. (1937). *Ibid.*, **2**, 1119.

ELECTRICALLY INDUCED CONVULSIONS IN TREATMENT OF MENTAL DISORDERS *

BY

Wm. H. SHEPLEY, M.B., Ch.B., D.P.M.

Senior Assistant Medical Officer, Warlingham Park Hospital

AND

J. S. MCGREGOR, M.B., Ch.B.

Assistant Medical Officer, Warlingham Park Hospital

(WITH SPECIAL PLATE)

It has long been observed that any severe shock, traumatic, pathological, or psychological, may occasionally lead to dramatic improvement in certain cases of mental disorder. Within the field of psychiatry the term "shock therapy" may be taken to mean the purposive induction of a sudden and disturbing physical or mental impact with specific remedial intent, to alleviate a disordered state of mind. To-day the term is used with narrower connotation, and denotes the induction of a state of coma by the administration of insulin (Sakel) or the production of epileptiform convulsions by such drugs as cardiazol, triazol, or picrotoxin. Dr. Lazlo von Meduna, in 1934 at Budapest, was the first to practise convulsion therapy, by using a solution of camphor in oil administered intramuscularly. This was soon superseded by the intravenous injection of an aqueous solution of cardiazol, giving more certain and controlled action (von Meduna, 1934). Meduna adopted the latter method of treatment because it had been observed that when epilepsy supervened in some cases of schizophrenia the schizophrenic symptoms cleared up (Muller,

1930). This has since been recorded by other workers (Shepley, 1938). Von Meduna (1937) suggested hypothetically that a biological antagonism exists between schizophrenia and epilepsy, and although the validity of this hypothesis is still disputed it is agreed by most workers that, despite certain disadvantages, the treatment is justified by results.

A new method of inducing convulsions by means of electric current has recently been introduced to this country, and definite advantages are claimed for it. It originated at Rome, being evolved by the Italian workers Cerletti and Bini in an effort to overcome the disadvantages of the drug method. The introduction of the electric method may appear to require some justification, since it necessitates the use of considerable voltage. The disadvantages of the drug methods are well known, and it is only necessary to summarize them here. In our experience the principal drawback is the extreme fear and apprehension engendered in some patients, a fear which is so great as to induce Neustatter and Freeman (1939) to use general anaesthesia in an effort to overcome it. This fear is also a great disadvantage where it is necessary to repeat the injection in order to obtain the major fit. Again, extreme and distressing psychomotor excitement often lasting some hours is met with; also such sequels as vomiting, confusion, and irritability are common in spite of careful preparation of the patient. Transient loss of memory and even gross intellectual impairment have also been reported (Dynes, 1939). Dislocations of the jaw and fractures of spine and femora have occasionally been noted by some observers (Stalker, 1938; McGuinness, 1939), while cardiac complications such as arrhythmias, auricular fibrillation, and heart-block have been recorded (Dick and McAdam, 1938). Difficulties are also encountered from the thrombosis of veins, as well as from their inaccessibility in obese subjects and those whose veins are of poor calibre. To some extent these difficulties can be avoided by the use of triazol, but this carries certain new disadvantages.

Nor is the induction of convulsions by the drug methods devoid of mortality, though this is certainly of low incidence. The report of the Board of Control for the year 1938 gives the number of deaths from cardiazol as ten: no analysis of these figures, however, is given. Without any suggestion of comparison, but merely as an indication that electrically induced methods are reasonably safe, it can be stated that a considerable number of such convulsions have been induced without fatality in patients treated at the Rome clinic. In evolving the new technique Cerletti applied the experience gained in previous research upon the problems of epilepsy undertaken at Genoa. He had produced epileptic fits in animals, chiefly dogs, by means of the normal electric current (125 volts A.C.). This was applied for a fraction of a second, the electrodes being placed in the mouth and the rectum. The technique and the physio-pathological results were published by his pupil Chiauzzi (1934). Together with his assistant Bini, Cerletti continued his researches in Rome. It was Bini who constructed the simple but practical instrument used to regulate the voltage and the duration of the passage of the current. This when perfected was patented in Italy by Bini, and was described by him as an "apparatus for electric applications." In the clinic at Rome where Cerletti carried out his observations the histological examination of the cerebral cortices of dogs was made after the application of much higher voltages than necessary to induce a fit and following the induction of multiple fits. These examina-

* Received for publication December 1, 1939

- 52
W. B. GABRIEL: HAEMORRHOIDS

FIG. 1.

FIG. 2.

FIGS. 1 and 2.—Prolapsed and sloughing internal haemorrhoids in a man aged 26, showing appearance at first examination and after seven days' expectant treatment at home. Subsequently treated by injections.

FIG. 3.

FIG. 4.

FIGS. 3 and 4.—Prolapsed and thrombosed internal haemorrhoids in a woman aged 25, with much external thrombosis and oedema, before and after seven days' expectant treatment in bed at home. Later treated by injections. [The photographs were taken by Mr. H. J. R. Bussey at St. Mark's Hospital.]

W. H. SHEPLEY AND J. S. MCGREGOR: ELECTRICALLY INDUCED CONVULSIONS IN TREATMENT OF MENTAL DISORDERS

FIG. 1.—Instrument panel.

FIG. 2.—Electrodes.