success in overcoming difficulties at home, at school, in the swimming-pool, or on the sports field. No other form of treatment was employed during the time that the cases were under observation, the use of drugs, even during the asthmatic attacks, being discouraged.

Results

Restricting discussion for the time being to the effect of treatment on the asthmatic attacks, the results are shown in the following table. The classification used is that employed by Boland (1938), who graded cases into five groups according to the frequency, and to some extent the severity, of the attacks, Group I comprising the most severe cases.

Case No.	Group before Treatment	Group after Treatment	Places Promoted
1	IV	V	1
2	III	IV	1
3	II	III	1
4	II	IV	2
5	I	IV	3

It will be seen that in the first three cases described the patients while under treatment improved to the extent of being promoted one grade, in the fourth case two grades, while in the fifth the patient was promoted three grades. It must be remembered that the patients had previously experienced periods of freedom from asthmatic attacks ranging from one to six months. While it is within the bounds of possibility that in the first four cases the improvement observed in the severity and frequency of the attacks was unrelated to the line of treatment that had been adopted but was attributable to spontaneous recovery, in the fifth case improvement was manifested so rapidly and consistently from the very onset of treatment that the possibility of spontaneous recovery could hardly be entertained. Diminution in the frequency and the severity of the attacks was, however, but one aspect of the improvement obtained in these cases. An outstanding feature was the effect upon the patient's general physique as shown by his ability to take up and retain employment or to be in regular attendance at school, and still further by his increased capacity for taking part in physical activities or normal excitements without ill effect. A third characteristic, and perhaps the most noticeable of all, was the alteration which was brought about in the child's personality and in its outlook on life. Children formerly characterized by shyness, diffidence, anxiety, and over-conscientiousness became confident, happy, and care-free. The spontaneous remarks of parents, teachers, and others gave ample evidence of this alteration. A satisfactory response to treatment appeared to depend primarily upon the recognition and adjustment of adverse factors in the child's mental attitude towards his symptoms and in the parents' attitude towards the child's ill-health, and to have no direct relation to the duration of the symptoms, the severity of the attacks, or the degree of incapacity.

Discussion

Two main factors emerge from a study of recent literature on the subject of bronchial asthma. First, the growing tendency to consider psychological factors as being of significance in its aetiology; and, secondly, and arising therefrom, the necessity for employing appropriate measures of treatment for this aspect of the disorder. It would seem from the examples here described that psychological factors may be of importance either in predisposing to or in precipitating asthmatic attacks, and that the introduction of psychological methods of treatment in selected cases, whether this takes the form of simple reassurance, removal of adverse environmental attitudes,

re-education, or resolution of psychoneurotic problems, merits greater attention than it has hitherto received. Particularly would this appear to be so in patients who are still at an age when habits of mind and bodily reflexes are less firmly established.

Summary

Views expressed by contributors to medical literature on the significance of psychogenic factors in cases of bronchial asthma are recorded.

Five cases of asthma in childhood referred for psychological investigation are described.

Conclusions regarding the respective parts played by allergic, infective, and psychogenic factors in the aetiology of asthma in childhood are arrived at.

The methods of treatment adopted and the results obtained in specific cases are discussed.

The desirability of employing psychological methods of treatment in selected cases is stressed.

REFERENCES

- Boland, E. R. (1938). *Guy's Hosp. Rep.*, **88**, 218.
 Bray, G. W. (1937). *Recent Advances in Allergy*, 3rd ed., London.
 Coke, F. (1939). *Asthma*, 2nd ed., Bristol.
 Dejerine, J., and Gauckler, E. (1915). *Psychoneurosis*, translated by Jelliffe, 2nd ed., Philadelphia.
 Gillespie, R. D. (1936). *British Medical Journal*, **1**, 1285.
 Gordon, R. G. (1939). *A Survey of Child Psychiatry*, contributed by Contemporary British Authorities and edited on behalf of the Child Guidance Council, London.
 Hurst, A. F. (1929). *Practitioner*, **123**, 4.
 McGregor, H. G. (1938). *The Emotional Factor in Visceral Disease*, London.
 Pearson, R. S. B. (1938). *Clin. J.*, **61**, 363.
 Rogerson, C. H., Hardcastle, D. H., and Duguid, K. (1935). *Guy's Hosp. Rep.*, **85**, 289.
 Strauss, E. B. (1937). *Ibid.*, **87**, 273.
 Ziegler, L. H., and Elliott, D. C. (1926). *Amer. J. med. Sci.*, **172**, 860.

TREATMENT OF INGUINAL HERNIA IN ADULTS

BY

JOHN F. O'CONNOR, F.R.C.S.I.

Surgeon, Limerick County Hospital

A glance at the above title may, I feel, evoke the response on the part of many surgeons that "surely we now know all that is to be known on this subject." When we explore a little further into the matter and refresh our minds on the results of the treatment of inguinal hernia they are so disappointing as to justify fully an article of this nature. I will not attempt to describe a new operation for the cure of inguinal hernia (such a thing would now be almost impossible), but rather would make a plea for a rational procedure that will appeal to the average surgeon and ensure success. That such a plea is necessary is proved (1) by the large numbers of recurrent hernias that come to hospital; (2) by the available statistics on the subject, some of which are a blot on present-day surgery; (3) by the disinclination of many practitioners to advise operative treatment for their private patients; and (4) by the existence of numerous prosperous firms which daily advertise their wares for the "cure" of hernia of all types by means of a truss.

Operative Results in the Literature

A short survey of the results of operations by different men discloses very different results. Fallis (1936), reporting on 1,600 cases, stated that of direct hernias 11.6 and of indirect 7.4 per cent. recurred. Max Page (1934) gave the recurrence rate as 20.2 per cent. in indirect and 25.8 per cent. in direct inguinal hernia. The latter figures were compiled from a list of healthy policemen, on whom one should expect the best results. Grace and Johnson (1937) of New York reported results

of herniotomy in 1,032 patients over 50 years of age as 25 per cent. recurrence in primary and 34 per cent. in those treated for recurrent hernia. These figures were almost all for the Bassini type of operation. Keynes's (1937) figures showed no recurrence in ninety large indirect cases and only three in ninety-four direct hernias. Burdick and Higinbotham (1935) for recurrent inguinal and sliding or large scrotal hernias adopted the drastic procedure of cutting across the spermatic cord (in cases where the other testis was normal) and leaving the testis in the scrotum. They had six deaths in 200 cases, and of 169 cases followed up twenty recurred. In four cases the testis sloughed and 17 cases suppurated.

The Injection Method

Injection treatment, like many other new ideas, found a number of supporters, especially in the United States and to some extent in Europe. The idea appeals to the public, particularly to those who have had a hernia for many years and who now believe that a minor miracle can be performed for their benefit. It also appeals to the practitioner who cannot do major surgery. During a recent visit to the United States I found little enthusiasm for this method among surgeons of repute whom I met. Injection treatment, if successful, would be of considerable value for old people and for those on whom any major operation is contraindicated or refused. Unfortunately these are usually people who have large hernias with large rings, and their cases are the most difficult to treat by injection. Irreducible hernias and also those which cannot be effectively controlled by a truss are unsuitable for such treatment. The immediate aim of injection is to close the internal ring by injecting a sclerosing solution around its edges and then into the processus vaginalis in the inguinal canal. Numerous solutions have been described for the purpose. The majority contain phenol as the principal ingredient, or urea hydrochloride. Ten to twenty injections are required at intervals of four or five days, and a truss must be worn *all the time* between injections and for several weeks afterwards. Porritt (1939) reported that out of 246 cases there were four failures and twelve which required further injections after apparent cure. These results were excellent and show a high degree of skill in this technique, but to recommend it as a routine practice in less skilled hands is not justified in our present state of knowledge. Porritt himself considers it useful for selected cases. Great care must be taken not to enter the peritoneal cavity or disaster may follow. Berne (1938) of Los Angeles reports two deaths due to peritonitis caused by intraperitoneal injection. Burdick and Coley (1937) of New York reported that forty-seven out of fifty-six cases relapsed after treatment. The method is entirely a blind procedure, and while it may be safe in the hands of some surgeons it cannot be recommended for general acceptance. Operative treatment under local anaesthesia properly carried out is to my mind preferable to the prolonged discomfort of injections with their attendant risks and failures.

Apropos of the treatment of inguinal hernia in adults by means of a truss, little may be said except to condemn it in general terms. There are of course a few cases in which no other treatment can be carried out—persons whose hernial contents have forfeited all right to domicile in the abdominal cavity or who are very bad risks for any operation, or those on whom a necessarily incomplete operation has been performed, such as for strangulation. For the average man, especially labourers, the truss is a dangerous apparatus, and even for the

more sophisticated and leisured classes its use should be discouraged; rather let us improve our individual operative technique and instil confidence into the public regarding an operation which should occupy the premier place in the domain of surgery for safety and success.

Operative Repair

Progress in surgery must be maintained by the development of new ideas and new experiments, but the majority of operations described for the treatment of inguinal hernia during recent years have little to recommend them. Some are perhaps ingenious in conception or reflect in some way the genius of the author; others are an imitation or mutilation of an original; some are of academic interest; but the majority are of little practical importance.

The name Bassini will always retain its place in the history of surgery of hernia, and to him our debt of gratitude is acknowledged. His work, however, was the foundation-stone upon which Halsted built the edifice. The latter's technique was further improved by Andrews. I omit through lack of space references to other operations which I consider are mainly of theoretical interest, except that of Gillies. Apropos of the latter operation, I consider that it is often performed unnecessarily. It takes a considerable time and is apt to tear as much as repair in the hands of the average surgeon. In my own list of cases I undertook this operation only twice, as an experiment—unnecessary on each occasion; I admit, however, that it is useful in large hernias when properly done. Andrews's method should be successful in 95 to 100 per cent. of all adults on whom this operation can be undertaken.

The following is a list of non-urgent operations for inguinal hernia which I have performed during the ten-year period 1928–38 and examined since January 1, 1940.

Total number of operations performed	103
Number of patients examined after operation	84
Number who had left home (migratory labour)	15
Number who died from other diseases since discharge	4
Number of recurrences	nil

The youngest patient was 18 years of age and the oldest 85. Eleven operations were for recurrent hernia—patients previously operated on by other surgeons. It will be observed that of eighty-four cases examined none had recurred during an interval of from two to ten years. Almost all the patients were agricultural labourers or farmers whose work involved heavy strain, and therefore constituted a fairly severe test for the operation. The anaesthetic used in most cases was gas and ether, and local anaesthetic for those for whom a "general" was not considered suitable.

The technique carried out in all cases except two was similar to that of Edmund Andrews, but I left the cord outside both leaves of the external oblique aponeurosis. The following are the steps of the operation:

The canal is fully opened and the sac is dealt with in the usual manner. The transversalis fascia should be clearly defined if possible from the pubic spine to the internal ring, exposing the deep epigastric vessels. Hold the cord forwards and sew the transversalis fascia to the inner or shelving margin of Poupart's ligament with interrupted sutures of No. 1 chromic catgut. Suture in a similar way the edge of the upper leaf of the external oblique aponeurosis to Poupart's ligament in front of the preceding layer, then suture the edge of the lower leaf of the external oblique aponeurosis to the front of the upper leaf with interrupted sutures of No. 2 chromic catgut. These three layers are behind the cord and constitute a strong line of defence where it is required. Great care is needed in making the new aqueduct a close fit for the cord, particularly at the internal abdominal ring. The word

"canal" is a misnomer: the old canal is completely obliterated, and instead a new exit is provided which is almost straight through the abdominal wall, composed of three layers of white fascia, each fitting closely against the walls of the emerging cord.

In some patients it may not be possible, especially in direct hernia, to find anything but a remnant of the transversalis fascia, and in these cases one should mobilize and turn down a flap from the anterior rectus sheath and suture it into Poupart's ligament instead of the transversalis fascia. In small hernias it is unnecessary to get an overlap of the external abdominal aponeurosis; it is sufficient in these cases to suture a good "bite" of the lower and upper leaves together by interrupted sutures. Gentleness and the avoidance of all unnecessary trauma are strongly recommended, while the use of a gauze wire as a dissecting agent on the cord is deprecated.

As regards suture material I personally favour chromic catgut. The use of wire in any form is not merely unnecessary but may cause serious trouble. First, it is apt to tear the tissues or cut into them, and, secondly, it acts as a permanent foreign body that may promote sepsis. Kangaroo tendon has the former disadvantage, while silk sutures appear to have the second. I am not aware of any real evidence that catgut is not the best material available.

Causes of Failure

As to the cause of so many failures in the treatment of inguinal hernia by operation, I submit that they are mainly due to the following: (1) the performance of the Bassini operation; (2) insufficient and improper suturing at the internal abdominal ring; (3) tearing the edges of Poupart's ligament by means of thick hard catgut or tight ligatures; (4) omitting to warn the patient against any strain or hard work for three months after operation; (5) failure of appreciation on the part of many surgeons of the fact that the operation requires proper planning and careful execution in every detail. It is not to be regarded as an operation "anyone can do with his eyes almost closed," unless he is to "view" his results with his eyes completely closed.

Summary

A series of eighty-four cases of inguinal hernia treated by a simple operation, with 100 per cent. success, is published.

No distinction has been made between direct and indirect hernia, as in a few cases notes were incomplete.

No patient admitted under my care during a ten-year period with inguinal hernia was denied operation.

Several patients were admitted to hospital for treatment for other conditions and were found to be suffering from hernia, for which all agreed to have an operation.

The number of patients fitted with a truss by dispensary doctors in recent years appears to be *nil*. In hospital practice this meant that many of the patients in my list had small hernias, and an overlap of the external oblique aponeurosis was not feasible or practicable at operation.

[Since writing the above I have read the article by Mr. W. Gray in the *British Medical Journal* of April 6, 1940, but I have no reason to alter my views on the subject. Perhaps Irish fascia is tougher than that found elsewhere!]

BIBLIOGRAPHY

- Andrews, Ed. *Operative Surgery*. Bickham, W. S., 7, 343, Philadelphia.
Berne, C. J. (1938). *J. Amer. med. Ass.*, **110**, 1812.
Burdick, C. G., and Coley, B. L. (1937). *Ann. Surg.*, **106**, 322.
— and Higinbotham, N. L. (1935). *Ibid.*, **102**, 863.
Fallis, L. S. (1936). *Ibid.*, **104**, 403.
Grace, R. V., and Johnson, V. S. (1937). *Ibid.*, **106**, 347.
Gray, W. (1940). *British Medical Journal*, **1**, 568.
Keynes, G. (1937). *Proc. roy. Soc. Med.*, **30**, 529.
Maingot, R. (1937). *Postgraduate Surgery*, vol. 3, London.
Page, C. M. (1934). *British Medical Journal*, **2**, 896.
Porritt, A. E. (1939). *Proc. roy. Soc. Med.*, **32**, 893.

SCARLET FEVER IN A PUBLIC SCHOOL

BY

J. K. BOSTOCK, M.B.

An epidemic due to the haemolytic streptococcus occurred recently in a public school. Almost every case was swabbed and typed according to the Griffith method, and it was revealed that out of eighty-seven cases of this infection sixty-four were due to Type 18 (Griffith). The remaining twenty-three cases were composed of nine other different Griffith types.

Details of the Epidemic

Scarlatiniform rashes occurred in five of the sixty-four cases of Type 18 (a proportion of about 1 to 13) and in six of the twenty-three cases composed of various types (a proportion of about 1 in 4). Type 18 was the only one which showed epidemic qualities. It was of fairly low virulence and evinced no tendency to increase in virulence by passage. Complications were few and varied. A proportion of the patients were merely suffering from a nasal discharge, and did not complain of feeling ill.

The school has eight houses—six in a central block and two in separate buildings. The infection was at first confined to one house almost exclusively, eventually spreading slowly through the school. In this particular house twenty-three out of the twenty-four cases were Type 18. In another house eleven cases were all of Type 18. In the six other houses there was a preponderance of Type 18, but other types were also present. One of these houses had sixteen cases of Type 18 and five other cases each of a different type.

The details of the eighty-seven cases are set out below. With two exceptions—a case of streptococcal pneumonia and one of erysipelas—they were all nasopharyngeal infections.

TABLE I.—Incidence of Scarlatinal Rash among Various Types of Haemolytic Streptococcus

No. of Cases	Type (Griffith)	No. with Rash
3	1	—
2	2	—
1	3	1 case
1	5	1 case
4	12	—
3	13	1 case
1	17	—
64	18	5 cases
1	20	—
1	28	—
5	Undetermined. 3 sterile swabs; 1 not swabbed	3 cases
1	Throat swab negative; close contact with positive case	(Erysipelas)

TABLE II.—Analysis of Individual Types, showing Occurrence of Complications, including Rash

No. of Cases	Type (Griffith)	Complications	No.
3	1	—	—
2	2	Otitis media	1
1	3	Sc. rash only	1
1	5	Sc. rash only	1
4*	12	—	—
3	13	Otitis media	1
1	17	Sc. rash only	1
64	18	Effusion into joints	2
		Pneumonia with empyema	1
		Haematuria ? due to sulphanilamide, with sc. rash	1
		Albuminuria with sc. rash	1
		Sc. rash only	3
1	20	—	—
1	28	—	—
5	?	Otitis media	2
		Sc. rash only	3
1	?	Erysipelas	1

* A typical case of scarlet fever due to this type occurred in the previous term.