I am indebted to Messrs. Ciba, Ltd., and N.V. Organon for supplies of testosterone propionate issued under the trade names of perandren and neo-hombreol respectively.

REFERENCES

- Callow, R. K. (1938). *Proc. roy. Soc. Med.*, 31, 841.
 Hamilton, J. B. (1937). *Endocrinology*, 21, 744.
 Huhner, M. (1939). *Med. Rec.*, 149, 366.
 Miller, N. E., Hubert, G., and Hamilton, J. B. (1938). *Proc. Soc. exp. Biol., N.Y.*, 38, 538.
 Parkes, A. S. (1937). *Lancet*, 2, 902.

MARCH FRACTURE

REPORT ON CASE INVOLVING BOTH FEET

BY

R. DRUMMOND, M.R.C.S., L.R.C.P.

Assistant Lecturer in Pathology, Welsh National School of Medicine

(WITH SPECIAL PLATE)

March fracture (*pied forcé*, or Deutschlander's disease) is a condition in which fracture of the shaft or neck of the second or third metatarsal bones occurs, rarely of the fourth, without any history of trauma. Practically all the cases are unilateral, and fracture of more than one metatarsal in the same foot is exceptional. Cases in which both feet are involved are even rarer, and hence the reason for publishing this case. Further, the present case illustrates the salient features of the condition. This fracture is comparatively rare, and it is not infrequently missed. Few cases are seen in surgical or orthopaedic clinics. Thus in the Cardiff Royal Infirmary fracture clinic only two or three cases are met with annually (B. Thomas, in a personal communication). Its importance, especially at the present time, is considerable, in that it occurs in people whose occupation requires them to be about on their feet for long periods, and hence it is not infrequently seen in nurses, soldiers, and factory workers. It is characteristic of the condition that there is never any history of trauma to the affected foot, and in this respect only does it differ from the ordinary run of fractures.

Characteristics of March Fracture

Initially there is a complaint of dull aching pain in the affected foot at the end of a day's work. This pain becomes progressive, and sooner or later has to be alleviated by resting the affected foot; still later, even rest may not afford complete relief. After an interval of time varying from a few days to a few weeks oedema occurs on the dorsum of the affected foot, usually over the second or third metatarsal. This oedema is apt to be persistent. The affected metatarsal may be tender to pressure before the fracture occurs, and it becomes more so after fracture; crepitus is not constantly present. Radiography may reveal that in the early stages the fracture is not yet present, but that there is a little rarefaction of the shaft or neck of the affected metatarsal. A later radiograph will show a crack fracture with little displacement. At a still later stage a large spindle-shaped mass of callus will be seen surrounding the line of fracture. Finally, in the more advanced cases there may be only a characteristic spindle-shaped mass of callus to be seen, due to the line of fracture having disappeared owing to bony union having been effected.

In some cases several radiographs may be necessary before the fracture is detected, and in this respect Watson Jones (1940) points out the analogy to carpal scaphoid fracture, in which a number of radiographs may be required. It is, however, the appearance of a dense spindle-shaped mass of callus or bone on the shaft of the metatarsal, without a line of fracture, that has given rise to some confusion as to the precise nature of the condition

or to the exact sequence of events. Thus Mercer (1936) contends that initially an osteoma forms and that subsequently a fracture occurs through it. Yet osteomata are notoriously hard structures, and no other parallel condition arises. However, Mercer (personal communication) considers it possible that a fracture may occur first and callus form around it or that callus formation may precede fracture. The latter seems unlikely, and in the case reported below the sequence of events (that is, fracture followed by callus formation) is well illustrated.

Various theories are held as to causation, but none are definitely proven. Thus spasm of muscles interfering with blood and lymph supply, local myositis, focal sepsis, loss of muscle tone from debility, etc., are all cited. According to Watson Jones (1940), however, metatarsus atavicus—a congenital shortening of the first metatarsal—is frequently a predisposing cause of spontaneous fracture of the second metatarsal bone, and, indeed, may lead to recurrent fracture in the same bone after a variable period.

Certain facts stand out. The subjects are usually young and healthy, and they have spent long hours on their feet. There is no history of trauma. The strain of long standing or marching appears to be the precipitating factor. The onset is gradual, the fracture being preceded by pain, swelling, and tenderness. Moreover, pain may persist for some time after the fracture has united.

Case Report

A probationer nurse aged 23 complained of pain (which interfered with her sleep) and swelling of the dorsum of the right foot in December, 1938. These symptoms were present for two weeks before she reported for examination. Three months previously she had had a swelling of the dorsum of the left foot with pain, but she did not report this. There was no history of trauma. Her health was otherwise good and she had had no previous illnesses of significance.

An examination revealed oedema on the dorsum of the right foot over the metatarsal bones and tenderness to pressure. A radiograph disclosed a transverse "crack" fracture of the middle of the shaft of the second right metatarsal bone, without callus formation. In the left foot, which was affected three months previously, there was a spindle-shaped mass of callus surrounding a uniting fracture of the shaft of the third metatarsal bone (Plate, Fig. 1). Elastoplast strapping was applied to the feet, and the patient was kept in bed for three weeks and then sent home to rest. She subsequently returned to work as a nurse and has experienced no further trouble.

Ten months later a further radiograph (Fig. 2) of the feet was taken; it showed, on the left side, disappearance of the line of fracture, considerable diminution of the callus, and thickening of the cortex of the shaft of the bone. On the right side the line of fracture had disappeared, and there was a dense fusiform mass of ossified callus, whereas in the previous radiograph there was no callus.

Discussion

Though most cases occur in young adults, yet in ten cases reviewed by Meyerding and Pollock (1938) the average age was 40 years, the oldest patient being 61 and the two youngest being nurses aged 19 and 21. Of this series only three patients were males, one of whom was a doctor aged 34. Savés (1936) reported a case of bilateral fracture of the second metatarsal bones in a young soldier. A case of march fracture of the second and third metatarsals of the left foot was reported by Monteith (1934): there was fracture without callus of the third metatarsal, and an old fracture of the second metatarsal with good callus formation.

In regard to these cases generally, the severity of pain varies from a mild to severe persistent dull ache. The occurrence of oedema of the dorsum of the foot is constant, and is associated with pain and tenderness to pressure over the affected metatarsal. The interest of the

case reported here is that the second fracture occurred two months after the first, and the sequence of events (that is, fracture followed by callus formation) is well illustrated.

The condition does not appear to be so generally well known as it ought to be. It has been confused with osteoma, syphilitic periostitis, and sarcoma, and mistaken diagnosis has had unfortunate sequelae. Thus Watson Jones (1940) cites a case in which a wife instituted divorce proceedings, as she was told the lesion was a syphilitic periostitis. Straus (1932) reports a case in which the entire second metatarsal was excised because of an erroneous diagnosis of sarcoma. Dodd (1933) reports the case of a woman who had a classical Lisfranc amputation of the right foot performed for a march fracture of the second metatarsal mistakenly diagnosed as a sarcoma.

Such errors should not occur if due attention is paid to the various x-ray appearances which may be seen, especially that of a spindle-shaped mass of callus or bone on the shaft of the metatarsal, without a line of fracture. The clinical history with absence of trauma is characteristic. Treatment consists in the application of a walking plaster for a month, followed by strapping and foot exercises. The prognosis is good.

My thanks are due to Dr. W. Davies of Sully Hospital, Glamorgan, for permission to publish this case, and to Professor Duguid for help and criticism.

REFERENCES

- Dodd, H. (1933). *Brit. J. Surg.*, **21**, 131.
Jones, R. Watson (1940). *Fractures, and Other Bone and Joint Injuries*, Edinburgh.
Mercer, W. (1936). *Orthopaedic Surgery*, London.
Meyerding, H. W., and Pollock, G. A. (1938). *Surg. Gynec. Obstet.*, **67**, 234.
Monteith, W. B. R. (1934). *Brit. J. Surg.*, **21**, 708.
Saves, M. (1936). *Rev. Orthopéd.*, **23**, 529.
Straus, F. H. (1932). *Surg. Gynec. Obstet.*, **54**, 581.

REMOVAL OF A TERATOMATOUS "TWIN"

BY

VERNON PENNELL, M.A., M.B., F.R.C.S.

Surgeon to Addenbrooke's Hospital, Cambridge.

AND

L. C. MARTIN, M.D., M.R.C.P.

Research Assistant, Department of Medicine, University of Cambridge

(WITH SPECIAL PLATE)

The presence of teratomata in and about the human body is not uncommon, but those displaying elements from the three embryonic layers and developing well-formed organs are distinctly rare. The following case presented several features of interest.

Case Record

The patient was a female aged 5 weeks, the fourth child. (The previous child was said to have been anencephalic.) She was admitted to Addenbrooke's Hospital with a large mass, 6 by 4 by 3½ inches, over the lower lumbar and sacral regions (Fig. 1 on Plate). It presented two dimples, into one of which a probe could be passed for two and a half inches. There was a nipple-like process half an inch from this opening. The child was otherwise well formed and the movements of her lower limbs were normal. There did not appear to be any impulse in the tumour when the baby cried or strained. A radiograph showed a well-marked plaque of bone not unlike a foetal lower jaw or the upper end of the ulna. It also revealed a spina bifida of the lower lumbar region.

The diagnosis of teratoma was made and it was decided to remove it under local anaesthesia. This was carried out by one of us (V. P.). Sixty c.cm. of 0.5% novocain were injected round the base of the tumour. A rather unnecessarily large flap of skin was fashioned and undercut before the tumour was defined, after which a layer of fat which surrounded the

mass was incised and the separation of the twin begun. A cavity was opened and a somewhat disconcerting display of intestines was revealed. A short examination, however, reassured us that they belonged to the teratoma and not to the host. Towards the midline of the patient a meningeal sac was opened and the lumbar spinal cord was clearly seen well above the cauda equina.

Soon after the removal of the tumour it was noticed that the patient's legs had become flaccid and were very largely paralysed. As this rapidly passed off and by next day the child was moving the legs in a perfectly normal manner it would appear probable that the local anaesthetic had trickled into the theca after that membrane had been opened and had become, in fact, a spinal anaesthetic. The child stood the operation well, although somewhat shocked from the loss of so large a proportion of its own bulk. In view of the shock no time was wasted in trimming the rather large skin-flap, and the wound was sewn up as quickly as possible.

The next day the redundant skin was seen to be bulging, and a pair of sinus forceps introduced through the scar evacuated a large quantity of cerebrospinal fluid. For ten days leakage continued unabated. In spite of this, and because of the unremitting care and attention which she received from the sister and the nursing staff, the child made steady progress, except for a small ulcer in the thigh which followed an intramuscular injection of sulphapyridine (given as a prophylactic during the leakage of cerebrospinal fluid). This, however, healed in due course and the child was discharged normal in every way.

Characteristics of the Specimen

The following is the report (by L. C. M.) of the examination of the specimen:

External.—The specimen is covered externally by an oval area of skin 5½ by 3 inches. There is a blind dimple situated 2 inches from one end of the oval of skin, and at 4 inches is an excrescence. Half an inch from this is an opening, flanked by two small protuberances, ? mandibular processes. The opening communicates via an apparent oesophagus with the stomach C (see diagrammatic key to Fig. 2 on Plate). Lying beneath the oval skin is a layer of fat varying in thickness from 1½ to 3 inches.

Internal (Fig. 2).—Lying in the mass of fatty tissue is a smooth cavity approximately 2 by 1½ inches lined by either cartilage or bone, which is partially shown at A. There is a definite firm overhanging ridge B. The cavity contains a primitive stomach (C), communicating with the surface of the tumour at the dimple X by a tube T—presumably the oesophagus. (Probe in X passing into stomach.) The stomach is connected to coils of large intestine (E), about 7 inches in total length, which terminate in the blind caecum D. The large bowel is covered by fatty deposits and shows definite taenia. No trace of liver, spleen, or genital organs can be found by simple dissection, although two small buds distal to the stomach may possibly represent rudiments of liver and pancreas. In the regions P and Q there are apparent extensions of the osteo-cartilaginous roof of the cavity A; but elaborate dissection would have been necessary to display their form. So far as can be seen all three embryonic layers are represented, and the assumption is that this is a partially formed twin.

Diagrammatic key to Fig. 2
on Plate.

Commentary

The following points of interest are raised by this case:

First, the advisability, or the reverse, of operating at so early a date. Here was a child of five weeks

R. DRUMMOND: MARCH FRACTURE INVOLVING BOTH FEET

FIG. 1.—Radiograph of feet taken on December 23, 1938. There is an ununited "crack" fracture of the mid-shaft of the right second metatarsal bone without callus formation. Near the neck of the left third metatarsal bone is a uniting fracture surrounded by a partially ossified spindle-shaped mass of callus.

FIG. 2.—Radiograph taken on October 30, 1939. The line of fracture has disappeared on both sides, and complete bony union has been effected. There is well-marked residual thickening of cortex of left third metatarsal. A spindle-shaped mass of dense bone surrounds mid-shaft of right second metatarsal: it is this appearance which may be confused with new growth or periostitis.