case reported here is that the second fracture occurred two months after the first, and the sequence of events (that is, fracture followed by callus formation) is well illustrated.

The condition does not appear to be so generally well known as it ought to be. It has been confused with osteoma, syphilitic periostitis, and sarcoma, and mistaken diagnosis has had unfortunate sequelae. Thus Watson Jones (1940) cites a case in which a wife instituted divorce proceedings, as she was told the lesion was a syphilitic periostitis. Straus (1932) reports a case in which the entire second metatarsal was excised because of an erroneous diagnosis of sarcoma. Dodd (1933) reports the case of a woman who had a classical Lisfranc amputation of the right foot performed for a march fracture of the second metatarsal mistakenly diagnosed as a sarcoma.

Such errors should not occur if due attention is paid to the various x-ray appearances which may be seen, especially that of a spindle-shaped mass of callus or bone on the shaft of the metatarsal, without a line of fracture. The clinical history with absence of trauma is characteristic. Treatment consists in the application of a walking plaster for a month, followed by strapping and foot exercises. The prognosis is good.

My thanks are due to Dr. W. Davies of Sully Hospital, Glamorgan, for permission to publish this case, and to Professor Duguid for help and criticism.

REFERENCES

- Dodd, H. (1933). *Brit. J. Surg.*, **21**, 131.
Jones, R. Watson (1940). *Fractures, and Other Bone and Joint Injuries*, Edinburgh.
Mercer, W. (1936). *Orthopaedic Surgery*, London.
Meyerding, H. W., and Pollock, G. A. (1938). *Surg. Gynec. Obstet.*, **67**, 234.
Monteith, W. B. R. (1934). *Brit. J. Surg.*, **21**, 708.
Saves, M. (1936). *Rev. Orthopéd.*, **23**, 529.
Straus, F. H. (1932). *Surg. Gynec. Obstet.*, **54**, 581.

REMOVAL OF A TERATOMATOUS "TWIN"

BY

VERNON PENNELL, M.A., M.B., F.R.C.S.

Surgeon to Addenbrooke's Hospital, Cambridge.

AND

L. C. MARTIN, M.D., M.R.C.P.

Research Assistant, Department of Medicine, University of Cambridge

(WITH SPECIAL PLATE)

The presence of teratomata in and about the human body is not uncommon, but those displaying elements from the three embryonic layers and developing well-formed organs are distinctly rare. The following case presented several features of interest.

Case Record

The patient was a female aged 5 weeks, the fourth child. (The previous child was said to have been anencephalic.) She was admitted to Addenbrooke's Hospital with a large mass, 6 by 4 by 3½ inches, over the lower lumbar and sacral regions (Fig. 1 on Plate). It presented two dimples, into one of which a probe could be passed for two and a half inches. There was a nipple-like process half an inch from this opening. The child was otherwise well formed and the movements of her lower limbs were normal. There did not appear to be any impulse in the tumour when the baby cried or strained. A radiograph showed a well-marked plaque of bone not unlike a foetal lower jaw or the upper end of the ulna. It also revealed a spina bifida of the lower lumbar region.

The diagnosis of teratoma was made and it was decided to remove it under local anaesthesia. This was carried out by one of us (V. P.). Sixty c.cm. of 0.5% novocain were injected round the base of the tumour. A rather unnecessarily large flap of skin was fashioned and undercut before the tumour was defined, after which a layer of fat which surrounded the

mass was incised and the separation of the twin begun. A cavity was opened and a somewhat disconcerting display of intestines was revealed. A short examination, however, reassured us that they belonged to the teratoma and not to the host. Towards the midline of the patient a meningeal sac was opened and the lumbar spinal cord was clearly seen well above the cauda equina.

Soon after the removal of the tumour it was noticed that the patient's legs had become flaccid and were very largely paralysed. As this rapidly passed off and by next day the child was moving the legs in a perfectly normal manner it would appear probable that the local anaesthetic had trickled into the theca after that membrane had been opened and had become, in fact, a spinal anaesthetic. The child stood the operation well, although somewhat shocked from the loss of so large a proportion of its own bulk. In view of the shock no time was wasted in trimming the rather large skin-flap, and the wound was sewn up as quickly as possible.

The next day the redundant skin was seen to be bulging, and a pair of sinus forceps introduced through the scar evacuated a large quantity of cerebrospinal fluid. For ten days leakage continued unabated. In spite of this, and because of the unremitting care and attention which she received from the sister and the nursing staff, the child made steady progress, except for a small ulcer in the thigh which followed an intramuscular injection of sulphapyridine (given as a prophylactic during the leakage of cerebrospinal fluid). This, however, healed in due course and the child was discharged normal in every way.

Characteristics of the Specimen

The following is the report (by L. C. M.) of the examination of the specimen:

External.—The specimen is covered externally by an oval area of skin 5½ by 3 inches. There is a blind dimple situated 2 inches from one end of the oval of skin, and at 4 inches is an excrescence. Half an inch from this is an opening, flanked by two small protuberances, ? mandibular processes. The opening communicates via an apparent oesophagus with the stomach C (see diagrammatic key to Fig. 2 on Plate). Lying beneath the oval skin is a layer of fat varying in thickness from 1½ to 3 inches.

Internal (Fig. 2).—Lying in the mass of fatty tissue is a smooth cavity approximately 2 by 1½ inches lined by either cartilage or bone, which is partially shown at A. There is a definite firm overhanging ridge B. The cavity contains a primitive stomach (C), communicating with the surface of the tumour at the dimple X by a tube T—presumably the oesophagus. (Probe in X passing into stomach.) The stomach is connected to coils of large intestine (E), about 7 inches in total length, which terminate in the blind caecum D. The large bowel is covered by fatty deposits and shows definite taenia. No trace of liver, spleen, or genital organs can be found by simple dissection, although two small buds distal to the stomach may possibly represent rudiments of liver and pancreas. In the regions P and Q there are apparent extensions of the osteo-cartilaginous roof of the cavity A; but elaborate dissection would have been necessary to display their form. So far as can be seen all three embryonic layers are represented, and the assumption is that this is a partially formed twin.

Diagrammatic key to Fig. 2 on Plate.

Commentary

The following points of interest are raised by this case:

First, the advisability, or the reverse, of operating at so early a date. Here was a child of five weeks

with a tumour mass, which it was proposed to remove, three-quarters of the size of its head, and shock was likely to be, and indeed proved to be, considerable. On the other hand, if the child was allowed to grow, so, too, would its attached twin, and a proportionately more extensive operation would be called for, probably necessitating general anaesthesia. It was perhaps remarkable that so large a mass did not result in any difficulty at birth; but the mother was young and the child her fourth. Forceps were not used.

Secondly, the leakage of cerebrospinal fluid.—Although every attempt was made to close the spinal theca, a prolonged leakage of cerebrospinal fluid occurred. The fact that this did not become infected was primarily due to the very skilful and sympathetic nursing that the child received. It is, however, possible that the redundant skin flap, at first thought to be a disadvantage, may have partially contributed to this lack of infection by carrying the suture line well over to the left and nearly two and a half inches away from the midline.

Thirdly, the recognition of the spinal anaesthetic effect arising from local anaesthesia after the theca was opened and disappearing within a few hours.

BENIGN CYST OF THE PAROTID GLAND

BY

M. G. KINI, M.C., M.B., M.Ch.(Orth.), F.R.C.S.Ed.
Surgeon, King George Hospital, Vizagapatam, S. India

(WITH SPECIAL PLATE)

Parotid tumours are common, but simple cysts of the parotid gland are rare. Very few cases of the latter are reported in the literature, and fewer still of cysts communicating with Stensen's duct. Out of 17 cases of tumour of the parotid in my series 10 were benign mixed tumours and 5 malignant tumours, one of which had undergone cystic degeneration. Only 1 among the 17 cases was a simple cyst communicating with Stensen's duct.

The origin of simple cysts in the parotid gland is speculative. Dermoid cysts and cysts of branchial origin may occur, but very few cases are reported in connexion with the parotid gland. In a cyst of branchial origin the existence of lymphoid tissue underneath the stratified epithelium is diagnostic. Cysts may occur in connexion with benign adenomas, and the pathology of these adenomatous structures may be the same as that described by Lenthal Cheate (1931) in regard to breast tumours. In rare cases fairly large cysts in the parotid gland have been described by Hertzler (1937). With these the histological picture is always of an adenomatous structure with a cystic formation due to association with adenomatous neoplasia or to degenerative change leading to cystic formation in an adenoma or blocking of a main duct. Vitamin A deficiency has also an important bearing upon the nutrition of the epithelium of the lining membrane of the salivary gland. In experimental animals Wolbach and Howe (1925) have shown keratinizing metaplasia in mucous membrane, gland duct, and secreting epithelium in vitamin A deficiency. In children Wilson and Dubois (1923) have demonstrated lesions of xerosis and keratomalacia and also metaplasia of the lining membrane of trachea, bronchi, pancreatic ducts, uterus, and salivary glands. In cases of vitamin A deficiency a metaplasia of the epithelium of the duct to the squamous type, with retention

of secretion inside the cavity, may occur. In these cases such changes are general and not localized to one gland. This factor has to be borne in mind in India, as vitamin A deficiency is common in young and old. There is no suggestion, however, of such a deficiency in the following case, in which a cyst communicated with Stensen's duct.

Case Record

A Hindu male aged 60 was admitted with a swelling of two months' duration in the region of the parotid gland and extending into the left cheek. It began as a small swelling in the region of the parotid. He found that the swelling increased in size with the taking of food, but on applying pressure over it the saliva flowed into the mouth and the swelling was reduced to its original dimensions. He was treated by indigenous methods consisting of the application of irritants inside the mouth. Subsequently the pain and swelling increased. He could not without pain empty the swelling by pressure as before. On admission to hospital he was found to be healthy except for the swelling, which on examination was seen to extend from the front of the ear to half an inch behind the angle of the mouth. It was fixed to the parotid gland. It was warm to the touch and ovoid in shape. Its margins were easily defined. Its surface was smooth, and its consistency firm at the periphery and soft and cystic at the centre, with fluctuation. The skin over the swelling was adherent and tender. There was slight limitation of movements of the temporo-mandibular joint. On opening the mouth the left side of the cheek was dry, and on deep pressure, which caused great pain, a drop of yellowish fluid could be squeezed out. The opening of the left Stensen's duct inside the mouth was retracted. After dilatation lipiodol was injected, and a radiograph showed the lipiodol in the cavity in droplets revealing the communication with Stensen's duct (Plate, Fig. 1). Under general anaesthesia the cyst was removed by a transverse incision parallel with the nerves and blood vessels over the most prominent part of the cyst. The cyst wall, which was thin and adherent to the skin, burst during dissection and was found to contain yellow fluid. At the anterior margin of the masseter the cyst was fixed to the duct by a narrow stalk. This was cut across and the cyst removed. A pedicle ligature was applied round the stump, held by forceps near Stensen's duct, and the wound was sutured in layers. After removal of the cyst a persistent parotid fistula was feared, but the wound healed by first intention without formation of a parotid fistula.

Pathological Report.—Section showed that the cyst wall was lined by stratified epithelium, with fibro-fatty tissues in places. No lymphoid tissue was found underneath. There were gland tubules suggestive of its origin from the parotid gland. Nothing was observed that suggested a branchial origin (Fig. 2).

The patient on inquiry after one year states that he is perfectly fit, with no recurrence of the swelling.

Points of Interest

A case of simple cyst communicating with Stensen's duct is reported.

The mode of origin of this cyst is difficult to appreciate, but it is presumed that, being a simple cyst communicating with Stensen's duct, and having some glandular structure round the periphery of the duct without any evidence of lymphoid tissue underneath the stratified epithelium, it began as a cyst in an adenoma arising near Stensen's duct from the glandular structure at the socia parotidis communicating with the duct.

My thanks are due to Dr. T. Bhaskara Menon for his pathological report and photomicrograph.

REFERENCES

- Cheate, G. L., and Cutler, M. (1931). *Tumours of the Breast*, London.
Hertzler, A. E. (1937). *Surgical Pathology of Diseases of the Neck*, Philadelphia.
Wilson, J. R., and Dubois, R. O. (1923). *Amer. J. Dis. Child.*, **26**, 431.
Wolbach, S. B., and Howe, P. R. (1925). *J. exp. Med.*, **42**, 753.

DUNCAN C. L. FITZWILLIAMS: A PLEA FOR A MORE LOCAL OPERATION IN REALLY EARLY
CARCINOMA OF BREAST

FIG. 1.—Diagnosis with no symptoms. Tiny carcinoma under right nipple well shown.

FIG. 2.—Photograph of patient taken thirteen years after operation.

VERNON PENNELL AND L. C. MARTIN: REMOVAL OF A TERATOMATOUS "TWIN"

FIG. 1.—A case of teratoma in a female child aged 5 weeks.

FIG. 2.—Contents of the teratoma. (See diagrammatic key in text.)