occasionally delayed. Figs. 3 and 4 show the condition of a knee-joint two years after arthrodesis. Table IV is a summary of results obtained by arthrodesis during the years 1931-5 inclusive.

TABLE IV

Result of Operation	No. of Cases	Quiescence Maintained	Reactivated and Amputated	Deaths	Cause of Death	Interval between Operation and Death
Bony union	49	46	—	3	Renal tuberculosis Tuberculous meningitis Heart disease	1 year 5 months 8 years
Firm fibrous union	5	5	—	—	—	—
Incomplete fibrous union	2	2	—	—	—	—
Non-union	3	—	2	1	Phthisis	13 months

The table shows that bony union was obtained in 83% of 59 cases, which closely agrees with the results published by Henderson and Fortin (1927), who reported bony union in 85.9% of 211 resections. The 5 cases in which firm fibrous ankylosis resulted have remained functionally excellent, but from the operative point of view the result obtained must be regarded as having fallen short of the objective. The cases of incomplete fibrous union and of non-union, and those which subsequently required amputation, were chiefly due to the common cause of operating in the face of smouldering disease—a mistake not always easy to avoid.

PRIMARY AMPUTATION

Primary amputation was performed in ten cases of this series in an attempt to save the patient's life or on account of the diseased joint being beyond hope of recovery. Three patients died within a few days of operation. Three more survived operation and left hospital, but have died at varying intervals since. The remaining four patients are well.

Amputation is a last resort, but the tendency is to delay it too long. A patient suffering from profound toxæmia has often passed the stage when amputation could have saved his life. Conversely, it is by no means uncommon for widespread disease to exist with remarkably little constitutional disturbance, and in such cases amputation should not be performed until prolonged conservative treatment has failed to yield any hope of recovery of the diseased joint.

Summary

Some general aspects of tuberculosis of the knee-joint in adults are reviewed, with special reference to 108 cases treated at St. Luke's Hospital.

The association of the local lesion with other tuberculous foci, active and quiescent, is shown in 37.75% of cases.

Reference is made to the early presence of bone lesions and to the high proportion of cases so affected.

Symptomatology, x-ray appearances, and diagnosis are discussed. Mention is made of the difficulties of early diagnosis of disease in some cases.

An outline of some of the principles of local treatment is given. Results of conservative treatment in a small series of cases are stated. The results in 59 cases of arthrodesis are quoted, and the opinion is expressed that this treatment is the most suitable for the majority of adult cases, following conservative treatment.

Indications for primary amputation are mentioned. Ten cases are recorded.

My thanks are due to Dr. H. J. A. Colvin, medical superintendent of St. Luke's Hospital, for giving me the benefit of his advice and wide experience. I also have to thank the Medical Officer of Health to the London County Council for his kind permission to publish this article.

BIBLIOGRAPHY

- Boyd, William (1933). *Surgical Pathology*, Philadelphia.
 Brav, E. A. (1933). *J. Bone Jt. Surg.*, **15**, 631.
 Ghormley, R. K., Kirklin, B. R., and Brav, E. A. (1933). *Amer. J. Roentgen.*, **33**, 747.
 Girdlestone, G. R. (1932). *Brit. J. Surg.*, **19**, 488.
 Henderson, M. S. (1936). *Austral. New Zeal. J. Surg.*, **6**, 27.
 — and Fortin, H. J. (1927). *J. Bone Jt. Surg.*, **9**, 700.
 Illingworth, C. F. W., and Dick, B. M. (1935). *Surgical Pathology*, London.
 Tregubov, S. (1937). *J. Bone Jt. Surg.*, **19**, 734.

TESTOSTERONE PROPIONATE IN
FUNCTIONAL IMPOTENCE

BY

A. W. SPENCE, M.D., F.R.C.P.

Assistant Physician, St. Bartholomew's Hospital

All observers are agreed on the successful results obtained with testosterone propionate in the treatment of impotence in castrated or hypogonadal patients. In view of this a trial was made of its effect in six patients with impotence unassociated with any obvious endocrine disorder and probably due to psychological causes. As the following case reports show, negative or unsatisfactory results were obtained in every case.

Case Reports

Case 1.—The patient, a single man aged 48, had been impotent for six years. He associated his trouble with frequent masturbation as a youth. At the age of 18 he began to have frequent involuntary nocturnal emissions, with dreams of a sexual type. During the course of years these became more pronounced, and later he had a constant feeling of exhaustion, weakness, and discomfort in his testes, which affected his work and happiness. There was no change in hair growth or in the size of his genitalia. He consulted Dr. Henry Wilson, who made the diagnosis of psychogenic impotence. Psychological treatment was unsuccessful. On examination he was seen to be a somewhat undersized nervous man, with a hesitant speech. The contour of his body had a slightly feminine tendency. His hair growth, penis, scrotum, and testes were normal. His trunk and legs were affected with psoriasis. The remainder of the physical examination revealed nothing abnormal. He was treated with 100 mg. of testosterone propionate intramuscularly three times a week for two weeks without any change in his condition.

Case 2.—The patient was an unmarried man aged 25. Ten years ago the left testis was knocked; it became very painful and later gradually diminished in size. Three years ago erections became less frequent and less powerful, and during the last six months three attempts at intercourse were unsuccessful because of inability to obtain an erection. On examination his appearance and growth of hair were normal, his voice was broken, the penis and right testis were normal, but the left testis was as small as a pea. The remainder of the physical examination was normal. He was treated at first with intramuscular injections of olive oil as a control measure, without effect. He was then given 100 mg. of testosterone propionate intramuscularly three times a week for two weeks. No improvement in his erections resulted.

Case 3.—A widower aged 50, in whom the first signs of impotence appeared between 1920 and 1925. He was married in 1914 and for several years intercourse was normal, but since 1925 this was impossible owing to his inability to obtain an erection. His wife died two years ago, and he would now like to marry again, but hesitates because of his condition. He was seen in the department of psychological medicine by Dr. E. B. Strauss, who diagnosed psychogenic impotence. Treatment by hypnotic suggestion brought about no improvement. Physical examination revealed nothing abnormal. As a control measure he was given intramuscular injections of olive oil for two weeks, during which he had frequent erections at night, but an attempt at coitus was unsuccessful. Treatment with 50 mg. of testosterone propionate intramuscularly twice a week for four weeks and, subsequently, with 100 mg. twice

a week for two weeks did not improve his condition. Unlike patients with hypogonadism he experienced no improvement in his general health, and his depression persisted.

Case 4.—This patient, aged 29, wished to become engaged to be married, but hesitated because of impotence. Occasionally he masturbated, but during the last seven years four attempts at intercourse were unsuccessful owing to his being unable to obtain an erection. At the age of 16 he had mumps, but there was no evidence of orchitis. Treatment for two weeks with 100 mg. of testosterone propionate three times a week was quite unsuccessful.

Case 5.—A married man aged 35. For the last six years he had been gaining in weight and was impotent. During the last few months he lost sexual desire. He was plump, but could not be considered obese. Examination revealed that the growth of hair, penis, and scrotum were normal, but the testes were slightly smaller than normal. The remainder of the examination disclosed nothing abnormal. He was first treated with subcutaneous injections of distilled water twice a week for a month. After the second injection he said that he felt more alert, and after ten days he had coitus successfully on one occasion, an act which he had been unable to accomplish during the previous five or six years. This improvement, however, was not maintained. He was then given 50 mg. of testosterone propionate intramuscularly three times a week for three weeks. During the second night after the first injection coitus was successful, but after that he was quite impotent. Subsequent treatment with gonadotrophic extract of the urine of pregnancy for two months was also unsuccessful.

Case 6.—This patient, a married man aged 43, complained of obesity and impotence. Twenty-two years ago, while receiving a low-rationed diet as a prisoner of war in Germany, he began to put on weight, and has gradually increased in weight ever since. Ten years ago he began to fall asleep at his work as a lorry-driver, and six years ago he had at first premature ejaculation on intercourse, and, later, erections failed, although they would occur on a tram or bus. During the war of 1914-18 he had gonorrhoea and syphilis, both of which were successfully treated. On examination he was a cheerful red-faced man, weighing 19 st. 6½ lb.; his obesity was generalized. His voice, distribution of hair, penis, and testes were normal. His lungs showed evidence of chronic bronchitis; his cardiovascular and nervous systems were normal. There were numerous small varicose veins on his legs. X-ray examination showed a small pituitary fossa. His Wassermann reaction was negative; the blood count was normal; the basal metabolic rate on two occasions was +16% and +8%. No definite glandular disturbance could be found as a cause of his condition. He was treated with a reducing diet by Dr. George Graham, and lost 3 st. in three and a half months. His impotence was at first treated with intramuscular injections of olive oil twice a week for two weeks. As this had no effect he was given intramuscular injections of 100 mg. of testosterone propionate three times a week for two weeks, but no improvement was obtained, and attempts at intercourse were still unsuccessful. He was then given one dose of 200 mg. of testosterone propionate. During the second and fourth nights after this he had a full erection and intercourse was successful. After a second dose of 200 mg. a week later no response was obtained.

Discussion

Psychological and endocrine disturbances are probably two of the most important causes of impotence, but whereas a disturbance of the psychological factor will always cause impotence, this is not always so in patients in whom testicular hormone is apparently lacking, for it is well known that occasionally some eunuchs are potent. Callow (1938) has shown that eunuchs excrete in their urine an amount of androgen which is within normal limits, and Parkes (1937) and Callow (1938) have suggested that the adrenals may be an additional source of supply. It is possible that in some eunuchs this is enough to maintain potency. In patients whose impotence is due solely to deficient testicular function treatment with testosterone

propionate will establish normal potency. The histories of the patients reported above demonstrate that impotence which is not due to an endocrine cause but rather is psychogenic in origin is unaffected by doses of testosterone propionate which had produced a considerable response in eunuchs. So certain is the response to testosterone propionate in impotence due to deficient testicular function that it may be asserted that if such a response is not obtained in a patient with underdeveloped genitalia his impotence is due also to other causes. The following case report illustrates this.

Case 7.—An unmarried Indian aged 24 complained of underdeveloped genitalia and inability to obtain an erection. He had never had full erections; on attempting intercourse he would have a slight erection and ejaculate before penetration. His voice did not break, and he shaved about twice a week. For many years he suffered from nervous dyspepsia; and he was unable to micturate when standing up. On examination the configuration of his body was normal; there was only a scanty growth of hair on the face, the axillary hair was normal, there was no hair on the trunk, and the pubic hair was shaved. His voice was high-pitched. The penis and testes were small, and the prostate was just palpable. He was treated with 100 mg. of testosterone propionate intramuscularly twice a week for two months. Two days after the first injection he began to have nocturnal emissions which occurred about twice a week, whereas previously these did not occur. Treatment caused slight increase in growth of the penis, but had no effect on his impotence or on his general condition.

There are, however, reports in the literature that are not in agreement with these findings. Thus, Hamilton (1937) brought about potency, with doses of 20 mg. of testosterone propionate three times a week, in a man whose impotence was not due to an organic cause and in whom a course of injections of non-medicated oil as a control measure had had no effect. Miller, Hubert, and Hamilton (1938) have reported that one patient with psychogenic impotence was restored to "a low normal sex drive and erectile capacity," and that another, who before treatment "escaped from the environmental situation partly responsible," was restored to complete normality. This patient, however, remained normal while receiving injections of an inert oil. The response obtained in these two patients appeared to be less than that in two eunuchs similarly treated. The most striking results with testosterone propionate in impotence not due to endocrine deficiency are those reported by Huhner (1939), who produced full potency in all the seven patients treated, with doses ranging from 5 mg. to 35 mg. three times a week for one to six months.

From my own observations it may be concluded that an adequate secretion of male sex hormone is necessary to ensure erection on psychogenic stimulation; if, however, the psychogenic stimulus be absent or disturbed excess of the hormone does not compensate for this deficiency. On the other hand, patients with impotence due to hypogonadism seem to be unduly sensitive to the influence of testosterone propionate, for whereas large doses have little or no effect on the erections of normal men, in eunuchs they may cause long-sustained priapism.

Summary

Six patients with functional impotence unassociated with any evidence of testicular dysfunction failed to respond to testosterone propionate given intramuscularly in doses varying from 50 to 100 mg. three times a week for two to six weeks. A seventh patient with hypogonadism also failed to respond after treatment for two months, probably because of a super-added psychological disturbance. As these doses are sufficient to cause prolonged erections in eunuchs, it is concluded that testosterone propionate has no effect on impotence due to psychological disturbances.

I am indebted to Messrs. Ciba, Ltd., and N.V. Organon for supplies of testosterone propionate issued under the trade names of perandren and neo-hombreol respectively.

REFERENCES

- Callow, R. K. (1938). *Proc. roy. Soc. Med.*, 31, 841.
 Hamilton, J. B. (1937). *Endocrinology*, 21, 744.
 Huhner, M. (1939). *Med. Rec.*, 149, 366.
 Miller, N. E., Hubert, G., and Hamilton, J. B. (1938). *Proc. Soc. exp. Biol., N.Y.*, 38, 538.
 Parkes, A. S. (1937). *Lancet*, 2, 902.

MARCH FRACTURE

REPORT ON CASE INVOLVING BOTH FEET

BY

R. DRUMMOND, M.R.C.S., L.R.C.P.

Assistant Lecturer in Pathology, Welsh National School of Medicine

(WITH SPECIAL PLATE)

March fracture (*pied forcé*, or Deutschlander's disease) is a condition in which fracture of the shaft or neck of the second or third metatarsal bones occurs, rarely of the fourth, without any history of trauma. Practically all the cases are unilateral, and fracture of more than one metatarsal in the same foot is exceptional. Cases in which both feet are involved are even rarer, and hence the reason for publishing this case. Further, the present case illustrates the salient features of the condition. This fracture is comparatively rare, and it is not infrequently missed. Few cases are seen in surgical or orthopaedic clinics. Thus in the Cardiff Royal Infirmary fracture clinic only two or three cases are met with annually (B. Thomas, in a personal communication). Its importance, especially at the present time, is considerable, in that it occurs in people whose occupation requires them to be about on their feet for long periods, and hence it is not infrequently seen in nurses, soldiers, and factory workers. It is characteristic of the condition that there is never any history of trauma to the affected foot, and in this respect only does it differ from the ordinary run of fractures.

Characteristics of March Fracture

Initially there is a complaint of dull aching pain in the affected foot at the end of a day's work. This pain becomes progressive, and sooner or later has to be alleviated by resting the affected foot; still later, even rest may not afford complete relief. After an interval of time varying from a few days to a few weeks oedema occurs on the dorsum of the affected foot, usually over the second or third metatarsal. This oedema is apt to be persistent. The affected metatarsal may be tender to pressure before the fracture occurs, and it becomes more so after fracture; crepitus is not constantly present. Radiography may reveal that in the early stages the fracture is not yet present, but that there is a little rarefaction of the shaft or neck of the affected metatarsal. A later radiograph will show a crack fracture with little displacement. At a still later stage a large spindle-shaped mass of callus will be seen surrounding the line of fracture. Finally, in the more advanced cases there may be only a characteristic spindle-shaped mass of callus to be seen, due to the line of fracture having disappeared owing to bony union having been effected.

In some cases several radiographs may be necessary before the fracture is detected, and in this respect Watson Jones (1940) points out the analogy to carpal scaphoid fracture, in which a number of radiographs may be required. It is, however, the appearance of a dense spindle-shaped mass of callus or bone on the shaft of the metatarsal, without a line of fracture, that has given rise to some confusion as to the precise nature of the condition

or to the exact sequence of events. Thus Mercer (1936) contends that initially an osteoma forms and that subsequently a fracture occurs through it. Yet osteomata are notoriously hard structures, and no other parallel condition arises. However, Mercer (personal communication) considers it possible that a fracture may occur first and callus form around it or that callus formation may precede fracture. The latter seems unlikely, and in the case reported below the sequence of events (that is, fracture followed by callus formation) is well illustrated.

Various theories are held as to causation, but none are definitely proven. Thus spasm of muscles interfering with blood and lymph supply, local myositis, focal sepsis, loss of muscle tone from debility, etc., are all cited. According to Watson Jones (1940), however, metatarsus atavicus—a congenital shortening of the first metatarsal—is frequently a predisposing cause of spontaneous fracture of the second metatarsal bone, and, indeed, may lead to recurrent fracture in the same bone after a variable period.

Certain facts stand out. The subjects are usually young and healthy, and they have spent long hours on their feet. There is no history of trauma. The strain of long standing or marching appears to be the precipitating factor. The onset is gradual, the fracture being preceded by pain, swelling, and tenderness. Moreover, pain may persist for some time after the fracture has united.

Case Report

A probationer nurse aged 23 complained of pain (which interfered with her sleep) and swelling of the dorsum of the right foot in December, 1938. These symptoms were present for two weeks before she reported for examination. Three months previously she had had a swelling of the dorsum of the left foot with pain, but she did not report this. There was no history of trauma. Her health was otherwise good and she had had no previous illnesses of significance.

An examination revealed oedema on the dorsum of the right foot over the metatarsal bones and tenderness to pressure. A radiograph disclosed a transverse "crack" fracture of the middle of the shaft of the second right metatarsal bone, without callus formation. In the left foot, which was affected three months previously, there was a spindle-shaped mass of callus surrounding a uniting fracture of the shaft of the third metatarsal bone (Plate, Fig. 1). Elastoplast strapping was applied to the feet, and the patient was kept in bed for three weeks and then sent home to rest. She subsequently returned to work as a nurse and has experienced no further trouble.

Ten months later a further radiograph (Fig. 2) of the feet was taken; it showed, on the left side, disappearance of the line of fracture, considerable diminution of the callus, and thickening of the cortex of the shaft of the bone. On the right side the line of fracture had disappeared, and there was a dense fusiform mass of ossified callus, whereas in the previous radiograph there was no callus.

Discussion

Though most cases occur in young adults, yet in ten cases reviewed by Meyerding and Pollock (1938) the average age was 40 years, the oldest patient being 61 and the two youngest being nurses aged 19 and 21. Of this series only three patients were males, one of whom was a doctor aged 34. Savés (1936) reported a case of bilateral fracture of the second metatarsal bones in a young soldier. A case of march fracture of the second and third metatarsals of the left foot was reported by Monteith (1934): there was fracture without callus of the third metatarsal, and an old fracture of the second metatarsal with good callus formation.

In regard to these cases generally, the severity of pain varies from a mild to severe persistent dull ache. The occurrence of oedema of the dorsum of the foot is constant, and is associated with pain and tenderness to pressure over the affected metatarsal. The interest of the