A CASE OF GLOMUS TUMOUR WITH OBSERVATIONS ON VARIATIONS OF PAIN WITH TEMPERATURE

BY

J. D. N. NABARRO, M.B., B.S.*Late House-physician at University College Hospital*

(WITH SPECIAL PLATE)

The following case of subungual glomus tumour was seen at University College Hospital, and the variations in the pain with changes of temperature were investigated.

Case History

The patient, a single woman aged 38, had suffered from pain in the left ring-finger radiating up the arm to the shoulder for ten to twelve years. The attacks of pain had become more frequent recently, and the patient was seldom free for more than a day. No injury to the finger could be remembered. Repeated examinations for cervical ribs had been made; these proved negative.

The pain started in the nail of the left ring-finger and spread up the front of the forearm and outer side of the arm to the shoulder and occasionally up the side of the neck to the jaw. Squeezing or accidentally knocking the finger gave intense pain of a burning type for several seconds. The pain was provoked by cold and was also worse in winter; it was alleviated by warmth, but heat brought on a throbbing type. The patient also complained of tenderness after experiencing bad pain. This was on the inner side of the middle finger and over the whole of the ring-finger and little finger, the ulnar area of the hand, and up the ulnar side of the forearm to the elbow. These areas were tender to light friction, and movement of the clothing led to soreness.

Examination showed that in the centre of the ring-finger nail there was a small purple patch, 3 mm. long and 2 mm. broad, very sensitive to pressure, surrounding parts being less sensitive. The surface of the nail was quite smooth and showed no distortion. There was no apparent atrophy and no unnatural moisture of the finger: the skin colour was normal. The patient had first noticed the spot twelve years previously, and it had continued unchanged. She stated that her fingers were nearly always cold.

The following were the effects of immersing the left hand in a water bath at various temperatures:

Before Immersion.—No throbbing or pain. Local pressure or tapping the nail produced local pain only, without radiation.

In Water at 14° C. for 60 seconds.—Patient would not tolerate a longer immersion or allow of immersion in water at a lower temperature on account of the severe ache in the ring-finger and little finger and on the ulnar side of the biceps. There was some hyperalgesia to light touch on the ulnar side of the ring-finger.

In Water at 20° C. for Four Minutes.—Slight ache in the ring-finger only; no hyperalgesia.

In Water at 25° C.—Aching quite gone after four minutes.

In Water at 30° C.—Complained of slight fullness in the ring-finger; pressure produced local pain only.

In Water at 35° C.—Still complained of feeling of fullness; pressure gave local pain only.

In Water at 39° C.—Complained of throbbing pain in the nail of the affected finger. Pressure on the nail now caused pain, which radiated up the ulnar side of the forearm to the elbow.

In Water at 45° C.—This gave the same effects.

Reflex vasodilatation was produced by immersing the right hand in water at 47° C. Before immersion the left hand showed no capillary pulsation; there was no pain or abnormal sensation, and pressure produced local pain only. When the right hand had been immersed for sixteen minutes there was capillary pulsation in all the finger-tips of the left hand. Throbbing was felt in the whole hand and was not increased in the affected finger. Pressure on the nail produced pain, which radiated up the finger. A radiograph of the affected

digit showed a small excavation on the dorsum of the terminal phalanx.

Operation was performed by Prof. R. S. Pilcher on May 26, 1939, with a nerve-block anaesthesia round the base of the finger. A small piece of nail was removed (about one-third) and the tumour shelled out quite easily, the pedicle being identified in the process. Microscopical examination revealed a typical glomus tumour (see Plate).

Post-operative progress was as follows: On May 27 the patient complained of two types of pain in the arm. One was a shooting pain in the middle, ring, and little fingers radiating up to the palm only, less severe than the pre-operative pain, but without tenderness when pressing on the finger-tip. There was also a constant burning pain in the upper forearm, the lateral aspect of the arm, and over the pectoralis major. On May 30 there were still the same pains, but less severe. On May 31 occasional shooting pains were felt, also a burning pain in the finger, in the arm, and over the front and back of the chest, accompanied by some hyperalgesia. On June 2 the burning pain had gone: there were occasional shooting pains only. On July 26 the nail had not yet regrown and the finger was slightly tender to the touch. There were now no shooting pains, but a little ache in the finger and arm was experienced after using the arm.

Discussion

The lesion seemed to cause most discomfort at high and low temperatures; this is in keeping with Grant's (1930) observation that the arterio-venous anastomoses from which these tumours are derived and which he studied in rabbits are open only at high and low temperatures. Why high temperatures should produce a throbbing and low temperatures an ache is not easily explained.

The lateness of the diagnosis may be emphasized, the patient's case having been investigated over a period of twelve years. The widespread tenderness of which she complained does not seem to be a common accompaniment of these lesions. The post-operative remnant of pain is in keeping with the two cases of Stabins, Thornton, and Merle Scott (1937), in which pain persisted for three and four months respectively. Vasodilatation of the affected hand has been mentioned by these three authors, and Kolodny (1938) reported a case in which stimulation of the lesion produced sweating of the affected digit. The present case gave no evidence of any such change: there was no unnatural moisture of the digit, and the fingers were cold.

Summary

A case of glomus tumour is reported with widespread tenderness in addition to widely radiating pain. The variations of pain with temperature were investigated.

I have to thank Sir Thomas Lewis, Dr. F. M. R. Walshe, and Prof. R. S. Pilcher for permission to publish this case, and Dr. Freudental for the section.

REFERENCES

- Grant, R. T. (1930). *Heart*, **15**, 281.
Kolodny, A. (1938). *Ann. Surg.*, **107**, 128.
Stabins, S. J., Thornton, J. J., and Merle Scott, W. J. (1937). *J. clin. Invest.*, **16**, 685.

E. Nobel (*Ann. Pediatr.*, 1940, **155**, 160), who records two illustrative cases in a boy aged 4 and a girl aged 12, maintains that enlargement of the spleen is by no means a constant feature in Still's disease. According to him there is no causal connexion between Still's disease and tuberculosis, as suggested by Kienböck. Repeated examinations of the blood during many months show that normal leucocyte counts may alternate with marked leucocytosis. He suggests that a deficiency of vitamin B₁ plays a much greater part in the causation of Still's disease and primary chronic arthritis in children than is generally recognized, as he has found that treatment with the anti-neuritic vitamin is very successful in such diseases, as well as in polyneuritis, gastro-enteritis, neuralgia, and sciatica.

M. G. KINI: BENIGN CYST OF PAROTID GLAND

FIG. 1.—Radiograph showing the lipiodol in the cyst.

FIG. 2.—Photomicrograph of the cyst wall, showing the stratified epithelium.

J. D. N. NABARRO: GLOMUS TUMOUR

The tumour in section.

H. KENRICK CHRISTIE: NEUROFIBROMATOSIS, OR VON RECKLINGHAUSEN'S DISEASE

FIG. 1.

FIG. 2.