Treatment

Treatment has hitherto been purely symptomatic and of little avail. In view of the probability that the disease is due to some disorder of the sympathetic nervous system, it would appear to be justifiable to treat the severe intractable cases by section of the sympathetic chain.

Since 1936 we have treated three cases by lumbar cord ganglionectomy; the fourth patient was not so treated because of a background of thyrotoxicosis. The results in the three operated patients have been excellent: in each case the symptoms have completely disappeared and the patient has remained well. Short notes of the cases submitted to operation are as follows:

Case 1.—A woman aged 24, a machinist, had had hot and burning feet when she was a girl at school. The trouble steadily got worse and the attacks began to come on in the winter. In these attacks, which lasted from three to four weeks and had intervals of only a few days between them, she says the pain was very severe and the feet "felt on fire." She could not walk, and had spent the previous six months lying in bed with the bare feet in the open. She was a thin pale woman, much worn by pain and loss of sleep. Both feet were bright red in colour—not sweating, very tender, with prominent veins; arterial pulsation was present but not exaggerated. A left-sided lumbar cord ganglionectomy was performed in July, 1936. The unilateral operation was done in order that the unoperated limb might serve as a control. The result in the left foot was entirely satisfactory. After watching the case for a year, the operation was done on the right side with equally good results. The patient has been seen since on many occasions, and remains entirely free from her former symptoms.

Case 2.—A man aged 55, a clerk, began to suffer five years ago from a hot patch on the right heel. This quickly spread to the whole foot, and shortly involved the left one. He then complained of burning pain, which varied little in summer or winter. He was quite incapacitated by the pain and tried to obtain sleep by bathing the feet in cold water two or three times during the night. There was no evidence of any disturbance in the blood supply to the feet; all pulses were present, but the skin was red, dry, intensely tender, and felt hot. A bilateral lumbar cord ganglionectomy was done in February, 1940, after which the symptoms at once disappeared; the feet returned to normal and he is perfectly well.

Case 3.—A woman aged 26, a supervisor, complained of burning pain in the feet, made much worse by warmth. The condition was typical of erythromelalgia. She was kept under observation for twelve months, during which time she became worse. A bilateral cord ganglionectomy was done in July, 1940. The pain was at once abolished and the feet became normal. She was seen a few days ago, and remains quite free from her trouble.

Summary

An account of erythromelalgia is given and its diagnosis from painful rubor is discussed.

Treatment by sympathectomy is advised and records of three successful cases are given.

REFERENCES

- Burger, L. (1924). *The Circulatory Disturbances of the Extremities*, Saunders, Philadelphia.
Collier, J. (1898). *Lancet*, 2, 401.
Lewis, Sir T. (1933-4). *Clin. Sci.*, 1, 175.
Mitchell, S. Weir (1878). *Amer. J. med. Sci.*, 76, 17.

J. G. Rushton, R. W. Cragg, and L. K. Stalker (*Arch. intern. Med.*, 1940, 66, 531) give an account of a young woman who suffered from spontaneous attacks of hypoglycaemia. Operation failed to reveal the pancreatic tumour which was expected. The patient died after the operation, and necropsy showed extreme atrophy of the adrenal glands and "extensive inflammatory destruction of the thyroid of the type described by Hashimoto." No tumour of the pancreas was found, and the authors consider the hypoglycaemia to have been due to the adrenal changes.

REVERSED ROTATION OF THE INTESTINE ACCOMPANIED BY ENCAPSULATING PERITONITIS*

REPORT OF A CASE

BY

J. W. GEARY GRANT, F.R.C.S.

Consulting Surgeon, Royal Infirmary, Cardiff

The occurrence in the same patient of the above two conditions, which are decidedly rare and are liable to cause some embarrassment to the operator, appears to be worth recording.

Case Report

On May 5, 1939, assisted by Dr. R. Nelson of Treherbert, I operated on a patient of his, a miner aged 62. The history given was that two days previously he was seized with sudden abdominal pain followed by complete constipation. The abdomen was moderately distended. Nothing was felt per rectum, and the history was negative. Under general anaesthesia the abdomen was opened by a right paramedian incision. All that was at first seen was a large cyst with smooth shining white walls; it resembled an ovarian cyst, and occupied the greater part of the left hypochondrium, epigastrium, and left lumbar region (Fig. 1). The only portion

FIG. 1.—Diagram of condition found at first operation for acute intestinal obstruction.

of the intestines visible was the descending colon, which appeared to come out from under the cyst in the left hypochondrium. The rest of the abdomen being carefully walled off, a small incision was made in the cyst, and out of this a quantity of foul-smelling blood-stained fluid escaped. On enlarging the opening it was found to be occupied by the whole of the intestine from the duodenum to the left colon, which emerged in the left hypochondrium from the sac. The gut was entirely free from the inner wall of the sac, and no difficulty was experienced in cutting away the sac from its attachment to the posterior abdominal wall. There was then found to be a volvulus of the mid-gut loop, which had undergone rotation of about 270 degrees in a clockwise direction. This was undone. The distended caecum showed a gangrenous patch the size of a shilling a Paul's tube was fixed in this opening and the caecum was fixed in the abdominal wound, which was closed.

The abdominal wound healed after some suppuration, but a large opening in the caecum remained from which all the intestinal contents escaped. The man's condition having greatly improved, an attempt was made to close the caecostomy; this was immediately followed by symptoms of obstruction. It was then decided to reopen the abdomen. On August 2, the opening in the caecum being temporarily closed, the abdomen was opened under general anaesthesia. It was found that the caecum and ascending colon came out

* Abstract of a lecture given in the Welsh National School of Medicine.

through a tunnel under the root of the mesentery (Fig. 2). (It will be observed from the diagram that the colon is crossed by the superior mesenteric artery and the duodenum. The caecum has nearly reached its proper position, but has

FIG. 2.—Showing the condition of parts found at second operation. Note colon crossed by duodenum and superior mesenteric vessels. The pre-arterial mesentery has contracted its normal but limited adhesion to the posterior abdominal wall, thus forming a narrow tunnel through which the caecum and right colon pass, and is trapped. The caecum and post-arterial mesentery have not become adherent, but at first sight the intestines appear to be in their normal position—caecum on right, small intestines on left.

taken the wrong route to do so, passing behind the duodenum instead of in front. Neither the pre-arterial nor the post-arterial mesentery has contracted any adhesion to the posterior abdominal wall, and the whole of the mid-gut loop swings free and is liable to torsion, as occurred in this case.) The diagnosis was that of reversed rotation of the mid-gut loop, which from the condition seen at the first operation would appear to have been completely enclosed in a cyst-like structure, everywhere free except at its points of attachment in the neighbourhood of the duodeno-colic isthmus. As obstruction had taken place when an attempt was made to close the caecostomy it was decided to carry out a right hemicolectomy. The ileum was divided five or six inches from the ileo-caecal valve and, the proximal end being closed, was anastomosed to the sigmoid. The colon was divided at its emergence from the tunnel under the mesentery, the proximal end closed, and the caecum and ascending colon removed. The patient made an uninterrupted recovery, and five months later resumed his work as a collier. I am informed that he has been perfectly well since.

Discussion

Whilst non-rotation is not a very rare condition—I have myself reported two cases and have met with one or two others (Grant, 1928, 1930-1)—reversed rotation appears to be decidedly rare. Dott (1923-4) reported one and referred to a second. Donald (1927-8) describes one, and the illustration in his article shows exactly the same condition as that found at the second operation on my case.

With regard to the condition of cyst formation present in this case, a good description is given by Wilmoth and Patel (1931). They call it "la péritonite chronique encapsulante," and one of the illustrations to their article has an appearance similar to that of mine on opening the abdomen, although the contents were different. Their description is that of a membrane-like pigskin, smooth, white, and shining—a very good one. They refer to a case in which the operator, seeing what he thought was a cyst, attempted its total extirpation, and they remark, "On peut deviner ce qu'il en advint."

Most of the cases reported by Wilmoth and Patel, Devine (1932-3), Soupault (1940), and various German surgeons, under the title of Zuckergussdarm, Zuckerguss-leber, and Zuckergussmilz, and by other French authors as

"intestin sucre glacé," describe a membrane more or less closely attached to intestine, liver, spleen, etc., as the case may be. I would like to stress that in the case I report there was no such attachment, the bowel in the sac being perfectly free. As Prof. Lambert Rogers remarked to me, one almost wonders whether the extra-embryonic coelom could possibly have been drawn in, forming a covering to the mid-gut loop.

BIBLIOGRAPHY

- Devine, H. B. (1932-3). *Brit. J. Surg.*, **20**, 204.
Donald, Charles (1927-8). *Ibid.*, **15**, 269.
Dott, Norman M. (1923-4). *Ibid.*, **11**, 251.
Grant, J. W. Geary (1928). *Surg. Gynec. Obstet.*, **46**, 133.
— (1930-1). *Brit. J. Surg.*, **18**, 166.
Hunter, J. I. (1922). *British Medical Journal*, **2**, 800.
Soupault, R. (1940). *J. Chir., Brux.*, **55**, 355.
Wilmoth, P., and Patel, J. (1931). *Ibid.*, **37**, 341.

Clinical Memoranda

March Fracture

As medical officer of an infantry battalion I have had a high incidence of march fracture—namely, five cases in the past nine months. Before joining the Army I had not seen a case and had heard very little about this interesting condition. Textbooks tend to dismiss it in a few lines.

CASE REPORTS

Case 1.—An N.C.O. reported sick on November 5, 1939, complaining of pain in the left foot radiating up to the ankle when walking. On examination there was slight oedema of the dorsum of the foot and marked tenderness over the head of the second metatarsal. Abnormal movement and crepitus were elicited. A diagnosis of ? march fracture was made. X-ray examination showed the condition to be Freiberg's infraction of the head of the second metatarsal. An orthopaedic surgeon advised a metatarsal bar on the shoe. The condition improved considerably, and the man was able to walk about normally but could not proceed more than a mile without a rest. His occupation was changed to that of a clerk, and he was able to perform his full duties. About April 1 he was transferred to another unit and I have not seen him since.

Case 2.—A private reported sick on January 6, 1940, complaining of pain on walking and slight swelling of the dorsum of the right foot. There was extreme tenderness over the second metatarsal. An x-ray examination revealed a transverse crack of the second metatarsal with no displacement. The foot was immobilized in plaster-of-Paris for twenty-one days. The plaster was then removed and the patient instructed to walk about. One week later he was discharged to duty in employment as a clerk. He has had no further trouble with his foot.

Case 3.—This man reported sick on July 22, 1940, with pain in the right foot on marching and swelling of the dorsum of the foot. There was marked tenderness over the second metatarsal. A radiograph taken on July 23 revealed no bony injury or abnormality. His foot was rested for a week with no improvement. A further radiograph taken on August 2 showed a typical march fracture. The foot was immobilized in plaster-of-Paris for twenty-one days. On removal of the plaster the patient was instructed to walk around. He complained of pain on walking for a week or so afterwards and was very loath to put any weight on the foot. A ten-day course of foot exercises by the physical training instructor restored his confidence. He was discharged to duty on September 13 and has not complained since.

Case 4.—This man reported sick on August 20, 1940, with a complaint of pain in the left foot and slight swelling of the dorsum on walking. There was marked tenderness over the third metatarsal. A diagnosis of ? march fracture was made and the foot was radiographed. The x-ray film showed slight thickening of the shaft of the third metatarsal on the medial side but no evidence of a fracture. The foot was rested for ten days and another radiograph was taken. This time the film revealed a typical march fracture with a spindle-shaped