Published in final edited form as:

Neuroscience. 2007 December 5; 150(2): 478–486.

Restraint-induced *fra-2* and *c-fos* expression in the rat forebrain: relationship to stress duration

Marc S. Weinberg, Milena Girotti, and Robert L. Spencer

Department of Psychology, University of Colorado at Boulder, 345 UCB; Muenzinger D244, Boulder, Colorado 80303

Abstract

The protein product of the *fra-2* gene (Fra-2), a *fos*-family member, can compete with Fos protein for participation in AP-1 transcription factor complexes and each protein can contribute different transactivational consequences to an AP-1 complex. To date, there is limited characterization of *fra-2* mRNA expression in the rat forebrain. We examined basal and restraint-induced mRNA expression (*in situ* hybridization) of *fra-2* in the rat forebrain and compared its temporal-spatial pattern to *c-fos*. In contrast to the very low basal expression of *c-fos*, *fra-2* basal expression was moderately high throughout cortex and some subcortical structures, including prominent basal expression in the hypothalamic paraventricular nucleus (PVN). Restraint-induced *fra-2* expression was quantified in the prefrontal cortex (PFC), lateral septum (LS) and PVN. Maximal *fra-2* gene induction in the PFC and LS was delayed (60 min) after restraint onset with respect to *c-fos* (15 min), whereas in the PVN, *fra-2* mRNA increased within 15 min of restraint. Additionally we compared *c-fos* and *fra-2* gene expression in rats given shorter or longer restraint durations, but equal total time from stress onset to sample collection, to determine the extent to which the kinetics of gene induction matched that of a hypothalamic-pituitary-adrenal axis hormone response. Rats given 45 min recovery after 15 min restraint showed less *c-fos* expression in the PVN, less *fra-2* expression in the prelimbic and infralimbic PFC, and no difference in the LS compared with rats restrained for 60 min. Thus, the expression of both genes was sensitive to stressor duration, but this sensitivity varied with brain region. Differential basal and stress-induced expression patterns of the *fra-2* and *c-fos* genes are likely to have important functional consequences for AP-1 transcription factor dependent regulation of neural plasticity.

Keywords

ACTH; CORT; hypothalamus-pituitary-adrenal axis; IEG

Introduction

Inducible immediate-early genes (IEGs), or genes which respond rapidly and transiently to a variety of stimuli (Robertson, 1992), may serve as markers of neuronal activity and thereby aid in our understanding of the neural basis of behavior. Of these IEGs, neuroscientists have paid particular attention to *c-fos* gene expression (Sheng and Greenberg, 1990). Another member of the *fos*-family, *fra-2*, was isolated in 1990 (Nishina et al., 1990) and subsequently characterized in terms of promoter elements and pharmacological expression profile in the

Email: marc.weinberg@colorado.edu, Phone: (303) 492-5853, Fax: (303) 492-2967.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

chicken embryo (Yoshida et al., 1993) and mouse (Foletta et al., 1994). The general structural organization of the *fra-2* gene is very similar to the *c-fos* gene (Nishina et al., 1990, Yoshida et al., 1993, Foletta et al., 1994). There are some indications, however, that *fra-2* gene expression patterns and protein function differ from *c-fos*, and that *fra-2* expression may be especially important for long-term neural and endocrine adaptation to repeated stress (Sabban et al., 2004, Hebert et al., 2005), drugs of abuse (Liu et al., 2005), brain injury (Pennypacker et al., 2000, Butler and Pennypacker, 2004), photoperiod phase shift (Engel et al., 2005), and recovery from sleep deprivation (Terao et al., 2003).

Both physical and psychological stressors produce rapid and transient induction of *c-fos* gene expression in a variety of rodent brain regions (Kovacs, 1998). These expression patterns have been used to assist with delineating stress-responsive neural circuits (Pacak and Palkovits, 2001). Moreover, the induction of *c-fos* is believed to be important for coordinating adaptive cellular changes within these structures in response to stress (Herman and Cullinan, 1997). To date, there is very limited characterization of the basal and stimulus-induced spatial-temporal expression pattern of *fra-2* in the rat brain. Early northern blot studies revealed readily detectable basal levels of *fra-2* in whole mouse brain homogenates, contrasting with the much lower basal expression of other members of the *fos* family (including *c-fos*, *fra-1*, *fra-2*, and *fosB*) (Foletta et al., 1994). *fra-2* mRNA expression, as detected by *in situ* hybridization, has only been described for the hippocampus (Beer et al., 1998), and the suprachiasmatic (SCN) and paraventricular nuclei (PVN) of the hypothalamus (Honkaniemi et al., 1994, Schwartz et al., 2000).

In this study we have used *in situ* hybridization to directly compare the time-course of *fra-2* and *c-fos* gene induction in rat forebrain in response to acute psychological stress (restraint). The only other study to examine stress-induced *fra-2* mRNA in the brain used systemic injection of capsaicin as its stimulus, and the analysis was restricted to the PVN (Honkaniemi et al., 1994). Additionally the neural response to capsaicin, a putative systemic stressor, may differ substantially from that of processive stressors such as restraint, which require higher-order processing (Herman and Cullinan, 1997). For this study we have examined parallel *fra-2* and *c-fos* gene expression at three rostral-caudal levels of rat forebrain that contain key structures believed to be part of a network of stress-reactive forebrain structures - prefrontal cortex, lateral septum and PVN (Herman et al., 2003). Because of potential differential mechanisms of transcriptional control in the *fra-2* and *c-fos* gene, studying their spatial-temporal expression patterns on adjacent brain sections from the same animals may reveal new information regarding neural recruitment patterns of the brain undergoing stress. An additional component of this study is that we directly compared gene expression in rats at the same point in time after restraint onset, but with different durations of acute stress (continuous versus some period of restraint recovery). This comparison can be informative for both intrinsic gene expression kinetics as well as system-level differences in ongoing neural activity as a result of different durations of restraint.

A portion of this manuscript has appeared previously in abstract form (Weinberg et al., 2005).

Experimental Procedures

Animals

Male Sprague–Dawley rats (Harlan Laboratories, Indianapolis, IN, USA) were allowed a two-week acclimation period after arrival to the animal facilities at the University of Colorado before experimental use (10 weeks old, weight range 282–329 g at experimental onset). Rats were housed in pairs in polycarbonate tubs with wood shavings, and were given food (Purina Rat Chow; Ralston Purina, St. Louis, MO, USA) and tap water *ad libitum*. The colony room

lights were regulated on a 12-h light/dark cycle, with lights on at 07:00 h. Procedures for ethical treatment of animals conformed to the guidelines found in the "Guide for the Care and Use of Laboratory Animals," DHHS Publication No. (NIH) 80-23, revised 1996 ed. and were approved by the University of Colorado Institutional Animal Care and Use Committee. All efforts were made to minimize the number of animals used and their suffering.

Grouping and Testing

Rats were divided into six groups designated according to their duration of restraint and post-restraint recovery time, respectively: 0-0, 15-0, 15-45, 60-0, 60-120, and 180-0. Group 0-0 rats were taken from their home cages and killed without any restraint exposure. Groups 15-0, 60-0, and 180-0 were taken from their home cages and restrained for 15, 60, or 180 min and then immediately killed. Rats in groups 15-45 and 60-120 were restrained for 15 or 60 min, replaced in their home cages for 45 or 120 min, after which time they were killed. The 15-45 and 60-120 groups were established for comparison to the 60-0 and 180-0 groups, respectively.

Restraint involved taking rats from the home cage and placing them in adjustable length (15.5 +/- 2.5 cm long and 6.3 cm diameter) Plexiglas tubes with air holes in the front, top and back. This stressor is considered to be primarily psychological in nature because it does not produce pain or direct physical insult (Herman and Cullinan, 1997). Restraint took place in a separate room adjacent to the home cage room. The 180-0 group consisted of four rats. Otherwise, all groups consisted of six rats. Cage-mates were restrained simultaneously.

Upon rapid decapitation, brains were collected, flash-frozen in an isopentane bath maintained between -40°C and -30°C and stored at -80°C. Trunk blood was taken in EDTA coated tubes, kept on wet ice and centrifuged at 5000g for ten minutes within forty-five minutes of sacrifice. Plasma was then rapidly frozen and stored at -80°C until hormone assay.

Corticosterone and ACTH radioimmunoassay

Corticosterone (CORT) and ACTH plasma concentrations were measured by radioimmunoassay, as described previously (Girotti et al., 2006). Assay sensitivity for CORT was approximately 0.5 µg/100 ml for a 20 µl sample. The detection limit for ACTH was approximately 15 pg/ml for a 50 µl sample.

In situ hybridization histochemistry

Ten micrometer brain sections were cut on a cryostat (Leica model 1850) through the extent of the prefrontal cortex (approximately 3.20 mm anterior to bregma (Paxinos G, 1998)), rostral septum (approximately 0.20 mm anterior to bregma) PVN (approximately 1.80 mm posterior to bregma), thaw-mounted onto poly-L-lysine-coated slides and stored at -80°C. *In situ* hybridization for *c-fos* or *fra-2* mRNA was performed as described previously (Girotti et al., 2006). For generation of the *c-fos* probe, plasmids containing a fragment of *c-fos* cDNA (courtesy of Dr. T. Curran, St. Jude Children's Research Hospital, Memphis) were used. For the generation of *fra-2* ribonucleotide probe, a 450 bp fragment (Exon 4) was amplified from GenBank accession number **NM_012954** using the primer set: 5'-gcagaaggagaaggagaag-3' (forward) and 5'-agagtgggggagttcaag-3' (reverse). Primer products were cloned into pCR-II TOPO vectors (Invitrogen) to generate RNA probes. Direct sequencing revealed the correct sequence and orientation of the cloned fragment. [³⁵S]-labeled complementary RNA probes were generated using standard *in vitro* transcription reagents (Promega, Madison, WI, USA). After the hybridization assay procedure, sections were dehydrated and exposed to X-ray film for 1-4 weeks.

Image Analysis

Semi-quantitative analyses of autoradiographs were performed on digitized images from X-ray films (NIH Image) as described (Campeau and Watson, 1997). For the prefrontal cortex, a 25 x 25 pixel square was placed within the prelimbic (PL), infralimbic (IL) or ventral orbital (VO) prefrontal cortex, as determined by using the Paxinos and Watson (Paxinos G, 1998) atlas for guidance. For the lateral septum (LS) and paraventricular nucleus of the hypothalamus (PVN), regions of interest (ROIs) were drawn around the brain structure using the Paxinos and Watson atlas for guidance. The ROIs were first drawn on images with positive signals then saved and positioned on control brain images according to identifiable landmarks. Background areas were chosen in the white matter or in a cell-poor area close to the ROIs. Signal pixels of the brain area of interest were defined as being 3.5 standard deviations above the mean of the set background. The number of pixels and the average pixel values above the set background were then computed and multiplied, giving an integrated densitometric measure of arbitrary units (integrated gray level).

Statistical Analyses

Time course data (groups 0–0, 15–0, 60–0, and 180–0) were analyzed with separate one-way analyses of variance (ANOVA) tests using a Statistical Analysis System (SAS) package for UNIX. When testing group differences in ACTH, CORT, and gene expression (*in situ* hybridization data), *post-hoc* tests (Fisher's least significant difference test) were performed to determine points of significant difference between treatment groups and the results are indicated on the data figures. When comparing groups of rats of different stressor durations but same total time from stimulus onset to death (i.e. 15–45 versus 60–0 and 60–120 versus 180–0) groups were compared by two-tailed independent sample t-tests (Microsoft Excel). In all statistical comparisons, an α level of $P < 0.05$ was used to determine statistical significance. All data were $\log(10)$ transformed prior to statistical analysis due to heterogeneity of variance. Data presented represent group mean $\pm$ S.E.M.

Results

Effect of restraint on HPA axis hormone secretion

One-way ANOVAs indicate a significant main effect of time on plasma ACTH [$F(3,18) = 6.23$, $p < 0.005$], and a significant main effect of time on CORT levels [$F(3,18) = 9.79$, $p < 0.0001$]. *Post-hoc* tests indicate that circulating plasma levels of ACTH and CORT (Fig 1) were significantly higher than control after both 15 and 60 min of restraint. CORT, but not ACTH, was also elevated after 180 min restraint. Rats given a 45 min recovery period (15–45) did not have significantly different plasma levels of ACTH or CORT than rats restrained for the full duration of time (60–0), however rats given 180 min of restraint (180–0) had higher plasma CORT than rats given 60 min of restraint followed by 120 min of recovery (60–120).

Basal c-fos and fra-2 expression

Under no-stress conditions (0–0) there was very little discernable *c-fos* mRNA expression on coronal sections at the level of the prefrontal cortex, lateral septum (with the exception of some signal within the dorsal endopiriform nucleus) and the PVN. In contrast, at each rostral-caudal level examined there was clearly visible *fra-2* mRNA expression throughout cortex with especially prominent expression in the outer layers of neocortex, in the innermost layer of the prelimbic and infralimbic area and layer two of the piriform cortex (Fig 2). At the level of the lateral septum there was also prominent *fra-2* mRNA expression in the shell of the nucleus accumbens and the dorsal endopiriform nucleus. At the level of the PVN there was also pronounced *fra-2* mRNA expression within the parvo and magnocellular regions of the PVN, the supraoptic nucleus and the principal cell body layer of the hippocampus and dentate gyrus.

Relative *c-fos* and *fra-2* mRNA levels were quantified for each restraint condition within the prelimbic, infralimbic and ventral orbital regions of prefrontal cortex, the lateral septum and the PVN.

Effect of restraint on *c-fos* and *fra-2* expression in the prefrontal cortex

One-way ANOVAs of *c-fos* mRNA levels indicate a significant effect of time of restraint on *c-fos* gene expression in the prefrontal cortex: PL [$F(3,17) = 63.98$, $p < 0.0001$], IL [$F(3,17) = 61.51$, $p < 0.0001$], and VO [$F(3,17) = 134.62$, $p < 0.0001$]. *Post-hoc* tests indicate that *c-fos* gene expression in the PL, IL, and VO (Fig 3) was significantly greater in rats restrained for 15 min than in home-cage controls, and remained elevated after 60 min of restraint. Although this *c-fos* gene expression was still significantly higher than basal by 180 min of restraint it had greatly diminished from peak expression levels. Rats given a post-restraint recovery period (15–45 and 60–120) did not have significantly different levels of *c-fos* expression in the prefrontal cortex than rats restrained for the full duration of time (60–0 and 180–0, respectively).

One-way ANOVAs of *fra-2* mRNA levels indicate a significant main effect of time on *fra-2* gene expression in the PL [$F(3,16) = 4.95$, $p < 0.05$], a significant main effect in the IL [$F(3,16) = 3.30$, $p < 0.05$], and a trend in the VO [$F(3,16) = 2.58$, $p = 0.09$]. *Post-hoc* tests indicate that *fra-2* gene expression in the PL, IL and VO (Fig 3) was significantly greater than control expression after 60 min of restraint. This expression declined by 180 min of restraint, however in the PL *fra-2* expression at 180 min of restraint was still significantly greater than control. In both the PL and IL prefrontal cortex, rats restrained for 15 min and given a 45 min recovery period (15–45) expressed significantly less *fra-2* than rats restrained for the full duration (60–0). Figure 4 shows sample autoradiograms of *c-fos* and *fra-2* expression in the prefrontal cortex for rats given 0–0, 15–0, 60–0, or 180–0 min of restraint.

Effect of restraint on *c-fos* and *fra-2* expression in the lateral septum

A one-way ANOVA of *c-fos* mRNA expression indicates a significant effect of time on *c-fos* gene expression in the lateral septum: LS [$F(3,18) = 14.57$, $p < 0.0001$]. *Post-hoc* tests reveal that *c-fos* gene expression in the lateral septum (Fig 5) was significantly greater in rats restrained for 15 min than in home-cage controls, and remained elevated after 60 min of restraint. By 180 min of restraint *c-fos* expression was no longer significantly different than home-cage control expression. Rats given a post-restraint recovery period (15–45 and 60–120) did not have significantly different levels of *c-fos* expression in the lateral septum than rats restrained for the full duration of time (60–0 and 180–0 respectively). The LS showed a highly significant main effect of time on *fra-2* expression: LS [$F(3,18) = 14.23$, $p < 0.0001$]. *Post-hoc* tests reveal that *fra-2* gene expression in the lateral septum (Fig 5) was significantly greater than control expression after 60 min of restraint and remained significantly elevated relative to control at 180 min of restraint. Rats given a post-restraint recovery period (15–45 and 60–120) did not have significantly different levels of *fra-2* expression in the lateral septum than rats restrained for the full duration of time (60–0 and 180–0, respectively). Figure 6 shows sample autoradiograms of *c-fos* and *fra-2* expression in the lateral septum for rats given 0–0, 15–0, 60–0, or 180–0 min of restraint.

Effect of restraint on *c-fos* and *fra-2* expression in the PVN

A one-way ANOVA indicates a significant main effect of time on *c-fos* gene expression in the PVN: [$F(3,17) = 63.03$, $p < 0.0001$]. *Post-hoc* tests show that *c-fos* gene expression in the PVN (Fig 7) was significantly greater in rats restrained for 15 min than in home-cage controls, and remained elevated after 60 min of restraint. By 180 min of restraint *c-fos* expression was significantly reduced from expression at 60 min of restraint. Nonetheless, PVN *c-fos* expression at 180 min of restraint was greater than expression in control rats. Rats restrained for 15 min

and given a 45 min recovery period (15–45) had significantly less *c-fos* expression in the PVN than rats restrained for the full duration of time (60–0). Rats restrained for 60 min and given a 120 min recovery time (60–120) had a trend ($p = 0.07$) for less *c-fos* expression in the PVN than rats restrained for the full duration (180–0). Within the PVN there was also a significant main of time on *fra-2* expression: [$F(3,17) = 5.16, p = 0.010$]. *Post-hoc* tests reveal that *fra-2* gene expression in the PVN (Fig 7) was significantly greater than control expression after 15 min of restraint and remained significantly greater than control at 60 min of restraint. This expression declined by 180 min of restraint. Rats restrained for 15 min and given a 45 min recovery period (15–45) did not have significantly different expression of *fra-2* in the PVN than rats restrained for the full duration of time (60–0). Rats given a long recovery period (60–120) had significantly greater levels of *fra-2* expression in the PVN than rats restrained for the full duration of time (180–0). Figure 8 shows sample autoradiograms of *c-fos* and *fra-2* expression in the PVN for rats given 0–0, 15–0, 60–0, or 180–0 min of restraint.

Discussion

This study investigated the temporal-spatial expression pattern of the *fra-2* gene in rat forebrain in response to restraint. Expression of the *fra-2* gene was directly compared to the more widely studied gene family member *c-fos*. Extensive characterization of restraint-induced *c-fos* expression in rat forebrain has previously been reported (Imaki et al., 1993, Cullinan et al., 1995, Girotti et al., 2006, Trneckova et al., 2006). We found that *c-fos* gene expression in the PFC, LS, and PVN peaked at 15 min of restraint, declined some by 60 min of restraint and had reduced to near no-stress levels by 180 min of continuous restraint. A notable difference between *fra-2* and *c-fos* gene expression was the relatively high basal levels of *fra-2* mRNA in cortical and some subcortical structures. Under AM basal conditions, *fra-2* expression was moderate in the PFC and undetectable in the LS. In the PVN, however, there were high levels of basal *fra-2* gene expression. This finding contrasts with the low basal *fra-2* expression in the PVN described in a previous study (Honkaniemi et al., 1994). Both studies looked at unhandled adult male Sprague-Dawley rats during the A.M., so the discrepancy is not readily explainable. Because a number of immediate early genes (e.g. *c-fos*, *zif268*, *NGFI-B*) have very low basal expression in the PVN (Rivest and Rivier, 1994, Imaki et al., 1996, Umemoto et al., 1997), the relatively high basal *fra-2* gene expression in the PVN may reflect phenotypic specific control of *fra-2* basal expression rather than a downstream intracellular signaling response to the tonic presence of an extrinsic signal.

Restraint (60 min) produced a significant increase in *fra-2* mRNA in the PFC, septum and PVN. A trend for an increase was apparent in each of these brain regions within 15 min after restraint onset, but this increase was only statistically significant within the PVN. By 180 min of restraint, *fra-2* expression returned to near-basal levels in all brain regions examined, although it was still significantly elevated in prelimbic PFC and LS compared to no-stress. Thus, *fra-2* gene expression in rat brain appears to have immediate early gene properties, as described *in vitro* (Nishina et al., 1990), but the delay to reach peak expression may reflect an additional increase secondary to other IEG induction.

Although *fra-2* and *c-fos* are members of the same gene family (fos-family), as described above, basal and restraint-induced temporal induction patterns between the two genes were different. There is considerable structural similarity between promoters of the two genes. Both genes share a number of sequentially ordered putative enhancer elements in the 5' regulatory region (Yoshida et al., 1993, Foletta et al., 1994, Beer et al., 1998), allowing for the possibility of some overlap in *trans*-activation profiles. However, there are several key differences in the promoters of the two genes as well. The *c-fos* promoter contains two cyclic adenosine monophosphate (cAMP) binding protein response (CRE)-like elements upstream of the transcription start site (TATA box) that confers enhancement of *c-fos* expression via cAMP

(Hartig et al., 1991). The *fra-2* promoter lacks a TATA box and has only one CRE (Yoshida et al., 1993). The absence of this additional CRE may explain our observed relative lack of immediate (within 15 min of restraint) *fra-2* expression compared to *c-fos*. Interestingly, the *fra-2* promoter contains an AP-1 site next to the principal transcription start site (Yoshida et al., 1993) that may allow for *trans*-activation by other fos-family members participating in an AP-1 transcription factor complex. This may explain the late induction of *fra-2* after restraint onset: Fos, the protein product of *c-fos* mRNA, may confer (via an AP-1 element in the *fra-2* promoter) late induction of *fra-2*.

Differences in *fra-2* and *c-fos* basal and stress-induced gene expression patterns are likely to have important functional consequences. The protein products of both *c-fos* and *fra-2* share five key functional domains, although other protein subregions are much less similar (Nishina et al., 1990). These two proteins compete with one another to form one half of the heterodimeric AP-1 transcription factor complex, leading to changes in downstream expression of genes containing AP-1 elements in their promoter regions (Karin, 1995). Different AP-1 dimer combinations vary in their response element utilization and transactivation potency (Cuevas et al., 2005). For example, an AP-1 complex containing Fra-2 may have much less transactivational efficacy than one containing Fos, depending on the phosphorylation state of Fra-2 (Murakami et al., 1999). Consequently, the presence of Fra-2 in an AP-1 complex may antagonize Fos actions (Herdegen and Leah, 1998).

An additional feature of our study was the direct comparison of restraint duration on HPA axis hormone and gene expression levels at a given point in time after restraint onset. It has been observed before that HPA axis hormone secretion declines over the course of a single session of restraint, with some decrease evident within 60 min of restraint onset and substantial decreases after two hours of continuous restraint (Pecoraro et al., 2006). Although this within restraint-session habituation could be due to an exhaustion of CRH and/or ACTH peptide hormone reserves, one study found that after four hours of continuous restraint the exposure of rats to a novel stressor restored ACTH and corticosterone secretion to initial peak levels, indicating that some neural adaptation had occurred over the four hours of restraint (Dhabhar et al., 1997). It has similarly been observed that *c-fos* mRNA levels peak relatively soon (15–30 min) after the onset of restraint or immobilization regardless of stressor duration (Umemoto et al., 1997, Trneckova et al., 2007), as was also evident in this study. There is some evidence for *c-fos* expression to be inhibited through an autoregulatory negative feedback effect (Sassone-Corsi et al., 1988). Thus, the temporal pattern of stress-induced *c-fos* expression may be limited in the extent to which it reflects sustained neuronal activity. In this study we found that rats that had only 15 min of restraint had somewhat lower ACTH and CORT levels 60 min after restraint onset than rats that were restrained for the entire 60 min. Additionally, rats restrained continuously for 180 min had greater CORT levels than rats restrained for 60 min followed by 120 min of recovery. Rats restrained for only 15 min also had lower *c-fos* mRNA levels in the PVN and *fra-2* mRNA levels in the PFC than rats restrained continuously for 60 min, whereas in the lateral septum restraint duration did not affect gene expression. Because gene expression was greater in rats that were restrained continuously for 60 min than for 15 min, the expression time-course of both these genes must be able to somewhat reflect more than just the first 15 min of neural activation after stress onset. Crane et al (Crane et al., 2005) also observed a greater number of Fos positive CRF cells in the PVN of rats given 60 min of continuous restraint compared to rats given 15 min of restraint. Interestingly, our data suggests that the sensitivity of *c-fos* and *fra-2* expression to stress duration varies with brain region, and not necessarily in a parallel manner.

By 180 min after restraint onset there appears to be substantial within-session neural response habituation, such that in general there were limited differences in hormone or gene expression responses in rats restrained continuously compared to those restrained for only 60 min. This

eventual neural habituation to prolonged stressor exposure may account for the failure of another study to see differences in PVN *c-fos* mRNA levels of rats immobilized continuously for 120 min compared to rats immobilized for just the first 60 min of the 120 min period (Trneckova et al., 2007).

In summary, *fra-2*, compared to *c-fos*, has greater basal gene expression in select cortical and subcortical forebrain regions, with notably high basal expression in the PVN. Expression of *fra-2* gene expression is further increased by restraint in forebrain cortical and subcortical structures including the PFC, LS, and PVN. The relatively delayed induction pattern of *fra-2* compared with *c-fos* in these brain regions may suggest some divergent dependence of these two genes on cellular factors for *trans*-activation. The effect of stressor duration on *fra-2* and *c-fos* gene expression also differed across brain regions examined. Given the different transcriptional regulatory effects of Fra-2 or Fos protein participation in an AP-1 complex, these different basal and stress-dependent *fra-2* and *c-fos* gene expression patterns are likely to have important functional consequences for transcriptionally-dependent aspects of neuroplasticity.

Acknowledgements

We would like to thank Dr. Gregory Carey for excellent statistical advice. This work was supported by United States Public Health Service Grants MH075968 and MH065977.

References

- Beer J, Mielke K, Zipp M, Zimmermann M, Herdegen T. Expression of c-jun, junB, c-fos, fra-1 and fra-2 mRNA in the rat brain following seizure activity and axotomy. *Brain Res* 1998;794:255–266. [PubMed: 9622645]
- Butler TL, Pennypacker KR. Temporal and regional expression of Fos-related proteins in response to ischemic injury. *Brain Res Bull* 2004;63:65–73. [PubMed: 15121240]
- Campeau S, Watson SJ. Neuroendocrine and behavioral responses and brain pattern of c-fos induction associated with audiogenic stress. *J Neuroendocrinol* 1997;9:577–588. [PubMed: 9283046]
- Crane JW, French KR, Buller KM. Patterns of neuronal activation in the rat brain and spinal cord in response to increasing durations of restraint stress. *Stress* 2005;8:199–211. [PubMed: 16236624]
- Cuevas BD, Uhlik MT, Garrington TP, Johnson GL. MEKK1 regulates the AP-1 dimer repertoire via control of JunB transcription and Fra-2 protein stability. *Oncogene* 2005;24:801–809. [PubMed: 15558021]
- Cullinan WE, Herman JP, Battaglia DF, Akil H, Watson SJ. Pattern and time course of immediate early gene expression in rat brain following acute stress. *Neuroscience* 1995;64:477–505. [PubMed: 7700534]
- Dhabhar FS, McEwen BS, Spencer RL. Adaptation to prolonged or repeated stress--comparison between rat strains showing intrinsic differences in reactivity to acute stress. *Neuroendocrinology* 1997;65:360–368. [PubMed: 9158068]
- Engel L, Gupta BB, Lorenzkowski V, Heinrich B, Schwerdtle I, Gerhold S, Holthues H, Vollrath L, Spessert R. Fos-related antigen 2 (Fra-2) memorizes photoperiod in the rat pineal gland. *Neuroscience* 2005;132:511–518. [PubMed: 15802201]
- Foletta VC, Sonobe MH, Suzuki T, Endo T, Iba H, Cohen DR. Cloning and characterisation of the mouse fra-2 gene. *Oncogene* 1994;9:3305–3311. [PubMed: 7936655]
- Girotti M, Pace TW, Gaylord RI, Rubin BA, Herman JP, Spencer RL. Habituation to repeated restraint stress is associated with lack of stress-induced c-fos expression in primary sensory processing areas of the rat brain. *Neuroscience* 2006;138:1067–1081. [PubMed: 16431027]
- Hartig E, Loncarevic IF, Buscher M, Herrlich P, Rahmsdorf HJ. A new cAMP response element in the transcribed region of the human c-fos gene. *Nucleic Acids Res* 1991;19:4153–4159. [PubMed: 1651478]

- Hebert MA, Serova LI, Sabban EL. Single and repeated immobilization stress differentially trigger induction and phosphorylation of several transcription factors and mitogen-activated protein kinases in the rat locus coeruleus. *J Neurochem* 2005;95:484–498. [PubMed: 16190871]
- Herdegen T, Leah JD. Inducible and constitutive transcription factors in the mammalian nervous system: control of gene expression by Jun, Fos and Krox, and CREB/ATF proteins. *Brain Res Brain Res Rev* 1998;28:370–490. [PubMed: 9858769]
- Herman JP, Cullinan WE. Neurocircuitry of stress: central control of the hypothalamo-pituitary-adrenocortical axis. *Trends Neurosci* 1997;20:78–84. [PubMed: 9023876]
- Herman JP, Figueiredo H, Mueller NK, Ulrich-Lai Y, Ostrander MM, Choi DC, Cullinan WE. Central mechanisms of stress integration: hierarchical circuitry controlling hypothalamo-pituitary-adrenocortical responsiveness. *Front Neuroendocrinol* 2003;24:151–180. [PubMed: 14596810]
- Honkaniemi J, Kononen J, Kainu T, Pyykonen I, Pelto-Huikko M. Induction of multiple immediate early genes in rat hypothalamic paraventricular nucleus after stress. *Brain Res Mol Brain Res* 1994;25:234–241. [PubMed: 7808222]
- Imaki T, Shibasaki T, Chikada N, Harada S, Naruse M, Demura H. Different expression of immediate-early genes in the rat paraventricular nucleus induced by stress: relation to corticotropin-releasing factor gene transcription. *Endocr J* 1996;43:629–638. [PubMed: 9075602]
- Imaki T, Shibasaki T, Hotta M, Demura H. Intracerebroventricular administration of corticotropin-releasing factor induces c-fos mRNA expression in brain regions related to stress responses: comparison with pattern of c-fos mRNA induction after stress. *Brain Res* 1993;616:114–125. [PubMed: 8358602]
- Karin M. The regulation of AP-1 activity by mitogen-activated protein kinases. *J Biol Chem* 1995;270:16483–16486. [PubMed: 7622446]
- Kovacs KJ. c-Fos as a transcription factor: a stressful (re)view from a functional map. *Neurochem Int* 1998;33:287–297. [PubMed: 9840219]
- Liu Y, Chen GD, Lerner MR, Brackett DJ, Matsumoto RR. Cocaine up-regulates Fra-2 and sigma-1 receptor gene and protein expression in brain regions involved in addiction and reward. *J Pharmacol Exp Ther* 2005;314:770–779. [PubMed: 15879001]
- Murakami M, Ui M, Iba H. Fra-2-positive autoregulatory loop triggered by mitogen-activated protein kinase (MAPK) and Fra-2 phosphorylation sites by MAPK. *Cell Growth Differ* 1999;10:333–342. [PubMed: 10359014]
- Nishina H, Sato H, Suzuki T, Sato M, Iba H. Isolation and characterization of fra-2, an additional member of the fos gene family. *Proc Natl Acad Sci U S A* 1990;87:3619–3623. [PubMed: 2110368]
- Pacak K, Palkovits M. Stressor specificity of central neuroendocrine responses: implications for stress-related disorders. *Endocr Rev* 2001;22:502–548. [PubMed: 11493581]
- Paxinos, GWC. The rat brain in stereotaxic coordinates. Academic Press; San Diego: 1998.
- Pecoraro N, Ginsberg AB, Warne JP, Gomez F, la Fleur SE, Dallman MF. Diverse basal and stress-related phenotypes of Sprague Dawley rats from three vendors. *Physiol Behav* 2006;89:598–610. [PubMed: 16935312]
- Pennypacker KR, Eidzadeh S, Kassed CA, O'Callaghan JP, Sanberg PR, Willing AE. Expression of fos-related antigen-2 in rat hippocampus after middle cerebral arterial occlusion. *Neurosci Lett* 2000;289:1–4. [PubMed: 10899394]
- Rivest S, Rivier C. Stress and interleukin-1 beta-induced activation of c-fos, NGFI-B and CRF gene expression in the hypothalamic PVN: comparison between Sprague-Dawley, Fisher-344 and Lewis rats. *J Neuroendocrinol* 1994;6:101–117. [PubMed: 8025563]
- Robertson HA. Immediate-early genes, neuronal plasticity, and memory. *Biochem Cell Biol* 1992;70:729–737. [PubMed: 1482550]
- Sabban EL, Hebert MA, Liu X, Nankova B, Serova L. Differential effects of stress on gene transcription factors in catecholaminergic systems. *Ann N Y Acad Sci* 2004;1032:130–140. [PubMed: 15677400]
- Sassone-Corsi P, Ransone LJ, Lamph WW, Verma IM. Direct interaction between fos and jun nuclear oncoproteins: role of the 'leucine zipper' domain. *Nature* 1988;336:692–695. [PubMed: 3143919]
- Schwartz WJ, Carpino A Jr, de la Iglesia HO, Baler R, Klein DC, Nakabeppu Y, Aronin N. Differential regulation of fos family genes in the ventrolateral and dorsomedial subdivisions of the rat suprachiasmatic nucleus. *Neuroscience* 2000;98:535–547. [PubMed: 10869847]

- Sheng M, Greenberg ME. The regulation and function of c-fos and other immediate early genes in the nervous system. *Neuron* 1990;4:477–485. [PubMed: 1969743]
- Terao A, Greco MA, Davis RW, Heller HC, Kilduff TS. Region-specific changes in immediate early gene expression in response to sleep deprivation and recovery sleep in the mouse brain. *Neuroscience* 2003;120:1115–1124. [PubMed: 12927216]
- Trneckova L, Armario A, Hynie S, Sida P, Klenerova V. Differences in the brain expression of c-fos mRNA after restraint stress in Lewis compared to Sprague-Dawley rats. *Brain Res* 2006;1077:7–15. [PubMed: 16487948]
- Trneckova L, Rotllant D, Klenerova V, Hynie S, Armario A. Dynamics of immediate early gene and neuropeptide gene response to prolonged immobilization stress: evidence against a critical role of the termination of exposure to the stressor. *J Neurochem*. 2007
- Umemoto S, Kawai Y, Ueyama T, Senba E. Chronic glucocorticoid administration as well as repeated stress affects the subsequent acute immobilization stress-induced expression of immediate early genes but not that of NGFI-A. *Neuroscience* 1997;80:763–773. [PubMed: 9276492]
- Weinberg, M.; Girotti, M.; Francis, A.; VanElzakker, M.; Spencer, RL. Fra-1 in the prefrontal cortex: stress regulation of immediate early gene expression. *Proceedings of the Society for Neuroscience*; Washington, DC. 2005.
- Yoshida T, Suzuki T, Sato H, Nishina H, Iba H. Analysis of fra-2 gene expression. *Nucleic Acids Res* 1993;21:2715–2721. [PubMed: 7687344]

Abbreviations

ACTH	adrenal corticotropin hormone
CORT	corticosterone
CRE	cyclic adenosine monophosphate binding protein response element, HPA axis, hypothalamus-pituitary-adrenal axis
IEG	immediate-early gene
IL	infralimbic prefrontal cortex
LS	lateral septum
PFC	prefrontal cortex
PL	prelimbic prefrontal cortex
PVN	paraventricular nucleus of the hypothalamus
SCM	sis-conditioned medium
SCN	suprachiasmatic nucleus
SRE	

VO

serum response element

ventral-orbital prefrontal cortex

Figure 1.

ACTH and CORT plasma levels after different periods of restraint. ACTH (top) and CORT (bottom) were measured in the plasma of rats at 0, 15, 60, or 180 min of restraint stress (●). Two additional data points represent plasma hormone levels of rats given a recovery period post-restraint but equal total time from restraint onset to death. One group was given 15 min of restraint followed by 45 min of recovery (Δ), and another group was given 60 min of restraint followed by 120 min of recovery (▲).*, significantly different from home-cage control (0 min); Fisher's LSD, $P<0.05$. #, significantly different from the group given equal total time from restraint onset to death; independent samples t-test, $P<0.05$.

Figure 2.

Basal *c-fos* and *fra-2* mRNA expression in the rat forebrain. Top row illustrates neuroanatomical subregions of coronal sections collected at the rostral-caudal level of the prefrontal cortex (PFC), lateral septum (LS), and hypothalamic paraventricular nucleus (PVN) (adapted from Paxinos and Watson (Paxinos G, 1998)). Numbers indicate regions of interest for densitometry analysis: 1 = prelimbic PFC, 2 = infralimbic PFC, 3 = ventral orbital PFC, 4 = LS, 5 = PVN. Letters indicate additional brain regions with relatively high basal *fra-2* mRNA: a = piriform cortex, b = nucleus accumbens shell, c = endopiriform nucleus, d = supraoptic nucleus, e = hippocampus, f = dentate gyrus. The bottom row presents representative autoradiogram images of *c-fos* expression (left half of each coronal section) and *fra-2* expression (right side of each coronal section) for comparison purposes.

Figure 3. *c-fos* and *fra-2* mRNA relative levels in the prefrontal cortex after different periods of restraint. *c-fos* (top) and *fra-2* (bottom) expression were measured in the PL (left), IL (middle), and VO (right) of the prefrontal cortex of rats at 0, 15, 60, or 180 min of restraint stress (●). Two additional data points represent mRNA levels of rats given a recovery period post-restraint but equal total time from restraint onset to death. One group was given 15 min of restraint followed by 45 min of recovery (Δ), and another group was given 60 min of restraint followed by 120 min of recovery (▲).*, significantly different from home-cage control (0 min); Fisher's LSD, $P < 0.05$. #, significantly different from the group given equal total time from restraint onset to death; independent samples t-test, $P < 0.05$.

Figure 4.

Representative autoradiograms of stress-induced *c-fos* and *fra-2* gene expression in the prefrontal cortex. Rats were exposed to 0, 15, 60, or 180 min of restraint. *c-fos* expression (top) is evident throughout the PL, IL, and VO by 15 min of restraint and has diminished by 180 min of restraint. *fra-2* expression (bottom) significantly increases above control levels in the PL, IL, and VO after 60 min of restraint, and diminishes by 180 min of restraint.

Figure 5.

c-fos and *fra-2* mRNA relative levels in the lateral septum (LS) after different periods of restraint. *c-fos* (top) and *fra-2* (bottom) expression were measured in the LS of rats after 0, 15, 60, or 180 min of restraint stress (●). Two additional data points represent mRNA levels of rats given a recovery period post-restraint but equal total time from restraint onset to death. One group was given 15 min of restraint followed by 45 min of recovery (Δ), and another group was given 60 min of restraint followed by 120 min of recovery (▲).*, significantly different from home-cage control (0 min); Fisher's LSD, $P < 0.05$. #, significantly different from the group given equal total time from restraint onset to death; independent samples t-test, $P < 0.05$.

Figure 6.

Representative autoradiograms of stress-induced *c-fos* and *fra-2* gene expression in the lateral septum. Rats were exposed to 0, 15, 60, or 180 min of restraint. *c-fos* expression (top) is evident by 15 min of restraint and has diminished by 180 min of restraint. *fra-2* expression (bottom) significantly increases above control levels after 60 min of restraint, and diminishes by 180 min of restraint.

Figure 7.

c-fos and *fra-2* mRNA relative levels in the hypothalamic paraventricular nucleus (PVN) after different periods of restraint. *c-fos* (top) and *fra-2* (bottom) expression were measured in the PVN of rats at 0, 15, 60, or 180 min of restraint stress (●). Two additional data points represent mRNA levels of rats given a recovery period post-restraint but equal total time from restraint onset to death. One group was given 15 min of restraint followed by 45 min of recovery (Δ), and another group was given 60 min of restraint followed by 120 min of recovery (▲).*, significantly different from home-cage control (0 min); Fisher's LSD, $P < 0.05$. #, significantly

different from the group given equal total time from restraint onset to death; independent samples t-test, $P < 0.05$.

Figure 8.

Representative autoradiograms of stress-induced *c-fos* and *fra-2* gene expression in the hypothalamic paraventricular nucleus. Rats were exposed to 0, 15, 60, or 180 min of restraint. *c-fos* expression (top) is evident by 15 min of restraint and has diminished by 180 min of restraint. *fra-2* expression (bottom) is evident even in home-cage control animals (0 min restraint). *fra-2* expression significantly increases above control levels within 15 min of restraint, and diminishes by 180 min of restraint.