Persistent Calcium Elevation Correlates with the Induction of Surface Immunoglobulin-mediated B Cell DNA Synthesis

By Hidehiro Yamada,^{*1} Carl H. June,^{*1} Fred Finkelman,[‡]
Mark Brunswick,[§] Michael S. Ring,^{*||} Andrew Lees,^{*‡}
and James J. Mond^{‡1}

From the Immune Cell Biology Program, Naval Medical Research Institute, Bethesda, Maryland 20889-5055; the [‡]Department of Medicine, Uniformed Services University of the Health Sciences, Bethesda, Maryland 20814-4799; the [§]Laboratory of Cell Biology, Food and Drug Administration, Bethesda, Maryland 20892; and ^{||}Geo-Centers, Inc., Ft. Washington, Maryland 20744

Summary

Surface immunoglobulin (sIg)-mediated stimulation of B lymphocytes induces a tyrosine kinase-dependent sequence of events leading to rapid and large elevations in intracellular ionized calcium ($[Ca^{2+}]_i$). These early biochemical events do not necessarily lead to proliferation of B cells, however, and conversely, the absence of or inhibition of these events does not necessarily prevent cellular proliferation. We now show by digital image analysis of single B cells that conditions which lead to B cell proliferation are associated with low-level but persistent sustained or cyclic elevations in $[Ca^{2+}]_i$. In marked contrast, early and nonsustained elevations in $[Ca^{2+}]_i$ are induced in B cells by stimuli that lead to G1 transition but fail to progress to DNA synthesis. Thus, when B cells were stimulated with mitogenic and nonmitogenic anti-IgD antibodies, both of which induce entry of cells into G1 and early calcium transients of comparable magnitude, persistent low-level calcium elevations were only detected in cells stimulated with the mitogenic antibody. Furthermore, persistent calcium elevations were also seen when B cells were stimulated with a multivalent dextran-anti-Ig conjugate which induced very high levels of B cell proliferation in the absence of detectable phosphatidylinositol 4,5-bisphosphate hydrolysis or elevations in $[Ca^{2+}]_i$ as detected by flow cytometry. Finally, B cells from X-linked B cell-defective mice, which do not proliferate in response to anti-Ig antibody, show marked and early increases in $[Ca^{2+}]_i$, but do not show persistent calcium elevations. These data suggest that the rapid and large increases of $[Ca^{2+}]_i$ seen in lymphocytes within seconds after antigen receptor ligation may be associated with entry in G1, whereas low-level but persistent elevations may be the hallmark of a cell destined to synthesize DNA.

Crosslinking of surface immunoglobulin (sIg)² receptors in B cells initiates a cascade of biochemical events that includes mobilization of intracellular ionized calcium $[Ca^{2+}]_i$ and activation of protein kinase C (PKC) and tyrosine kinases (1-7). Activation of PKC and tyrosine kinase in turn activates a number of other enzyme systems, including phosphatidylinositol 3 and mitogen-activated protein kinases, guanosine triphosphate-activating protein, and raf-1 protein

kinase (8-10), all of which are intimately involved in cell growth of B cells as well as other cell types. Although it is clear that the summation of these events reflects an early response to sIg-mediated signal transduction in B cells which will lead to cell growth, the specific role of each event in regulating entry of the cell into G1 or S of the cell cycle is not known. Experiments both in human and in murine systems suggest that there is no correlation between the ability of an antigen or of anti-Ig antibody to induce phosphatidylinositol 4,5-bisphosphate (PIP₂) activation or tyrosine kinase activation and its ability to induce B cell DNA synthesis (11-14). Thus, for example, inhibition of sIg-mediated PIP₂ hydrolysis does not prevent B cell proliferation (15). Furthermore, stimulation of these early events by anti-Ig antibodies does not necessarily lead to B cell proliferation (13, 16, 17).

¹ H. Yamada, C. H. June, and J. J. Mond contributed equally to the work.

² Abbreviations used in this paper: $[Ca^{2+}]_i$, intracellular free calcium concentration; PIP₂, phosphatidylinositol 4,5-bisphosphate; PKC, protein kinase C; sIg, surface Ig; TI, T cell-independent antigen.

An added level of complexity in understanding the role for these early biochemical events is created when trying to dissect differences between sIg-mediated events stimulated by T cell-dependent (i.e., mono or paucivalent) as compared with T cell-independent (multivalent and higher molecular weight) antigens (TI). Although both antigen types are similar in their abilities to induce the early events in B cell activation, T cell-dependent but not -independent, antigens require B cells to interact with T cells or T cell-derived cytokines to stimulate proliferation and Ig secretion.

We have recently described a model system utilizing dextran-conjugated anti-Ig antibodies which stimulates polyclonal activation of B cells via sIg ligation and crosslinking and mimics, in many respects, activation that is stimulated by TI type 2 antigens (18). These dextran conjugates stimulate high levels of B cell proliferation at picogram per milliliter concentrations that are not accompanied by detectable PIP₂ hydrolysis and tyrosine phosphorylation whereas higher concentrations of this ligand stimulate PIP₂ hydrolysis and tyrosine phosphorylation (11, 12, 16). Digital image analysis has made it possible to examine, at the level of the single B cell, some early activation events that may be masked when they are studied in cell populations (19). The experiments described in this manuscript demonstrate that sIg crosslinking-mediated induction of low-level and temporally persistent cyclic elevations in intracellular free calcium concentration ($[Ca^{2+}]_i$) is a more critical determinant than the early induction of large increases in $[Ca^{2+}]_i$ for stimulating B cells to synthesize DNA.

Materials and Methods

Mice. DBA/2 mice (the Jackson Laboratory, Bar Harbor, ME) were used at 8 wk of age.

B Cell Purification. Suspensions of single spleen cells were washed

three times with RPMI 1640 (M.A. Bioproducts, Walkersville, MD) plus 10% FCS (GIBCO, Grand Island, NY) and treated with the following antibodies: monoclonal anti-Thy-1.2 (1 μ g/ml) (20), anti-CD4 (2.5 μ g/ml) (21), and anti-CD8 (1:400 dilution of culture supernatant) (20), respectively, per 10⁷ spleen cells for 30 min on ice. This was followed by treatment with newborn rabbit complement (10%) (Pel-Freez Biologicals, Rogers, AR), in the presence of a 1:10 dilution of tissue culture fluid containing the anti-rat κ chain mAb, mouse anti-rat 18.5 (22), at 37°C for 45 min.

Cells were then fractionated into low- and high-density populations by Percoll gradient centrifugation (Pharmacia Fine Chemicals, Piscataway, NJ; 3,000 rpm for 15 min). Gradients consisting of 70, 65, 60, and 50% Percoll (with densities of 1.086, 1.081, 1.074, and 1.062 g/ml, respectively) were used. The high-density cells (resting) were collected from the 70–65% interface. The average percentage of Ig⁺ cells in the 70–65% fractions was 90–95%.

Fluorescence Digital Microscopy. Details of the image analysis system have been described (23, 24). Briefly, 5×10^6 cells were loaded with 1.5 μ M fura-2 for 30 min at 31°C, washed, and suspended in HBSS with 0.1% FCS, 5×10^5 M 2-ME and 20 mM Hepes, pH 7.4. Cells were allowed to settle onto glass coverslips and equilibrated in a 2-ml microscope chamber at 37°C. Agonistic antibodies were gently added as a 10 $\times$ solution to the chamber. The fluorescence digital image processing system used is similar to that described previously by Tsien and Harootunian (23). Hardware consists of an image processor (model FD5000; Gould Inc., Glen Burnie, MD), a microscope (Axiovert; Zeiss, Oberkochen, Germany) with a 1.3 numerical aperture $\times 40$ objective, and a filter changer (Ludl Electronic Products, Hawthorne, NY) with excitation filters centered at 350 ± 10 nm and 380 ± 10 nm. The image processor and filter changer are interfaced to a host computer (Microvax; Digital Equipment Corp., Marlboro, MA). 16 images were acquired at each wavelength at 30/s through a camera (CCD; Cohu Inc., San Diego, CA) and image intensifier (Video Scope Intl., Ltd., Wash, DC). The images were averaged, background subtracted, and a shading correction applied. The ratio of the 350- and 380-nm images is displayed according to a code kindly provided by R. Y. Tsien (University of California, San Diego, CA). Images

Figure 1. Fura-2 B cells were settled on coverslips and stimulated with 0.5 μ g/ml or 0.2 ng/ml of H δ 1/-dextran. Results show the $[Ca^{2+}]_i$ of individual B cells for the first 6 min after stimulation; there were 35 and 36 cells in the left and right panels, respectively. Each line represents the $[Ca^{2+}]_i$ of an individual cell analyzed over 6 min. This experiment is one of three representative experiments.

Figure 2. B cells were treated as in Fig. 1 and evaluated by image analysis at 90 min after the addition of $H\delta^A/1$ -dextran. (Circles) The $[Ca^{2+}]_i$ of an individual cell at 90 min; 105–120 cells per condition are plotted. Bars represent the mean $[Ca^{2+}]_i \pm$ SEM. This experiment is one of four representative experiments.

were acquired and stored digitally at 15-s intervals. The fura-2 ratio is converted to $[Ca^{2+}]_i$ as described (23).

Representation of Data. The error bars in the graphs represent the SE of the mean $[Ca^{2+}]_i$ of individual cells.

Results

Short-Term Calcium Transients Stimulated by Picogram Quantities of Anti-Ig Antibody. We have previously reported that picogram per milliliter concentrations of dextran-conjugated anti-Ig antibody (anti- δ -dex), which stimulate high levels of B cell DNA synthesis, do not stimulate detectable increases

Figure 3. Frequency of responding cells late after stimulation. Data from the experiment shown in Fig. 2 were analyzed for the cumulative frequency of responding cells. Cells in one microscopic field were analyzed beginning 90 min after stimulation for a further 5 min. Cells were considered as responding cells if the $[Ca^{2+}]_i$ exceeded $0.3 \mu M$. Cells were cultured in medium (.....), $H\delta^A/1$, 0.2 ng/ml (----), or $H\delta^A/1$, $0.5 \mu g/ml$ (—). The total number of cells per field for the three conditions was 35, 40, and 35, respectively.

in $[Ca^{2+}]_i$ as measured by flow cytometry (11). In contrast, microgram per milliliter concentrations of this antibody retained mitogenicity and stimulated early onset $[Ca^{2+}]_i$ elevations that were high in magnitude. To evaluate whether analysis of a population of cells by standard flow cytometric approaches might be masking small but significant perturbations of $[Ca^{2+}]_i$ after B cell activation, we studied sIg-mediated B cell activation by digital image analysis. This technique allows one to study events of single cells over real time and to detect oscillatory responses that may not be apparent by flow cytometry (23, 24). Purified resting B cells were allowed to settle for 5 min onto coverslips and then placed in thermostatically controlled chambers at $37^\circ C$. Anti- δ -dex at $0.5 \mu g/ml$ stimulated rapid increases in $[Ca^{2+}]_i$ in $>90\%$ of the B cells to levels that were, on average, 8–10-fold greater than those of unstimulated B cells (Fig. 1, left). This result is similar to that which we reported previously using flow cytometric analysis of indo-1-loaded B cells (11). As has previously been described, B cell $[Ca^{2+}]_i$ levels showed either sustained or oscillatory patterns of calcium elevation (19). The magnitude of the response of each cell and its oscillatory frequency exhibited substantial heterogeneity for the ~ 200 cells examined. When B cells were stimulated with highly mitogenic but 1,000-fold lower concentrations of anti- δ -dex (0.2

Figure 4. Histograms of peak $[Ca^{2+}]_i$ late after stimulation. Cells from one microscopic field from the experiment shown in Figs. 2 and 3 were analyzed 90 min after stimulation for a further 5 min. Data were obtained at 15-s intervals, and histograms of the peak $[Ca^{2+}]_i$ for each cell are plotted. Cells were cultured in medium, $H\delta^A/1$, 0.2 ng/ml , or $H\delta^A/1$, 500 ng/ml . The total number of cells per field for the three conditions were 35, 40, and 35, respectively.

Figure 5. B cells were stimulated with the mitogenic H δ a/1 anti- δ antibody or the nonmitogenic FF1 anti- δ antibody. $[Ca^{2+}]_i$ was evaluated immediately after the addition of anti-Ig antibody (three pseudo-color photos, *right*) or 60 min after the addition of antibody (three pseudo-color photos, *left*). Chambers were kept in an incubator at 37°C, 5% CO₂ for the interim 60 min. Because of the necessity of maintaining cells in an incubator for this prolonged period of time, a different field of cells was analyzed at 0 time and at 60 min. However, for each time point, three separate fields of at least 30 cells/field were analyzed. Results are depicted for one representative field. The color legend (*right*) reflects the $[Ca^{2+}]_i$ of each cell, with blue being the lowest and violet the highest $[Ca^{2+}]_i$. The halo at the periphery of some cells is an artifact and does not indicate a spatial intracellular $[Ca^{2+}]_i$ gradient.

ng/ml), smaller increases in $[Ca^{2+}]_i$ were also consistently observed (Fig. 1, *right*). In our previous flow cytometric studies, we were unable to detect a response in B cells stimulated with this low dose of anti- δ -dex (11). The response detected by digital microscopy was more apparent when the results were depicted using a smaller scale (Fig. 1, *inset*). Whereas, at any given time, <25% of B cells stimulated with low concentrations of anti- δ -dex showed increases in $[Ca^{2+}]_i$, >90% of B

cells showed significant responses in the five min after stimulation. Although the amplitude of the oscillations under these conditions was small, the frequency was not appreciably different than that stimulated by the higher concentrations of anti- δ -dex. This suggested that the persistence of oscillations may be a more critical factor than the magnitude of initial calcium elevation in determining the calcium-dependent events that lead to B cell activation.

Figure 5. (continued)

Prolonged Calcium Oscillations Stimulated by Mitogenic but Not by Nonmitogenic Concentrations of Anti- δ Antibodies. The reports that B cell proliferation stimulated by anti-Ig antibody appears to require prolonged exposure to antigen or anti-Ig antibody suggested that persistent signalling was a critical factor in directing a B cell to DNA synthesis (25). To study this point directly, we evaluated calcium transients at both early and late time points after B cell stimulation by mitogenic amounts of anti- δ -dex antibody at high (500 ng/ml) or low (0.2 ng/ml) concentrations. By 90 min after activation, the mean increase in $[Ca^{2+}]_i$ stimulated by the high concentration of anti- δ -dex had fallen significantly from

its peak level seen seconds after stimulation, whereas that stimulated by the low concentration stayed relatively constant over the 90 min of observation (Fig. 2). In six experiments of this design, the mean increase in $[Ca^{2+}]_i$ in cells 90 min after stimulation with 500 ng/ml was either the same as, or only marginally greater than that stimulated by 0.2 ng/ml.

The above results were obtained by observing three microscopic fields for each condition at one time point 90 min after stimulation. It was necessary to sum the results from several fields of cells for statistical purposes. However, observation of the same field of cells as a function of time at 15-s

intervals supplies additional information, and reveals that the response pattern of individual cells is dynamic, and consists of cells with both sustained elevations of $[Ca^{2+}]_i$ and cells with oscillatory elevations (19). When the response of a particular field of cells was analyzed as a function of time, observations were necessarily limited to the number of cells contained in one field of cells. Typically, there are 30–50 cells. In one experiment, it was found at the first observation point that there were 23% of cells with increased $[Ca^{2+}]_i$ in the group cultured for 90 min in $H\delta^a/1$ -dex at 0.5 $\mu\text{g/ml}$, whereas there were no cells with elevated $[Ca^{2+}]_i$ in the groups cultured in low-dose $H\delta^a/1$ -dex 0.2 $\mu\text{g/ml}$ or in medium. However, over the next 5 min, there was a progressive increase in the fraction of cells that had elevated $[Ca^{2+}]_i$, reaching 69 and 38% in the high- and low-dose $H\delta^a/1$ conditions, whereas only 2 of 35 cells (6%) of cells cultured in medium had elevated $[Ca^{2+}]_i$ (Fig. 3). Histograms of these responses showed that the magnitude of peak calcium increases in the time period of 90–95 min after stimulation was similar in the low- and high-dose $H\delta^a/1$ -dex groups (Fig. 4). Thus, the magnitude of calcium elevations late after stimulation appears to be similar in both the high- and low-dose $H\delta^a/1$ -dex conditions. We are unable to determine if the frequency of responding cells differs between the two groups. During the 5 min of analysis, the high-dose condition appears to approach a plateau, whereas the plot of low-dose condition suggests that a higher fraction of responding cells might be observed with longer periods of observation (Fig. 3). Currently, we are unable to distinguish between these possibilities because of leakage of the calcium probe with prolonged cell culture. These results indicate that high-dose $H\delta^a/1$ causes large and early transient increases in B cell calcium whereas low-dose $H\delta^a/1$ results in only minimal early increases in calcium that are detectable by video microscopy (Fig. 1) and not by flow cytometry (11). In contrast, when B cells are examined 60–90 min after stimulation, both doses of $H\delta^a/1$ result in calcium elevations in a substantial fraction of cells (Fig. 3), of which the magnitude of the elevations is similar (Figs. 2 and 4). Together, these results suggested that persistent and prolonged signalling, as reflected by low-level calcium transients, was a critical element in determining whether a cell would proceed to mitogenesis. Alternatively, this form of signal transduction might serve as a marker of cells that are committed to mitogenesis.

To further test this, we compared the ability of a mitogenic ($H\delta^a/1$) and a nonmitogenic (FF1-4D5) monoclonal anti- δ antibody (26) to stimulate immediate as well as prolonged elevations in $[Ca^{2+}]_i$. Unconjugated $H\delta^a/1$, in contrast to the dextran conjugate, is mitogenic over a narrower concentration range of 1–100 $\mu\text{g/ml}$ (11, 18). Both $H\delta^a/1$ and FF1-4D5 stimulated comparable increases in $[Ca^{2+}]_i$ within seconds after stimulation (Fig. 5). In marked contrast, by 60 min after stimulation, cells stimulated by the mitogenic, but not the nonmitogenic, anti- δ antibody showed small but significant elevations in calcium (Figs. 5–7). These prolonged cyclic elevations were not seen when nonmitogenic concentrations of $H\delta^a/1$ (0.02 $\mu\text{g/ml}$) were employed (Fig. 7). When B cells from C57BL/6 mice, which express Ig of

Figure 6. Experimental design same as in Fig. 5, with results depicted as a scattergram. (Circles) The $[Ca^{2+}]_i$ of one cell at 60 min after stimulation. The bars represent mean $[Ca^{2+}]_i \pm \text{SEM}$. In the medium, $H\delta^a/1$, and FF1 cultures, there were 80, 120, and 100 cells, respectively.

the b allotype and thus are nonreactive to $H\delta^a/1$ (anti-allotype IgD^a) were examined, no short- or long-term increase in calcium transients was observed, indicating that these low-level oscillations are not attributable to artifacts of the in vitro culture system (Fig. 8). These results suggest that persistent but low-level elevations in $[Ca^{2+}]_i$ are associated with stimulation of B cells by mitogenic but not nonmitogenic anti-Ig antibodies.

Absence of Long-Term Calcium Transients in Anti-Ig-stimulated B Cells from *xid* Mice. CBA/N mice have been shown to have an X-linked immune defect (*xid*) and do not respond to immunization with TI type 2 antigens (27). Furthermore, whereas unconjugated anti-Ig antibody does not stimulate proliferation in B cells from *xid* mice, dextran-conjugated anti-Ig antibody stimulates high levels of proliferation (13). In view of this finding, we wished to test whether dextran-conjugated antibody, but not unconjugated anti-Ig, would stimulate long-term calcium transients in B cells from *xid*

Figure 7. Fura-loaded B cells were cultured in medium (0), or stimulated with a mitogenic (10 $\mu\text{g/ml}$) or nonmitogenic (0.02 $\mu\text{g/ml}$) concentration of unconjugated $H\delta^a/1$ and analyzed at 90 min. In the 0, 0.02, and 10 $\mu\text{g/ml}$ conditions, there were 100, 90, and 90 cells, respectively. This experiment was one of three representative experiments.

Figure 8. This experiment depicts the $[Ca^{2+}]_i$ response in 30 B cells from DBA/2 cells (allotype Ig^a) that are reactive with and stimulated by H δ^a /1 (top right), and in 20 B cells from C57BL/6 mice (allotype Ig^b) that are nonreactive and not stimulated by H δ^a /1 antibody (bottom right). The results (bottom left) demonstrate the $[Ca^{2+}]_i$ response in 25 B cells from DBA/2 mice 60 min after stimulation with the nonmitogenic FF1, an anti- δ antibody which stimulates the early but not the late phase of calcium elevation. (Top left) The calcium levels of 20 DBA/2 spleen B cells cultured in medium alone.

mice. In agreement with our previous flow cytometric studies (13), we found that both H δ^a /1 and H δ^a /1-dex stimulated rapid increases in $[Ca^{2+}]_i$ in B cells from normal and *xid* mice, albeit to a lesser degree in the latter (Fig. 9). By 90 min after stimulation, low-level but significant increases were observed in B cells from normal mice that were stimulated both with H δ^a /1 and with H δ^a /1-dex. In contrast, only H δ^a /1-dex, but not unconjugated H δ^a /1, stimulated long-term elevations in $[Ca^{2+}]_i$ in B cells from *xid* mice (Fig. 10). This finding lends additional support to the hypothesis that the ability of sIg ligation to stimulate B cell proliferation may depend on its ability to stimulate long-term calcium transients.

Figure 9. B cells from normal or *xid* mice were stimulated with H δ^a /1 or H δ^a /1-dextran, and the immediate increase in $[Ca^{2+}]_i$ was recorded for 6 min after stimulation. The data represents the mean increase in $[Ca^{2+}]_i$ of 50–60 individual cells followed for 6 min after stimulation.

Figure 10. B cells from normal or *xid* mice were stimulated with H δ^a /1 or H δ^a /1-dextran, and the $[Ca^{2+}]_i$ was recorded at 90 min after stimulation. The data shows the response of each individual cell, each circle representing the $[Ca^{2+}]_i$ of one cell analyzed at 90 min after stimulation. In the medium, H δ^a /1, and H δ^a /1-dextran conditions, there were 96, 156, and 78 cells (left) and 72, 99, and 180 cells (right), respectively. The bars represent mean $[Ca^{2+}]_i \pm$ SEM.

Discussion

The major parameters that influence sIg-mediated B cell activation are the extent and duration of crosslinking of sIg. These parameters are themselves influenced by the valency and avidity of the ligand for sIg and sIg receptor density (26, 28–31). At the onset of activation, when resting B cells express a high density of sIg, lower avidity antibodies are capable of initiating early steps in B cell activation. This enables paucivalent T cell-dependent antigens to crosslink sIg, albeit relatively ineffectively, and stimulate B cells to rapidly express increased MHC class II, which facilitates antigen presentation (32). In the absence of added cytokines or T cell help, these B cells do not progress to DNA synthesis. In contrast, multivalent antigens, as exemplified by the TI type 2 antigens, bind with higher avidity to sIg and establish effective domains of crosslinked sIg, even when ligand-induced modulation of sIg causes a decrease in Ig density. Stimulation of cells with TI type 2 antigens can therefore result in persistent and repetitive signalling, which leads to B cell proliferation even at extremely low antigen concentrations. In our previous studies with dextran-conjugated anti-Ig antibodies, we found that picogram per milliliter concentrations of conjugate stimulated high levels of B cell activation in the absence of detectable elevations in $[Ca^{2+}]_i$ (11). This suggested either that PIP₂-dependent pathways of activation were being stimulated that were below the threshold of our detecting systems, or that at such low concentrations of agonist different activation pathways were being recruited. To distinguish between these two possibilities, we used digital image analysis to study calcium transients in B cells stimulated by low concentrations of anti- δ dextran conjugates. Furthermore, we hoped to distinguish between the biochemical events stimulated by sIg ligation by nonmitogenic vs. mitogenic anti-Ig antibodies. As anticipated from earlier studies, both stimuli induced rapid and large early increases in $[Ca^{2+}]_i$. However, only the mitogenic antibodies stimulated prolonged increases that were ob-

served beyond 60 min after the initial activation. Nonmitogenic anti-Ig antibodies, which stimulated PIP_2 hydrolysis comparable to that induced by mitogenic anti-Ig antibodies, did not stimulate persistent increases in calcium. Likewise, neither persistent calcium transients nor DNA synthesis were observed in *xid* B cells incubated with anti-Ig antibody, although this treatment stimulated *xid* B cells to rapidly and transiently increase $[\text{Ca}^{2+}]_i$. Conversely, low concentrations of anti-Ig-dextran, which induced no detectable PIP_2 hydrolysis, stimulated prolonged and persistent increases in calcium as well as DNA synthesis.

We found that low concentrations of $\text{H}\delta^2/1$ -dextran induced $[\text{Ca}^{2+}]_i$ transients that were asynchronous and often oscillatory in nature (Fig. 1). After 90 min of stimulation, <5% of cells had elevated $[\text{Ca}^{2+}]_i$ at any given time. However, during a 5-min period of observation, about 40% of cells showed a response (Fig. 3). This provides a partial explanation as to why we were previously unable to detect this response by flow cytometry (11). Furthermore, for flow cytometric analysis, B cells were loaded with larger concentrations of indo-1 than the fura-2 concentrations used for our present studies. These larger concentrations may have buffered or dampened the low-level calcium responses.

The ability to stimulate persistent calcium transients is related to the ability of a ligand to bind and crosslink sIg over a prolonged period of time. This can be accomplished by high concentrations of bivalent anti-Ig antibody, which can create multiple domains of tightly crosslinked sIg even on partially activated B cells which express few membrane Ig receptors. At low ligand concentrations, however, this can only be accomplished by a multivalent ligand (anti- δ -dex), which binds with a high avidity to a large number of membrane Ig molecules. The ability of low concentrations of the multivalent ligands to crosslink multiple sIg molecules allows for effective signalling without modulating a large percentage of sIg molecules. Thus, when B cells are cultured with low concentrations of anti-Ig-dex, the density of sIg molecules that can be ligated as cell culture continues is maintained at a much higher level than when B cells are cultured with high

concentrations of unconjugated anti-Ig antibody. The importance of maintenance of a high sIg density is suggested by: (a) the much greater magnitude of DNA synthesis of B cells cultured with low concentrations of anti-Ig-dex than with high concentrations of unconjugated anti-Ig; (b) the ability of preincubation of B cells with unconjugated anti-Ig to inhibit anti-Ig-dex-induced B cell proliferation; and (c) the reduced ability of high concentrations of anti-Ig-dex, which modulate most sIg, to stimulate DNA synthesis. In this regard, it is interesting that unconjugated anti-Ig stimulates comparable increases in $[\text{Ca}^{2+}]_i$ at 90 min after stimulation to that stimulated by anti-Ig-dex. This suggests that analysis at time points after 90 min may reveal additional differences between these two mitogens or that other coassociated molecules may be more strongly stimulated when sIg is engaged by a strongly crosslinking molecule.

The finding of agonist-stimulated repetitive and asynchronous calcium oscillations within seconds after sIg crosslinking has been previously reported in many cellular systems, including B cells (19). Although other investigators studying other cellular systems have suggested that the frequency of oscillation may influence cell growth (for a review see reference 33), our studies do not at this point support this notion for B cell activation.

For many different cellular systems, rapid and large increases in $[\text{Ca}^{2+}]_i$ may be both necessary and sufficient. Thus, for example, the rapid release of histamine from mast cells may require only a rapid but not prolonged increase in $[\text{Ca}^{2+}]_i$. Likewise, rapid and large calcium elevations may be a critical element for B cells for the induction of events that may occur shortly after sIg crosslinking as, for example, enhanced antigen presentation or possibly lymphokine secretion. Our data indicate that persistent calcium elevations may be indispensable for cellular events that require prolonged signalling as, for example, induction of cellular mitogenesis. Alternatively, the present results are compatible with the notion that persistent calcium signal transduction is a marker of a cell committed to mitogenesis, and further studies will be required to distinguish between these possibilities.

We are indebted to Mary Chase for help in preparing the manuscript.

This work was supported in part by Uniformed Services University Protocol RO83DR; National Institutes of Health grants AI-27465 and AI-21328; and Naval Medical Research and Development Command grants N0007592WR00025, N0007592WR00024, and MR04120.001.1011. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of Defense, or the U.S. Government.

Address correspondence to Dr. C. H. June, Mail Stop 44, Immune Cell Biology Program, Naval Medical Research Institute, Bethesda, MD 20889-5055. H. Yamada is currently at the Institute of Medical Science, St. Marianna School of Medicine, Kawasaki 216, Japan.

Received for publication 22 June 1992 and in revised form 8 January 1993.

References

- Maino, V.C., M.J. Hayman, and M.J. Crumpton. 1975. Relationship between enhanced turnover of phosphatidylinositol and lymphocyte activation by mitogens. *Biochem. J.* 146:247.
- Ransom, J.T., L.K. Harris, and J.C. Cambier. 1986. Anti-Ig induces release of inositol 1,4,5-trisphosphate, which mediates mobilization of intracellular Ca^{++} stores in B lymphocytes. *J. Immunol.* 137:708.
- Bijsterbosch, M.K., C.M. Meade, G.A. Turner, and G.G.B. Klaus. 1985. B lymphocyte receptors and phosphoinositide degradation. *Cell.* 41:999.
- Fahey, K.A., and A.L. DeFranco. 1987. Cross-linking membrane IgM induces production of inositol trisphosphate and inositol tetrakisphosphate in WEHI-231 B lymphoma cells. *J. Immunol.* 138:3835.
- Gold, M.R., D.A. Law, and A.L. DeFranco. 1990. Stimulation of protein tyrosine phosphorylation by the B-lymphocyte antigen receptor. *Nature (Lond.)* 345:810.
- Brunswick, M., L.E. Samelson, and J.J. Mond. 1991. Surface immunoglobulin crosslinking activates a tyrosine kinase pathway in B cells that is independent of protein kinase C. *Proc. Natl. Acad. Sci. USA.* 88:1311.
- Campbell, M.A., and B.M. Sefton. 1990. Protein tyrosine phosphorylation is induced in murine B lymphocytes in response to stimulation with anti-immunoglobulin. *EMBO (Eur. Mol. Biol. Organ.) J.* 9:2125.
- Carter, R.H., D.J. Park, S.G. Rhee, and D.T. Fearon. 1991. Tyrosine phosphorylation of phospholipase C induced by membrane immunoglobulin in B lymphocytes. *Proc. Natl. Acad. Sci. USA.* 88:2745.
- Gold, M.R., L. Matsuuchi, R.B. Kelly, and A.L. DeFranco. 1991. Tyrosine phosphorylation of components of the B-cell antigen receptors following receptor crosslinking. *Proc. Natl. Acad. Sci. USA.* 88:3436.
- Hempel, W.M., R.C. Schatzman, and A.L. DeFranco. 1992. Tyrosine phosphorylation of phospholipase C- γ 2 upon crosslinking of membrane Ig on murine B lymphocytes. *J. Immunol.* 148:3021.
- Brunswick, M., C.H. June, F.D. Finkelman, H.M. Dintzis, J.K. Inman, and J.J. Mond. 1989. Dextran conjugated anti-Ig antibody stimulates B cells by repetitive signal transduction: a model for T cell independent B cell activation. *Proc. Natl. Acad. Sci. USA.* 86:6724.
- Brunswick, M., E. Bonvini, M. Francis, C.C. Felder, T. Hoffman, and J. Mond. 1990. Absence of demonstrable phospholipid turnover in B cells stimulated by low mitogenic concentration of dextran-anti-immunoglobulin conjugates. *Eur. J. Immunol.* 20:855.
- Lindsberg, M.-L., M. Brunswick, H. Yamada, A. Lees, J. Inman, C.H. June, and J.J. Mond. 1991. Biochemical analysis of the immune B cell defect in *xid* mice. *J. Immunol.* 147:3774.
- Francois, D.T., I.M. Katona, C.H. June, L.M. Wahl, N. Feuerstein, K.-P. Huang, and J.J. Mond. 1988. Anti-Ig-mediated proliferation of human B cells in the absence of protein kinase C. *J. Immunol.* 140:3338.
- Mond, J.J., N. Feuerstein, C.H. June, A.K. Balapure, R.I. Glazer, K. Witherspoon, and M. Brunswick. 1990. Bimodal effect of phorbol ester on B cell activation: implication for the role of protein kinase C. *J. Biol. Chem.* 266:4458.
- Padeh, S., A. Levitzki, A. Gazit, G.B. Mills, and C.M. Roifman. 1991. Activation of phospholipase C in human B cells is dependent on tyrosine phosphorylation. *J. Clin. Invest.* 87:1114.
- Mond, J.J., N. Feuerstein, F.D. Finkelman, F. Huang, K.-P. Huang, and G. Dennis. 1987. B-lymphocyte activation mediated by anti-immunoglobulin antibody in the absence of protein kinase C. *Proc. Natl. Acad. Sci. USA.* 84:8588.
- Brunswick, M., F.D. Finkelman, P.F. Highet, J.K. Inman, H.M. Dintzis, and J.J. Mond. 1988. Picogram quantities of anti-Ig antibodies coupled to dextran induce B cell proliferation. *J. Immunol.* 140:3364.
- Wilson, H.A., D. Greenblatt, M. Poenie, F.D. Finkelman, and R.Y. Tsien. 1987. Crosslinkage of B lymphocyte surface immunoglobulin by anti-Ig or antigen induces prolonged oscillations of intracellular ionized calcium. *J. Exp. Med.* 166:601.
- Ledbetter, J.A., and L.A. Herzenberg. 1979. Xenogeneic monoclonal antibodies to mouse lymphoid differentiation antigens. *Immunol. Rev.* 47:63.
- Dialynas, D.P., D.B. Wilde, P. Marrack, A. Pierres, K.A. Wall, W. Havran, G. Otten, M.R. Loken, M. Pierres, J. Kappler, and F.W. Fitch. 1983. Characterization of the murine antigenic determinant, designated L3T4a, recognized by monoclonal antibody GK1.5: expression of L3T4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. *Immunol. Rev.* 74:29.
- Lanier, L.L. 1982. Monoclonal antibodies against rat immunoglobulin kappa chains. *Hybridoma.* 1:125.
- Tsien, R.Y., and A.T. Harootunian. 1990. Practical design criteria for a dynamic ratio imaging system. *Cell Calcium.* 11:93.
- Niklinska, B.J., H. Yamada, J.J. O'Shea, C.H. June, and J.D. Ashwell. 1992. Tyrosine kinase-regulated and inositol phosphate-independent Ca^{2+} elevation and mobilization in T cells. *J. Biol. Chem.* 267:7154.
- DeFranco, A.L., J.T. Kung, and W.E. Paul. 1982. Regulation of growth and proliferation in B cell subpopulations. *Immunol. Rev.* 64:161.
- Goroff, D.K., A. Stall, J.J. Mond, and F.D. Finkelman. 1986. In vitro and in vivo B lymphocyte-activating properties of monoclonal antibodies. I. Determinants of B lymphocyte-activating properties. *J. Immunol.* 136:2382.
- Scher, I. 1982. The CBA/N mouse strain: an experimental model illustrating the influence of the X-chromosome on immunity. *Adv. Immunol.* 33:1.
- Mongini, P.K.A., C.A. Blessinger, and J.P. Dalton. 1991. Affinity requirements for induction of sequential phases of human B cell activation by membrane IgM-cross-linking ligands. *J. Immunol.* 146:1791.
- Goroff, D.K., J.M. Holmes, H. Bazin, F. Nisol, and F.D. Finkelman. 1991. Polyclonal activation of the murine immune system by an antibody to IgD. XI. Contribution of membrane IgD cross-linking to the generation of an in vivo polyclonal antibody response. *J. Immunol.* 146:18.
- Parker, D.C., D.C. Wadsworth, and A.B. Schneider. 1980. Activation of murine B lymphocytes by anti-immunoglobulin is an inductive signal leading to immunoglobulin secretion. *J. Exp. Med.* 152:128.
- Dintzis, R.Z., M. Okajima, M.H. Middleton, G. Greene, and H.M. Dintzis. 1989. The immunization of soluble haptened polymers is determined by molecular mass and hapten valence. *J. Immunol.* 143:1239.
- Chestnut, R.W. and H. Grey. 1981. Studies on the capacity of B cells to serve as antigen-presenting cells. *J. Immunol.* 126:1075.
- Berridge, M.J. 1990. Calcium oscillations. *J. Biol. Chem.* 265:9583.