(c) *Dye tests* have their value in surgical practice. In "medical" cases they yield results comparable to, but not so uniformly reliable as, blood urea estimations.

In conclusion it may be stated that special importance is attached to the following in the diagnosis and prognosis of chronic renal disease: (1) History of the case. (2) General condition of the patient. (3) State of the heart, blood vessels, and blood pressure (systolic and diastolic)—an estimate of cardio-vascular efficiency. (4) Condition of the retinae and discs. (5) General examination of the urine. (6) Range of urea concentration test. (7) Estimation of the blood urea and perhaps also the inorganic phosphorus in the plasma.

Summary.

Special attention has been directed to putting forward the plea that in estimating the function of the kidneys note should be taken not only of the degree to which concentration of waste products in the urine can occur, but also of the capacity of the kidney to produce a dilute urine. It has been shown that the power to concentrate urea is best determined over the night hours after a provocative dose of urea, and that the value of the result obtained is much enhanced by ascertaining the greatest extent to which water diuresis, most effective during the waking hours, can lower the concentration of urea. The minimum concentration provides a basis from which to judge the maximum, and vice versa, an expression of range of function or reserve power. The part played by the heart's action and blood pressure in determining these results has been discussed.

I wish to take this opportunity of expressing my indebtedness to Professor Langmead for his kindness in permitting me to make use of his cases and for much helpful advice.

REFERENCES.

- ¹ Maclean, H.: *Modern Methods in the Diagnosis and Treatment of Renal Disease*, Constable and Co., 1924. ² Cushny, A. R.: *Monograph—The Secretion of Urine*, 1917. ³ Orr and Innes: *Brit. Journ. Exper. Path.*, iii, 61, 1922. ⁴ de Wesselow: *Lancet*, 163, 1923.

THE RELATIVE LOSS OF HEAT AND WEIGHT IN THE NEWBORN, AND THE TREATMENT OF SHOCK.*

BY

A. LOUISE McILROY, M.D., D.Sc.,
PROFESSOR OF OBSTETRICS, UNIVERSITY OF LONDON.
(From the Obstetric Unit, Royal Free Hospital.)

THE RELATIVE LOSS OF HEAT AND WEIGHT IN THE NEWBORN.

The initial drop in the weight of the newborn has hitherto been regarded as inevitable. As a rule the chief fall occurs in the first three days. The average loss is about seven ounces, and some authorities state that half a pound is within normal physiological limits. Various causes are assigned, such as adaptation of ante-natal and post-natal metabolism, loss of urine and meconium, and deficient nutriment until the mother's milk is fully established.

There is no doubt that some loss of weight must occur in the majority of cases, but the problem is whether the loss cannot be diminished. Weight cannot be regarded as an

absolute indication of the infant's welfare, but is one of the best signs of progress.

Some time ago Professor Pembrey pointed out to me that the treatment of the normal newborn was irrational as compared with that of the premature, and deprecated the process of bathing as it entails loss of heat.

Acting on this suggestion, some nine months ago I began to try new methods of treating newborn infants. A series of normal babies at term were cleaned at birth with olive oil instead of by bathing.

The results have been satisfactory, and now oiling has

been instituted as the routine method in the Obstetric Unit at the Royal Free Hospital. The babies are oiled after birth and daily until the day before they leave the hospital, when the mothers are instructed how to bathe them. They lose less weight than bathed babies, and in most cases regain their birth weights or reach even higher weights before leaving hospital on the ninth or tenth day. The cleansing properties of oil are not well understood in this country: babies treated in the manner described are just as sweet and clean as if they had been bathed in the ordinary way. The oil is, of course, wiped off again after application.

After treating a large number of infants I felt that still further improvement could be effected by reducing exposure to the air after birth, and thus avoiding loss of heat. Immediately after delivery the infants in one ward were wrapped in warmed blankets. On the cord being cut the child was at once transferred to a cot with warmed blankets and hot-water bottles, and kept there until oiled and dressed. The results in these cases were still further improved.

Appended are two charts displaying results of 160 normal cases selected at random. For purpose of comparison the charts deal with 40 oiled and warmed babies and 40 bathed (controls) babies of multiparae, and 40 oiled and warmed babies and 40 bathed (controls) babies of primiparae. The oiled and warmed classes on the one hand, and the bathed on the other, correspond as nearly as possible in their birth months. Birth weights correspond as nearly as possible, with a maximum divergence of 4 oz.

The records of the oiled and warmed babies show a marked improvement, proving that loss of heat means a loss of weight in the newborn. Contrary to expectation, there appears to be little difference between first and subsequent babies, but in the oiled and warmed class first-born babies show a slight advantage.

Records of temperature (rectal) were kept of 100 warmed and oiled babies. The temperatures were taken at birth, one or two hours after, and after oiling. In almost every case there was a fall after birth and after oiling. The average temperature at birth was found to be between 99° and 100° F. In one or two hours it had dropped about 1 degree. After oiling it was further reduced 0.5 to 1 degree.

Strain during labour with shock to the infant is followed by a greater fall in weight than occurs after normal delivery. In forceps cases there is a considerable drop in the weight of the infant during the first three days. This is less marked when the child is not exposed to air or bathing. Premature babies and also those born by Caesarean section usually lose more weight than children born by normal deliveries. Early artificial feeding until the mother's milk is established in both these groups of cases has been found to give satisfactory results. The weight charts of twins show in many cases a parallel loss and gain of weight for each infant. In infants of mothers who have suffered from toxæmia the charts show a considerable loss of weight.

Feeding methods are not involved in the inquiry, but better results as to weight are obtained when the child is fed every three hours than when it is fed every four hours. The children were all breast-fed, but when the mother's milk supply was deficient supplementary feeds were given.

The decrease in infantile mortality during the first year has aroused much attention, but few realize that the decrease during the first month is comparatively small. To put the matter in another way, the improvement has been chiefly during the last eleven months of the infantile year. For instance, in 1905 the mortality per 1,000 for the first year was 128, between two and eleven months 86, and during the first month 42. In 1923 the corresponding

FIG. 2.—Charts showing the average loss of weight in 40 oiled and warmed babies and 40 bathed babies of multiparae, and 40 oiled and warmed babies and 40 bathed babies of primiparae (1 square = 1 oz.).

FIG. 1.—Charts showing initial loss of weight in normal newborn babies: A, of multipara (bathed); B, of primipara (warmed).

* Read at a meeting of the Obstetrical Section of the Royal Society of Medicine, December 4th, 1924.

figures were 69, 37.5, and 31.5. According to the Registrar-General's records, deaths during the first month are mainly due to birth injuries and prematurity.

Probably much can be done to reduce mortality by improved methods at and after birth, but ante-natal treatment is the real remedy. The future of successful obstetric practice lies in skilled supervision during the ante-natal period. It is obvious that an infant whose mother has suffered during pregnancy from toxæmias or any constitutional disease has a reduced vital capacity. Any form of ante-natal treatment, therefore, which improves the mother's general condition and reduces the possible complications of labour will have a marked effect upon the newborn child.

Ante-natal correction of malpresentations, the treatment of cardiac complications by rest, the reduction of pain by the administration of sedatives during labour, and the stimulation of a failing foetal heart by strychnine, strophanthus, etc., must have a beneficial effect upon the welfare of the infant.

SHOCK IN THE NEWBORN.

The treatment of white asphyxia demands fresh consideration. As a rule blue asphyxia can be treated by clearing out the air passages and stimulating the skin reflexes.

In white asphyxia the usual method is to clear the air passages and immerse the infant in a warm bath, sometimes with the application of cold water to the head, followed by artificial respiration of varying and progressive degrees of severity, finally culminating in the Schultze method. This arduous effort probably satisfies the obstetrician that if he has failed to restore the infant he cannot be blamed for not having exercised his energies to their utmost capacity.

The late Dr. Gordon Ley pointed out that in such cases attention should be directed, not to establishing respiratory movements, but to relieving cardiac failure due to shock, usually caused by cranial compression, or in some cases by pressure upon the umbilical circulation. This is, I think, the correct view. The greater the strain upon the mother during labour the greater the shock to the infant. Respiration cannot be induced until the heart has adapted itself. What is required is cardiac stimulation and not inhibition. Much depends upon the amount of carbon dioxide in the blood for the stimulation of the respiratory centres. It is possible that the reserves for this purpose may be wasted in useless muscular movements carried out upon an infant with an enfeebled heart.

As soon as the child is born it should be wrapped in a warm blanket and its air passages cleared. The mucus should be sucked out of the trachea by a rubber catheter with a glass window enabling the mucus to be observed before it reaches the operator's mouth. The infant should be then left at rest in a warm cot. Gentle massage of the heart can be carried out under the blanket, and stimulants such as pituitrin (2 minims) and camphor given hypodermically. No movements should be permitted which involve the head or spine. It is difficult at first to abstain from artificial respiration; but experience has proved the efficiency of the alternative method, and shown that results are better than when more vigorous means are employed. Time, warmth, and rest will usually restore the heart's action if the shock has not been so severe as to be incompatible with life. In cases of blocked air passages artificial respiration may be advocated, but the use of the mucus catheter will generally be successful. Heart failure is not usually the cause of death in such instances.

After the administration of scopolamine, or morphine, or even of ether, during labour, the infant is often born in an anaesthetized condition. It must therefore be allowed time to recover.

I am grateful to my obstetric residents, Miss Langston and Miss Salmond, and to the students and sisters for assisting me in these investigations. A record of several hundred cases has been charted and classified by Miss Pritchard, whose services were available through the assistance of the Medical Research Council.

EPIDEMIC ENCEPHALITIS SIMULATING ACUTE SURGICAL LESIONS OF THE ABDOMEN.

BY

G. H. STEVENSON, O.B.E., M.C., F.R.C.S.,
M.R.C.P.Ed., D.P.H.,

SURGEON TO OUT-PATIENTS, GLASGOW ROYAL INFIRMARY; SURGEON,
MINISTRY OF PENSIONS HOSPITAL, GLASGOW.

I AM taking the opportunity of reporting shortly on four cases observed by me during August, 1924, to put on record what appears to be a new development relating to the diagnosis of that extraordinary disease now so well known as "lethargic" encephalitis.

At present we are experiencing an epidemic of this disease in Glasgow, and it will be clear from my report how essential it is for those of us who are engaged in hospital practice to draw the attention of our colleagues engaged in general practice to the various forms which encephalitis may take. I may add that the diagnosis of the disease under the conditions of general practice met with in this city is of necessity a matter of the greatest difficulty, especially as many of the doctors will have had little opportunity of seeing cases formerly. It would seem that the qualifying adjective "lethargic" would hardly apply to the type of the disease I have seen lately, and the term certainly conveys a very wrong impression of the striking symptoms, at any rate in the earlier stages of the disease. It is pretty certain that many of the milder cases are treated for other conditions, and the danger of the disease becoming more widespread is very real.

All four cases described were admitted to hospital during the short period of one month (August 13th to August 28th as a matter of fact) as *acute abdominal* conditions, and subsequently three out of the four were diagnosed definitely as epidemic encephalitis and removed to isolation hospitals. The fourth case is as yet not finally diagnosed, but the diagnosis appears to rest between encephalitis and tuberculous meningitis. This case shows that even when the resources of a large general hospital are available diagnosis may be difficult. The notes on the cases are of necessity short as the cases were observed for a brief time only.

CASE I.

Serving soldier, aged 19. Admitted August 15th, 1924. The diagnosis on admission was *perforated gastric ulcer*, which had been changed from that of *strangulated hernia*. The patient gave a history of sudden onset of very severe pain in the left lower quadrant of the abdomen, and stated that he had received a blow in this region on August 13th. He had vomited freely prior to admission.

I did not see the patient until some hours after the diagnosis of perforated gastric ulcer had been made, and it was then quite clear that we were dealing with some acute nervous condition. My first impression was that the case was one of tetanus or strychnine poisoning, as on entering the door of the side-room in which he lay I found him in the position of opisthotonos and having marked spasm of almost all the muscles of the trunk and limbs. His mental condition was, however, lethargic, and in complete contrast to the spastic state of his body. Hyperaesthesia of the skin of the right side of the abdomen from nipple to groin was very marked, and on testing for this strong myoclonic movements were produced. The temperature was 97° F. and the pulse rate 60. He was unable to give a history of any eye symptoms. Having excluded the possibility of tetanus and strychnine poisoning so far as possible I made the provisional diagnosis of epidemic encephalitis, which was confirmed by a physician (Dr. Harrington) next morning. I was assisted in the diagnosis of this case by the fact that I had seen an almost similar case in the same room about a year previously. In this case, however, the pain started in the left arm, but was accompanied by similar generalized myoclonic movements.

The patient was removed in a few days, having run a completely apyrexial course with rather a slow pulse. Ophthalmic examination before removal showed oedema of both optic discs.

It is obvious in this case that the doctor who saw the case prior to admission had seen him before the onset of the typical myoclonic movements, which appear to follow and not precede the agonizing pain.

CASE II.

Miner, aged 41. Admitted August 20th, 1924. The diagnosis on admission was *right-sided renal colic*. The patient gave a history of having been seized with acute pain in the right lumbar region, which later shot down into the right testicle, simulating the pain of the passage of a stone. He had, however, no haematuria and no difficulty in passing urine.