I have tried to show how far the complications due to malpresentations and misfits, to ante-partum haemorrhage and convulsions, with consequent interference in labour, may be greatly lessened. Errors due to miscalculations will be with us so long as we remain erring mortals, but as we become more experienced they will become fewer. There will, however, be left over four-fifths of the interference in labour—the low forceps delivery, the “shoe-horn” operation of easing the head over the perineum because of failure of the forces of labour to overcome the final obstacle of the resistance of the levator ani muscles. Happily this procedure is a minor one, but even it calls for a fresh attitude of mind in regard to it, though that is another story which I cannot begin to-night. The forceps must be regarded as our second line of defence after our first line has given way, and we must try to strengthen the first line, which is to seek out the causes of the ineffectiveness of the powers and remove them—of which fatigue, long-continued pain, anxiety, and dread of the ordeal are the most prominent.

Lectures

ON

THE SYMPATHETIC INNERVATION OF STRIATED MUSCLE.*

BY

THE LATE JOHN IRVINE HUNTER, M.D., CH.M.,
CHALLIS PROFESSOR OF ANATOMY IN THE UNIVERSITY OF SYDNEY.

LECTURE II.—PART II.

THE FUNCTIONS OF THE TWO GROUPS OF MUSCLE FIBRES:

EXPERIMENTAL EVIDENCE (Continued).

EXPERIMENTS ON BIRDS.

THE folded attitude of the wings of birds when standing or walking affords an excellent example of what Sherrington has called an “active posture”—namely, the maintenance of position against the action of gravity by means of the activity of contractile tissues. I thought a study of the mechanism concerned in the production and maintenance of this posture would throw light on the postural activity of voluntary muscle generally, and provide a means of determining the exact part played by the dual nerve supply—somatic and sympathetic—of such muscle in the production and maintenance of posture.

The effect of gravity, which would be freely acting under normal conditions in some birds, would tend to cause drooping and falling outwards of one wing, if the mechanism for the maintenance of its posture were interfered with, and by comparison with the position of the normal wing of the opposite side the degree of interference is readily measurable. The wing offers advantages for study over the limbs of the cat, dog, and goat, in which the normal attitude is extension, for compensatory efforts are necessary in order to retain this position in the interests of maintaining the erect attitude. This tends to mask the effects of injury reducing the efficiency of the postural mechanism. Variations in the position of the wing may also be studied more easily than alterations in the degrees of flexion of the hind limbs of the frog after sympathetic ramisection, as suspension of the frog (which is necessary to allow the normal side and the injured side to be compared) introduces rotatory reflexes. This circumstance renders a careful analysis necessary in order to dissociate the effects of labyrinthine impulses from the proprioceptive effects produced in the musculature of the limb itself¹⁵ (cf. Langelaan, *Brain*, 1915 and 1922).

In his memoir, “The Postural Activity of Muscle and Nerve,”¹⁶ Sherrington says:

“Section of the afferent roots of the wing affects little, if at all, the folded posture maintained by the wing when not in flight, i.e., during standing or walking. Nor has the source of the postural

contraction of the flexors of the wing been found. Here, as in the case of the iris, the postural contraction, if, as is presumable, reflex, lies in receptors not those of the contracting muscle themselves—is, in short, alienogenous not autogenous.”

In view of our experiments on goats it appeared to me that a possible explanation for the failure to alter materially the folded position of the wing, by severing the posterior nerve roots corresponding to the origin of the brachial plexus, was due to the fact that in such an experiment the sympathetic innervation of the voluntary muscle of the wing was left intact. In previous papers (see, for example, *Surgery, Gynecology and Obstetrics*, December, 1924) I have argued that the function of the sympathetic nerve supply of skeletal muscle is the maintenance of posture once the position is attained—in other words, that it is responsible for the property of plastic tone. It appeared desirable, therefore, to ascertain the effect of removal of the influence of the sympathetic nerve supply of the muscle of the wing by section of the cervical sympathetic. The result of the first experiment of this kind has already been recorded.¹⁷ The wing was markedly drooped compared with the normal side when the body of the fowl was supported in the hand during recovery from the anaesthetic. Later the droop of this wing was definitely noticeable but not so marked, its tip being 1.5 cm. lower than the tip of the intact wing. The wing also was less firmly folded to the side of the body. This condition was still noticeable when the fowl was kept at rest for some minutes up to the time of being killed seventy-four days later. A *post-mortem* examination revealed that the vertebral sympathetic cord had been cut below (or behind) the upper two roots of the brachial plexus only, so that the effect was incomplete. Subsequent operations of a more complete character, in which the sympathetic was approached by dissection of the intertransverse spaces from behind, instead of from the front as in the first experiment, gave similar results.

Anatomical Features of the Cervical Sympathetic Trunk of Birds.

The vertebral sympathetic plexus or cord, which contains the fibres that interest us in these experiments, passes through the foramina of the transverse processes along the vertebral vessels on a plane ventral to the roots of the brachial plexus. On approaching the ribs the cord usually splits into dorsal and ventral trunks, which pass respectively dorsal and ventral to the neck of the rib. The operation usually performed on the fowls was to divide these trunks above the first true rib (at C in Fig. 7)—that is, caudal to the last large nerve (the last cervical) which contributes towards the formation of the brachial plexus. In sea-gulls I invariably cut the dorsal and ventral trunks immediately behind (or below) the penultimate cervical nerve as well.

This section produced a more marked effect on the eyelids than the more caudal section. Langley has shown that the last cervical nerve is the first nerve through which pre-ganglionic fibres of the sympathetic nervous system emerge from the spinal cord. He records that the criterion of movement of the feathers following stimulation shows that these fibres are distributed to the upper part of the neck and the head. It may be distributed to the anterior part of the wing, but the effect is only slight when it occurs there. Therefore the more caudal section of the sympathetic

FIG. 7.—Diagram of the fowl's brachial plexus. C, The spot where the whole sympathetic connexions of the wing can be severed. D, The line along which either the posterior or the anterior somatic roots or both can be cut.

* Delivered by Professor Elliot Smith at University College, January 26th, 1925.

chain satisfies the purpose of the present investigation. But the changes in the position of the eyelid are useful as indications that the chain has been successfully identified and cut. In the fowl the effect of section of the cervical sympathetic above the last cervical nerve (at *C* in Fig. 7) produces a raising of the lower eyelid and a more marked drooping and overhanging of the upper lid, so that the pupil is partially obscured. Normally the upper lid is retracted to such an extent that only its free edge is visible. After the operation, especially when viewed from above, it may be seen for an extent of several millimetres. When the effect is not so apparent as in the lower section alone a useful test is to draw the upper lid down over the eye on each side and compare the rate of its return on the two sides. It returns perceptibly more slowly on the operated side, and the return is, of course, not complete. The effects on the eyelid of the sea-gull are not so noticeable as in the fowl because they are smaller and less mobile. But the circular palpebral tissue becomes oval in shape following the operation.

The removal of the cervical sympathetic cord has an effect upon the pupil in the bird, as Langley has shown. The test of ruffling the feathers described by Langley is useful to determine whether the operation has been successful. On the normal side the feathers rapidly return to their normal depressed condition, whereas on the side from which the sympathetic innervation has been removed the return is slow.

In this series of experiments the effect upon the blood vessels was not used as a criterion to indicate whether the sympathetic had been severed or not. In one case a marked increase of temperature of the comb occurred on the side of the operation. In another marked pallor was observed in the vascular appendages of the head following the operation. This was probably due to a state of strong contraction of the capillaries following sympathetic denervation described by Krogh. On examination on the following day in the above instance it was again present.

The Effect of Section of the Cervical Sympathetic upon the Posture of the Wing.

The immediate effects upon the position of the wing of section of the cervical sympathetic cord have already been briefly noted. In the fowl the wing is somewhat drooped on the affected side. This persists after healing has completely occurred. For instance, in one of the fowls the distance of the tip of the wing from the mid-line, as measured by the callipers on the eleventh day following the operation on the left side, was 4.2 cm. on this side, compared with the corresponding measurement of 3 cm. on the normal side. When the wings were stretched outwards and then simultaneously released, an average of three measurements on the left and right sides were 4.2 cm. and 2.2 cm. respectively. These differences are in part due to the falling away of the limb from the side (abduction) in addition to dropping downwards. An estimate of the difference due to these causes is also obtainable by measuring the distance of the outer surface of the wing at the level of the elbow-joint from the mid-line on each side. On the day in question the left and right measurements were 10.1 cm. and 8 cm. respectively.

Another change due to the operation is the slowness in the return of the partially denervated limb to its final position when released after stretching outward or downward or after elevation. This effect is perceptible to the eye and is constant over numerous trials. Cinematographic records confirmed this observation. The greater part of the return movement takes place relatively rapidly, though not so rapidly as on the normal side. The limb remains somewhat abducted and elevated, and then slowly settles down to its final position, which is closer to the trunk and in a somewhat more depressed position. This latter element in the return movement may occupy several seconds. If the wing is passively placed in its normal position it rapidly falls to the depressed level. Sometimes the bird itself voluntarily lifts the wing to its normal position, but it quickly falls to its original position. This effect becomes more marked in fatigue—for example, after repeated examination.

In sea-gulls abduction and dropping of the elbow is apparent, but the effect is not so marked as in the fowl, because the tail is normally used to support the large wings, which in some of the specimens operated upon attain a span of a metre from tip to tip. In the normal position of the wings they lie crossed over one another resting on the tail. Subsequent to cervical sympathetic section the affected wing takes a position nearly parallel with the direction of the tail, and, following a two-sided operation, an interval may exist between the two wings which is not normally to be seen. When an affected wing is removed from the support of the tail it droops below the tail level approximately to the extent of 2.5 cm. The drooping of the wing is very marked during the process of recovery of the limb from the anaesthetic. Open ether was the anaesthetic employed in each experiment.

For purposes of control the vertebro-scapular muscles were severed on both sides, and both right and left sympathetic cords were exposed, but only the left was cut. The only difference between the two sides, therefore, is that section of the sympathetic cord was performed on the left side and not on the right side.

It is apparent from these remarks that the tone of the affected wing is reduced so that the normal posture is not perfectly maintained. But considerable tone remains, and it tends to impose a folded posture upon the wing. This influence must be due to the presence of the somatic proprioceptive reflex arcs which remain intact, but in the absence of the influence of the sympathetic innervation of the wing the maximum position of abduction and flexion can no longer be maintained against the action of gravity. This force leads to some degree of drooping and abduction of the wing. During light anaesthesia the effect is rendered more evident as the result of the partial removal of the influence of the somatic proprioceptive connexions. The influence of the somatic arc is also markedly reduced by fatigue, as already noticed following repeated stretching of the wing in the fowl. The natural effort of sustaining flight shows the result of fatigue to greater advantage. These observations were carried out with the bird attached to a light line mounted on a smoothly running reel. In the unilateral operation the position of the normal wing may be used as a control, as in sea-gull 3. Before flight only slight abduction and droop of the elbow region of the wing were apparent. After a flight of one hundred metres, in which the wings were maintained in constant movement so that no gliding occurred, the abduction and drooping became very marked. The wing no longer rested on the tail, and the bird made ineffectual attempts to place it in its usual position. On supporting the body of the bird in the hand and allowing the wings to drop freely a condition resembling that seen in similar circumstances during recovery from the anaesthetic resulted. Fatigue of the somatically innervated muscles has apparently produced this effect, so that the condition of the wing in such circumstances is not far removed from that seen in flaccid paralysis.

These observations are in accord with the results of our operations on goats—namely, that the sympathetic innervation of voluntary muscles is responsible for the maintenance of plastic tone. The defect following removal of the sympathetic influence consists in inability of the voluntary muscle to remain fixed at a given length—such, for instance, as the length required to maintain the extended position of the limbs in a quadruped when placed on its back, the flexed position of the frog's hind limbs, or the folded position of the wing normally assumed when a bird is walking or standing. The flexed posture of the wing, to take the latter instance, is conferred upon the wing by the intact somatic proprioceptive reflex arc; and this tendency disappears when this arc is interrupted. Therefore this arc, when intact, subserves contractile tone (Langelaan, 1922). The position of flexion of the wing imposed by the contractile fibres fails to be maintained to the full degree in the absence of plastic tone, which is subserved by the sympathetically innervated fibres of the striated muscle of the wing. Normally, therefore, plastic tone is the property of muscle enabling the flexed position to be maintained. The "catch" or "ratchet" mechanism of involuntary muscle described by Uexküll is beautifully illustrated. The fixed

position of flexion of the wing can no longer be maintained for an indefinite period with a minimum of fatigue in the absence of the sympathetic arc. Furthermore, after voluntary activity of such partially denervated muscle, of which flight provides the best example, exhaustion of the somatic reflex mechanism manifests itself to a striking degree. It is evident, therefore, that the sympathetic innervation plays a part in movement as well as at rest. To retain the analogy already mentioned, it would appear that an "infinitely fine ratchet" assists in voluntary movement. This helps to maintain the intermediate postures through which the muscle passes during the execution of an active movement.

In the absence of the sympathetic nerve supply it is apparent that greater work is imposed upon the somatic nervous system during voluntary activity, for it not only performs the voluntary movement, but alone supports the weight of the limb. When both wings have been partially denervated by removal of the sympathetic nerve supply (as in sea-gull 8) after several short flights both wings were drooped and the bird was apparently exhausted, the mouth was widely open, and respiration very rapid. On being approached the bird only attempted short low flights to avoid recapture.

The effect of the section of the cervical sympathetic trunk in narrowing the palpebral fissure is to be ascribed to loss of plastic tone of the striated muscle of the eyelid to a greater degree than to the denervation of involuntary muscle. Dr. Oliver Latham of Sydney has made gold chloride preparations which show the sympathetic non-medullated fibres terminating in characteristic grape-like endings on considerable bundles of the eyelids. Removal of the influence of these fibres affects the capability of the muscle to remain fixed at the length required to maintain the eyelids widely separated.

Section of the Afferent Limb of the Sympathetic Proprioceptive Reflex Arc.

By cutting the posterior nerve roots (see the line *D* in Fig. 7) Sherrington conclusively proved that tone is subserved by a proprioceptive reflex mechanism. The question arises, then, as to whether a sympathetic proprioceptive reflex arc is responsible for the production and maintenance of plastic tone. Langley showed that the feathers of the wing are influenced by stimulation of the last cervical nerve and the upper two thoracic nerves. Of these the first thoracic nerve has a strong effect on the wing. The anterior part of the wing may be affected by the last cervical nerve, but this effect when present is only slight. The second thoracic nerve has a moderate or slight effect, usually on the posterior part of the wing only. Acting on the supposition that the sympathetic innervation of the voluntary muscle of the wing would have the same origin as that of the feathers, an operation was performed on the first thoracic posterior nerve root. This may be done by enlarging the first foramen seen on the dorsal view of the fused thoracic vertebrae. *Post-mortem* examination showed that this root was completely severed without injury to the anterior nerve root. In this case the brachial plexus received no somatic contributions from the first thoracic nerve. The large size of this root in all the specimens operated on is a striking feature in view of the slight part, if any, this segment takes in the formation of the brachial plexus. This confirms the suggestion that this nerve contributes the principal sympathetic connexions to the wing, so accounting for its large size. The effect of removal of the posterior root of this nerve upon the posture of the wing is in all respects the same as that following section of the cervical sympathetic cord caudal to the brachial plexus. Drooping and abduction of the wing are apparent; and, on releasing the wings after stretching, the affected side was perceptibly slower in returning to its final position. On the fifth day the measurements were as follows: The tip of the wing was 2.5 cm. from the mid-line on the left side, and 0 cm. on the right side; or, measured by the callipers, the elbows were 80 cm. and 6.8 cm. from the mid-line respectively. When the wings were stretched outward and then released the tips were 2.7 cm. and 2 cm. from the mid-line on the left

and right sides respectively. It is evident, therefore, that if a sympathetic proprioceptive arc subserves plastic tonus afferent fibres must descend along the ganglionated (vertebral) chain to the first thoracic level and there pass to the nerve trunk at this level by way of one or other or both rami communicantes. In the experiment of section of the cervical sympathetic posterior to the brachial plexus both afferent and efferent fibres are therefore severed. The same effect follows section of the afferent limb of the arc only when the first thoracic posterior nerve root is cut through. Sherrington's preparations, in which the afferent nerves were cut, owe some of their features to removal of the influence of the sympathetic proprioceptive reflex arc subserving plastic tone, for in "deafferenting" the fore limb of the cat he removed *inter alia* the posterior nerve root of the first thoracic nerve, and in the case of the hind limb of that animal the fourth lumbar posterior nerve root was one of the two roots resected. In the cat the thoraco-lumbar outflow extends from the first thoracic to the fourth lumbar segment: hence Sherrington's operations included the region of the sympathetic innervation of the limbs.

Persistence of Plastic Tonus after Removal of all the Posterior Nerve Roots Connected with the Brachial Plexus.

The anatomical connexions of the somatic and sympathetic nervous systems and the wing musculature make it possible to remove the influence of the former without materially affecting the innervation derived from the latter source: for the most important sympathetic connexion consists of a reflex arc at the first thoracic level. On one side the posterior nerve roots of four segments above this level were removed with a view to removing the influence of the somatic reflex arcs. In addition to the absence of voluntary movement the result of this operation is to remove the tendency of the wing to take up a flexed position—that is, contractile tone is absent. When the wing is stretched outwards, therefore, and released, it drops to the side of the bird, and instead of taking up the flexed attitude characteristic of the normal wing it remains dependent. If the wing is passively flexed it will remain in this new position. In this way the wing may be moulded into the position normally occupied by it, and it will remain there indefinitely.

The wing may now be drawn down to a dependent position, where it will remain. It has, therefore, the properties of a plastic body, enabling it to retain any position in which it may be placed. In other words, the flexors of the wing exhibit the "shortening and lengthening reactions." These reactions indicate the presence of plastic tone.

It may be inferred, therefore, that contractile tone, which normally tends to impose the typical posture upon the wing, is absent in such a preparation, and is therefore a function of the somatic proprioceptive reflex arc that has been destroyed. Plastic tone, which fixes the length of the muscle fibres to maintain a posture once assumed, is a function of the sympathetic proprioceptive reflex arc that remains intact. Together these components of tone are responsible for posture, so that the combination of the two may be called "postural tone."

In one fowl in which the posterior roots of the brachial plexus had been severed, the relationship between plastic tone and the sympathetic innervation of the wing musculature was shown by a further experiment. The cervical sympathetic was severed immediately below (or behind) the last cervical nerve, directly after an examination that revealed the presence of plastic tone in the wing. After the operation the plastic tone had disappeared. The wing was heavy and dependent as in flaccid paralysis. It would not remain in any position of flexion in which it was placed. When folded over the back of the bird it dropped heavily to the ground. This experiment proves conclusively that the plastic tone exhibited after the first operation was maintained by the sympathetic proprioceptive reflex connexions of the wing musculature. As other experiments have shown, this is at the level of the first thoracic segment in the fowl.

*Compensatory Increase in Contractile Tone after
Sympathetic Denervation of the Skeletal
Muscle of the Wing.*

To test the duration of the effect upon the wing of severing the cervical sympathetic trunk caudal to the brachial plexus, the position of the normal and affected wings of some preparations were daily measured. I am indebted to Mr. R. E. Murray, who carried out these daily observations. Fig. 8 is a graphic record of these observations. One observation was made each day on the position of the tips of the wings from the median line was recorded daily. Both graphs show the same features. The contractile tone increases in the affected wing until both wings are approximately equidistant from the median line. This condition is gradually attained and the distances become equal in about sixty days. A similar result is found after removal of the afferent limb of the sympathetic proprioceptive reflex arc.

When this compensation has taken place the effect of cutting the cervical sympathetic is manifested under two conditions; when the bird is kept at rest for a period of approximately fifteen minutes or longer, the droop and abduction of the operated side reappear. Fatigue has a

similar effect. Starting with the elbows and tips of the two wings equidistant from the median line, the left wing and then the right wing were each folded across the back by passive movement fifty times. The difference in the distance of the elbows from the mid-line was then 2.2 cm. and the difference in the distance of the tips from this line was 2.9 cm., due to the droop of the wing on the operated side.

Plastic tone was therefore still absent. It remains to account for the increase in contractile tone. Magnus and de Kleijn have demonstrated that pathways other than the afferent nerves from the muscles themselves affect the tone of skeletal muscle. These accessory influences are the labyrinths and the proprioceptors of the neck muscles. In the resting conditions both these sources of impulses would no longer be effective, and consequently the contractile tone would be lessened, as in the experiment recorded above. It is noteworthy that Langelaan has shown that in frogs removal of

FIG. 8.—One of Mr. R. E. Murray's graphic records to express the compensatory increase in contractile tone in a fowl's wing after sympathetic denervation.

plastic tone from the hind limb leads to domination of the labyrinth of the opposite side. This leads to extension of the normal limb and increased flexion of the limb upon the operated side.

REFERENCES.

- ¹⁵ Compare Langelaan, *Brain*, 1915 and 1922. ¹⁶ *Ibid.*, 1915. ¹⁷ *Med. Journ. of Australia*, June 14th, 1924, p. 581.

THE HYPOTONIC (FLABBY) CHILD.

BY

DONALD PATERSON, M.B., M.R.C.P.,

PHYSICIAN FOR DISEASES OF CHILDREN, WESTMINSTER HOSPITAL;
PHYSICIAN TO OUT-PATIENTS, HOSPITAL FOR SICK CHILDREN,
GREAT ORMOND STREET; PHYSICIAN TO THE INFANTS
HOSPITAL, WESTMINSTER.

A CONSIDERABLE proportion of the children seen at the out-patient department of a children's hospital present such indefinite symptoms but such a uniform clinical picture that they were thought to be worth investigating in detail. These children are generally between the ages of 2 and 7 years, and are brought to hospital for such complaints as tiredness, lack of energy, pains in the limbs on continued exercise, knock-knee, flat-feet, or turning in of the toes when walking, and hence may be seen by either the general physician or the orthopaedic surgeon. Sometimes, however, the complaint was merely that the child had no appetite or was pale, and the usual diagnosis was "debility."

On examination of these children the most striking feature is their stance and their extreme lack of muscle tone. This is well shown in the figures. The large "pot" belly with compensating lordosis is the result of the extremely feeble development of the abdominal muscles. Hyper-extensible joints, knock-knee, flat-feet, turned-in toes, and winged scapulae are further expressions of this same lack of tone. The great majority of cases appear well nourished, but the limb muscles are so soft that it is sometimes difficult to judge whether they are really properly developed. It is interesting to note here that the calf and thigh muscles are always better

developed and have better tone than the arm muscles. Many have enlarged tonsils and adenoids, and some have a more generalized lymphoid enlargement.

Cameron¹ recently described a type of case frequently seen at the out-patient departments, which he terms the nervous child. This is not, however, the type of patient which the writer is attempting to describe and elucidate. The child which Cameron describes is as a rule an only child or the youngest child of elderly parents. He is thin, dislikes to hold himself in the upright position, and has what is

described in America as the "fatigue" stance. His day is spent in bursts of activity and periods of collapse. During a period of exhaustion his stance is like those under investigation, but his muscles never at any time show the hypotonia of a flabby child. The metabolism of the "nervous exhaustion" child is as a rule overactive, and what is required for him is mental and physical rest.

With the hope of elucidating the etiology of this clinical syndrome of flabbiness an investigation of a large number of cases was undertaken during the past year at the Hospital for Sick Children, Great Ormond Street. Early in the investigation it was clear that a number of factors were involved, and the following is a review of these factors and of the various theories that have been put forward to account for the condition.

DIET.

Recently Mellanby² has stated that by diet alone in dogs he can produce every degree of muscular weakness and wasting and can cure the same by restriction of cereals and