found in the spectra which prove beyond doubt that normal and pathological urobilin differ in certain particulars. The band of pathological urobilin at F is dark, broad, and shades off towards violet, while that of normal is narrow, is not deeply shaded, and does not shade off towards violet.

In acid solutions of pathological urobilin other bands are seen, which are missing in the corresponding solutions of the normal pigments.⁶⁴ The idea occurred to me in studying the spectrum produced by the action of nitric acid on solution of bile-pigment, that the stage in which the band at F is dark corresponds to the production of pathological urobilin, while that in which the band at F is faint corresponds to the production of the normal pigment. So it has turned out, but we are not to conclude that either of these pigments as it occurs in urine is necessarily derived from bile pigments. By the long continued action of sodium amalgam and of zinc and sulphuric acid, aided by heat on hæmatin, a substance is eventually formed which possesses many of the characters of (pathological) urobilin.⁶⁵ And by the action of hydrogen peroxide on hæmatin in acid solutions, another substance is found which possesses the character of normal urobilin, as I have proved, and by reducing this with sodium amalgam it changes into a substance resembling pathological urobilin.⁶⁶

For this and other reasons we may conclude that normal urobilin is a more completely oxidised substance than the pathological pigment, but it is possible—and, indeed, probable—that both are produced under different conditions of metabolism. In this they afford a parallel to urea and uric acid; we know that uric acid is a less completely oxidised substance than urea, yet uric acid is not produced by the same metabolism that produces urea. I believe that normal urobilin may be produced elsewhere than in the liver by the action of nascent oxygen on worn-out hæmoglobin and its decomposition products, and in many cases it does not originate from bile pigments. Pathological urobilin may result from the incomplete oxidation of worn-out hæmoglobin and its decomposition products, and may not in many cases have its origin from bile pigment. It would appear to be excreted, as is now well known, after extravasations of blood into the tissues, into the peritoneum, and elsewhere; in such cases, there is too much material for complete oxidation to take place. On the other hand, there is a pigment present in fæces—stercobilin—and this presents nearly all the characters of pathological urobilin,⁶⁷ so that in some cases the presence of the latter in the urine may indicate the absorption of fæcal matters from the intestine. I am convinced that some so-called bilious attacks are due to this cause, for if the pigment is absorbed it must be accompanied by ptomaines, which produce the headache, aching of the limbs, vomiting, and so on. Whether this absorption takes place owing to the liver not performing its guardian function owing to vasomotor disturbance, or whether absorption takes place directly from the intestine, is not clear at present. At all events, everyone knows the beneficial effects of a saline purge in such cases.

In spite of the assertion that the urobilin found in urine is always due to the absorption of hydrobilirubin from the intestine, this hydrobilirubin being supposed to originate from the action of nascent hydrogen, generated by putrefactive processes in the intestine, on the pigments of the bile (Maly and Hoppe-Seyler), I say, without fear of contradiction, that the former may arise in other ways, and that the urobilin of normal urine is a product of oxidation, and that pathological urobilin is an incompletely oxidised substance. By the action of zinc chloride, of that and ammonia, and of other reagents on those pigments, it is easy to prove, as I have elsewhere shown,⁶⁸ that hydrobilirubin is not identical with either normal or pathological urobilin. The presence in the urine of urohæmatoporphyrin and the replacement of this by pathological urobilin, and *vice versa*, teach, when taken in connection with other facts, that the latter may, as I said before, be derived in many cases from hæmatin, and may have no biliary origin. In some cases, of course, we may have urobilin derived from bile pigment as well as directly from hæmatin, but

in such cases there is generally evidence of the passage of bile pigment into the blood. It is a well-known fact that in recovery from jaundice the bilirubin may appear in the urine as urobilin.

The spectroscope is helpful, also, in enabling us to detect indican. If the urine, after applying Jaffe's test, is shaken with chloroform, the latter, if indican is present, will show a band before D due to indigo blue, and another after D, said to be due to indigo red. Or even when Jaffe's test fails, it may be detected by boiling the urine with common hydrochloric acid, cooling and agitating with chloroform; the latter shows the same bands as in Jaffe's test.⁶⁹

There are different pigments which become red on treating the urine with nitric acid and other acids, either at the ordinary temperature or after heating, but at present their origin is uncertain, although at least one is derived from a skatol compound.

Other Pigments.—I have now described some of the most important absorption spectra, speaking from a purely medical point of view, but the subject is far from being exhausted. I have not mentioned the lipoderms or fat pigments, which term includes under it the pigments formerly known as luteins, nor the black pigments known as melanines, some of which latter do not owe their origin to hæmoglobin, as is commonly supposed; nor have I been able to furnish more than a partial and unsatisfactory account of the other pigments. Still what I have said will serve to show that the spectroscope has taught us a good deal, and, I believe, will teach us more when it comes to be more generally used. In biology generally it has recently brought many new facts to light; these in turn must help to advance comparative physiology, which at present is not advanced enough to enable anyone to write even an elementary book upon it.

ESTIMATION OF URIC ACID BY HAYCRAFT'S PROCESS.

Read in the Section of Pathology at the Annual Meeting of the British Medical Association, held at Birmingham, 1890.

By A. HAIG, M.D.,

Assistant Physician Metropolitan Hospital.

THE process consists in (1) combining all the uric acid in a given amount (25 cubic centimetres) of urine with silver contained in a solution of nitrate of silver with addition of ammonia. The result is a gelatinous precipitate; (2) the gelatinous precipitate is collected and well washed on an asbestos filter; (3) the precipitate, after washing, is dissolved in nitric acid; (4) the silver thus got into solution is estimated by means of a colour test (Volhard's method). From the silver found the amount of uric acid combined with it is calculated.

Apparatus.—Flasks, pipettes, measuring-glasses, etc., a flask fitted with a rubber bung perforated in two places. Into one of the perforations fits the stem of a funnel containing broken glass and asbestos to form a filter. Into the other perforation fits a glass tube connected with a Sprengel's pump or other means of producing a vacuum.

[The working of the process was then shown on a specimen of urine.]

Addition of bicarbonate of soda—Haycraft says about 15 grains. Hermann considers that it is better to add about 30 grains; the object of the addition of the bicarbonate is to prevent the reduction of the silver combined with the uric acid.

It is always well to add an excess of the silver nitrate solution, as it is probable that when this is done the resulting urate of silver has a tolerably constant composition. When the patient is taking an iodide a precipitate of iodide of silver falls on adding the silver solution to the urine. This does not necessarily interfere with the process of estimating the uric acid; but it should always be an indication for adding more silver, so that the iodine may not deprive the uric acid of its share.

In collecting the urate of silver precipitate on the filter, if the filtrate is not quite clear it may be necessary to pass it a

⁶⁴ MacMunn, *loc. cit.*

⁶⁵ Hoppe-Seyler, *Physiol. Chemie, loc. cit.*

⁶⁶ MacMunn, *loc. cit.*

⁶⁷ MacMunn, *loc. cit.*

⁶⁸ Hoppe-Seyler: *Physiol. Chemie.*, p. 297; Maly: *Centra'bl. f. d. med. Wiss.*, 1871, No. 54.

⁶⁹ MacMunn: *Clinical Chemistry of Urine*, pp. 96-98.

second or even a third time through the filter; the urate clogs the filter, and thus aids in its own filtration. When the precipitate has been collected on the filter, wash with plenty of water till the washings give no turbidity on addition of solution of sodium chloride, showing the absence of silver, and that no silver remains in the filter except that combined with uric acid. When this has been done, dissolve the precipitate and wash it through the filter with nitric acid, using plenty and filling the filter up to the brim, so that none of the precipitate on the sides may escape its action. Having got all the urate in solution, the final process is to estimate the silver by a colour test, and the uric acid combined with it is then found by calculation.

The process takes from 30 to 40 minutes, but I never grudge time to the washing process, which, with urines containing much uric acid, may be very slow owing to great clogging of the filter by the gelatinous precipitate.

It is possible, however, to attend to other matters during this process, merely every now and then pouring a little more water over the filter. The same process may be applied to a watery extract of blood, liver spleen, muscle, or other tissues of the body.

The watery extract is obtained by boiling the blood or tissue, previously finely divided, in about ten times its volume of water. I always see that the water is boiling before I add the blood, and I add only a little at a time, so that it shall be coagulated as soon as possible. After boiling for ten minutes the heavy particles are separated by decantation, and the fluid filtered. It is then evaporated down to about the volume of the original blood used and Haycraft's process applied as for urine.

This process for obtaining the uric acid from blood and other tissues is given by Salkowski and Leube¹ and they, after evaporating, proceed to obtain uric acid in crystals, but I have preferred to apply Haycraft's process.

I have, however, obtained uric acid crystals in some cases with a view to confirming my results with the Haycraft process, and they can be obtained either before or after the application of the process.

The advantages of Haycraft's process are its comparatively easy methods and relatively short time required to carry it through, also that it only requires 25 cubic centimetres of urine, so that the excretion can be estimated every half hour and changes, either natural or artificial, watched.

But for this it would have been often quite impossible to estimate the excretions during short headaches, epileptic fits, and short periods of high arterial tension, in all of which the process has enabled me to obtain interesting results.

Hermann² praised Haycraft's process as giving results sufficiently accurate for clinical purposes with greater celerity and ease of manipulation than any other process. It has, however, been criticised adversely by Gossage³ and by Salkowski.⁴

The latter critic, we must remember, is the author of another process for estimating uric acid, which, however, is too long for clinical use and does not, to my mind, necessarily ensure any greater accuracy than that of Haycraft. Its final stage depends on the crystallisation of uric acid from an acidified solution, a process which does not seem to me to give absolute freedom from error. As to results to be obtained from known solutions of urate, Haycraft's figures, given when he published his process, seem to me to leave little to be desired.⁵

The possible error from xanthem mentioned by Salkowski⁶ has been already dealt with by Haycraft, who points out that though xanthem combines with silver, the compound it forms is insoluble in nitric acid,⁷ and in any case the error from xanthem could only be a very small one.

Variations are also considered by Salkowski to be introduced into the process by variations in the amount of silver which combines with the uric acid, but this is a chemical point which I would leave Haycraft to deal with. I have previously said, however, that it is well, so far as possible, always

to add an excess of the silver; so that there shall never be any lack of it present to combine with the uric acid, and when this condition is fulfilled the resultant urate of silver is probably fairly constant in its composition.

To me it appears quite clear that the errors or possible errors from these causes are either so small as to be neglected, or so constant as not to prevent the accurate comparison of results, *inter se*, which for clinical purposes is all that I am concerned with. Further, I would ask: Have those who criticise the process by the light of their own results with it avoided all errors of manipulation? They appear to have repeated the process only some fifteen or twenty times. I have sometimes repeated it nearly that number of times in a single day.

The more important of my results during the last five years agree completely with each other, and with the physiological results produced along with them, and they show that, given the same conditions and the same drug, the results in the human organism are as constant as regards the excretion of uric acid as those in a chemical test tube.

My main facts, that acids diminish, and alkalies increase, the excretion of uric acid and that salicylates and phosphate of soda increase it greatly, while iron and lead diminish it, are the concordant results of very numerous experiments (in the case of acids and alkalies probably extending to three figures) with Haycraft's process, and can be verified as I know in some cases, even by the rough process of direct precipitation (Heintz) which, though doubtless inaccurate, is nevertheless, I think, to be believed when it shows that the uric acid excretion on one day is twice as great as on the previous day.

And these results are, as will now be easily seen, in complete accord with the physiological and pathological effects of the drug above mentioned. In the process itself also there are certain safeguards; thus, after the addition of the silver solution to a urine that contains much uric acid, the fluid may be almost converted into a jelly by the copious urate precipitate; on the other hand, when there is very little uric acid the precipitate may be represented by a slight haze. By bearing these points in mind, a rough idea of the amount present may be formed to the extent of estimating fairly accurately that one urine will contain from five to six times as much as another. Again, the filtration of the one that contains most urate will take five or six times as long as that of the one which contains least. Again, specific gravity and its relation to urea varies to a certain extent with the amount of uric acid present, and though other things may occasionally interfere with this, it is often a useful guide, and I feel more confidence in the accuracy of my results when they accord with this. Thus, take a urine of specific gravity 1025, if there is much uric acid urea will probably be only about 1.5 per cent., whereas if there is little uric acid, urea may be 2.0 per cent. or more.

While my results are constant, and are thus safeguarded, I fail to see that objections to the process resting on more or less theoretical points in chemistry need greatly concern me. I can further show that my results for the most part agree with what was previously known and found out about uric acid by other processes, thus:

Sir A. Garrod⁸ pointed out that blood or serum drawn over an inflamed joint contains no uric acid. Similarly, I have found that blood drawn from the seat of a local inflammation contains almost no uric acid, and that blood drawn early in an acute fever, as pneumonia, contains little or none. My explanation of the fact differs from that of Sir A. Garrod, but the facts alone concern us here.⁹

Again, it is pointed out in textbooks of physiology that after death the spleen more constantly contains uric acid than any other gland. Recently I have examined the blood from the heart cavities and at the same time a bit of the spleen, and have found in several cases that the spleen contains from four to five times as much as the blood. The liver, again, I have found to contain from three to four times as much as the blood, a fact also previously pointed out by Sir A. Garrod,¹⁰ and his writings on this subject led me to suggest¹¹ that,

¹ *Die Lehre vom Harn*, p. 94.

² *Zeits. für physiol. Chem.*, July, 1888.

³ *Proc. Roy. Soc.*, vol. xlv, p. 284.

⁴ *Zeitsch. für physiol. Chem.*, vol. xiv, p. 31.

⁵ *BRITISH MEDICAL JOURNAL*, vol. ii, 1885, p. 1100.

⁶ *Op. cit.*

⁷ *BRITISH MEDICAL JOURNAL*, vol. 2, 1885, p. 1101.

⁸ *Gout and Rheumatic Gout*, pp. 187 and 274.

⁹ See paper in next volume of *Med.-Chi. Trans.*, on Salicin compared with Salicylate of Soda as to Effect on the Excretion of Uric Acid and Value in Treatment of Acute Rheumatism.

¹⁰ *Lumleian Lectures. BRITISH MEDICAL JOURNAL*, vol. i, 1883, p. 549.

¹¹ *Journal of Physiology*, vol. viii, p. 214, and elsewhere.

when an acid diminishes the excretion of uric acid in the urine, it does so by diminishing the solubility of uric acid and causing its retention, in accordance with Sir A. Garrod's observation, in the liver, spleen, and joints, which are naturally less alkaline than the blood and tissue fluids of the rest of the body; and my results are obviously in complete accordance with this suggestion.

I have examined upwards of sixty specimens of blood, some drawn during life, but most taken from the heart cavities after death, and all tend, I think, to show that there is excess of uric acid in the blood—some specimens containing as much as 0.07 to 0.08 per cent.—under exactly the same conditions in which there is excess of it in the urine, and recently I have drawn the urine from the bladder at the same time that I took the blood from the cavities of the heart, and my results seem to show that excess in the urine corresponds and coincides with excess in the blood, and scantiness in the urine with a very small quantity, or even none, in the blood.

Again, my results show that blood drawn during acute fever, as pneumonia, or at the seat of a local inflammation, contains almost no uric acid, and I have previously mentioned that Sir A. Garrod found none in the blood and serum drawn under similar conditions. On the other hand, blood obtained after death from chronic wasting disease, especially if there are urate deposits in the joints, may contain a very large amount, as 0.08 per cent.

From all these facts, I infer that it is safe to argue from the urine to the blood, and that an excess of uric acid in the urine above the relation to the urea of 1 to 33, comes from and is the index of some excess of uric acid in the blood—a fact which I was led to infer several years ago from the physiological and pathological results I can produce by influencing the excretion of uric acid by the urine. And results thus constant and in harmony with those previously obtained by others are, I submit, incompatible with the existence of errors in the process sufficient to vitiate it for clinical and comparative purposes; and, used with due care and sufficient knowledge and experience of the sources of error, I believe that Haycraft's is by far the best process for the clinical estimation of uric acid at present known.

TUBERCULOSIS IN THE SAUROPSIDA.

Read in the Section of Pathology at the Annual Meeting of the British Medical Association, held in Birmingham, July, 1890.

By WALTER K. SIBLEY, M.B.CANTAB.,
Assistant-Physician to the North-West London Hospital.

THE vertebrate kingdom of animals has been divided into three main divisions, namely, the ichthyopsida (fish and amphibians), the sauropsida (reptiles and birds), and the mammalia (mammals).

Tuberculosis, the commonest disease of domestication both in man and many of the lower animals, has not as yet been found in the ichthyopsida, and, with but very rare exceptions, in only one division of the sauropsida—that is to say, the birds. Until the last few months back no case had been described in the reptiles, but now we find isolated cases occurring in the ophidia when kept in confinement.

PART I. Birds.—The abdominal organs—namely, the spleen and liver—were the parts primarily affected; the abdomen often being bulged from the enormous size of the liver, whilst the spleen was usually enlarged many times its natural size. In some cases no deposits were seen in the intestines, at other times these organs were beset with hard nodules often as large as beans. Microscopically the deposits in an early stage consisted of small groups of epithelioid cells, which were often apparently lying in vessels and enclosed by the vessel wall. In a much more advanced stage, each deposit consisted of a central mass of caseation, which was very irregular in outline, and surrounded by a zone of giant and epithelioid cells undergoing degenerative changes. Surrounding this was a ring of granulation tissue, the whole being encapsuled by fibrous tissue. In a still more advanced stage there was a considerable development of fibromyxomatous tissue. In these last stages no bacilli were present in the newly-formed fibrous tissue, but they occurred in densely packed groups in the caseous masses. In the intestine the lesions varied from a

small ulcer with thickened edges to pouches occasionally as large as small oranges springing from the gut walls. The walls of these were formed chiefly of caseous tissue and fibrous tissue. Often the lungs presented no obvious lesion. In a few cases there were caseating lymphatic glands in the base of the neck. Tubercular disease of the bones was very rare. In one case of a fowl there was a caseous mass in the neck of the thigh bone, and in another case some caries of the vertebrae. The example of a peacock presented several tumours in connection with the bones and joints, such as the scapula, sternum, and knee-joint; several of these tumours were as large as walnuts. Microscopically, they were formed of granulation tissue, with groups of epithelioid cells, and also giant cells here and there. Tubercle bacilli occurred in parts of these growths.

The bacilli vary much in different classes of birds both as to their number and arrangement, and also their actual size. All of these I believe to be merely local varieties of the same bacillus, the result of the slight differences in the soil in which they are growing. In birds the masses of caseation are very firm; often a furrow appears around them, by the growth and contraction of the fibrous tissue, and so the deposits, which are close to the surface of an organ, may in this way become completely separated. In some birds I have found the nodes formed in the liver lying loose in the abdominal cavity. The deposits appear never to go beyond the stage of caseation; they do not of themselves break down and form cavities, nor do they become calcified. There is a tendency for the disease to extend by continuity and to a considerable new tissue formation.

In a large number of cases which occurred in a farmyard among the fowls there was very strong evidence that the disease was hereditary in its nature, in that the birds became affected during consecutive years when about the same age—that is, young adult life. Also the disease was limited to certain broods, others which lived along with these remaining free from the affection. So, also, pigeons and wild birds feeding in the same yard never became subject to the affection, and the disease was not present in the neighbouring farms.

I have thus far found tuberculosis in the following birds dying naturally in confinement, the diagnosis being in every case made from the microscopical appearances of granulomata undergoing central degeneration, together with the presence of the bacilli: canaries, finches, fowls, geese, guan, owls, prafowls, pheasants, pigeons, swans and vultures.

PART II. Snakes.—The first case I met with of tuberculosis in snakes occurred in an Italian snake (*Tropidonotus matrix*), which died in the Zoological Society's Gardens; it is described in full in *Virchow's Archiv* for 1889.¹ This case was marked by the number of tumours in various regions of the body together with general visceral tuberculosis, tubercle bacilli being abundantly present in most of the lesions. Here the seat of the deposits appeared to be in the lymphatic vessels, there being a complete tubercular thrombosis of the main lymphatic channels. The microscopical structure of the lymphatic deposits was the same as that of the visceral deposits.

In addition to this well-marked case of tuberculosis I have examined a few specimens of what appeared to be the same disease, in which the lesions were confined to the subcutaneous tissues. A viper had a large tumour on the side of the neck, which had been watched for over two years without showing any appreciable change in its size. This tumour was composed of very dry caseous tissue lying in definite areas of fibrous tissue.

The study of tuberculosis in reptiles is of great interest, in that the disease can be studied with a complete control of the animal's temperature, as the temperature of a snake's body does not differ much from that of the surrounding medium, so they may be kept at a given temperature and the disease under its various aspects so studied. With this object I have kept various members of the reptile class (inoculated with tubercle) in incubators regulated to definite temperatures. After considerable difficulty I have succeeded in producing tuberculosis in snakes by the inoculation of pure cultures of the bacilli and keeping the reptile at about a temperature of 35° C.

¹ Translated in the *Journal of Anatomy and Physiology*, 1889.