TABLE I.

Date.	Time of Day.	Quantity.	Acid, its Equivalent in Grains of Oxalic Acid.	Amount of Uric Acid.	Percentage of Urea.	Amount of Urea.	Proportion, Uric Acid Urea.	Albumen.	Position.	Diet.
		Ozs.		Grains.		Grains.				
April 3	7.30 A.M. to 1.30 P.M.	3	8.593	1.218	3.8	53.01	$\frac{1}{17}$	Present	Up	Fish.
" "	1.30 to 3.15 P.M.	4	5.27	0.933	2.1	39.06	$\frac{1}{11}$	Absent	"	"
" "	3.15 to 7 P.M.	4	8.784	0.684	2.9	53.94	$\frac{1}{18}$	"	"	"
" "	7 P.M. to 4.50 A.M.	9	19.725	1.539	3.4	147.69	$\frac{1}{27}$	"	In bed	"
" "	4.50 to 7.30 A.M.	5 $\frac{1}{2}$	4.003	0.689	1.8	46.03	$\frac{1}{26}$	"	"	"
April 5	7.30 to 11.15 A.M.	6	4.394	1.679	1.6	44.64	$\frac{1}{27}$	"	Up	Meat.
" "	11.15 A.M. to 5 P.M.	5	4.394	1.165	2.2	51.15	$\frac{1}{23}$	Present	"	"
" "	5 to 7.45 P.M.	7	4.589	1.197	1.7	55.33	$\frac{1}{28}$	Absent	"	"
" "	7.45 P.M. to 5.25 A.M.	13	19.032	2.628	2.8	169.26	$\frac{1}{61}$	"	In bed	"
April 8	7.30 A.M. to 12.30 P.M.	4	6.44	1.374	3.2	60.72	$\frac{1}{19}$	Present	Up	Meat.
" "	12.30 to 2.45 P.M.	4	5.56	?	1.9	35.34	$\frac{1}{20}$	"	"	"
" "	2.45 to 6 P.M.	5	9.07	0.933	2.7	62.77	$\frac{1}{22}$	"	"	"
April 16	7.30 A.M. to 1.45 P.M.	3	8.3	0.7	3.4	48.23	$\frac{1}{25}$	Absent	"	Milk.
" "	1.45 to 4.45 P.M.	4	11.12	0.995	3.0	55.8	$\frac{1}{26}$	"	"	"
" "	4.45 to 8.45 P.M.	6	6.14	1.213	2.6	72.54	$\frac{1}{28}$	"	"	"
" "	8.45 to 5.15 A.M.	8 $\frac{1}{2}$	25.77	2.365	3.3	125.16	$\frac{1}{37}$	"	In bed	"
April 20	7.30 to 10.15 A.M.	2 $\frac{1}{2}$	3.416	?	2.8	62.55	$\frac{1}{22}$	Present	Up	Milk.
" "	10.15 A.M. to 4.10 P.M.	3 $\frac{1}{2}$	5.856	1.202	2.6	42.38	$\frac{1}{27}$	"	"	"
" "	4.10 to 6.46 P.M.	2	4.687	0.497	3.0	27.9	$\frac{1}{28}$	"	"	"
" "	6.45 P.M. to 4.45 A.M.	6	?	0.788	2.9	80.91	$\frac{1}{27}$	"	In bed	"

albumen regularly while he was on fish diet; secondly, he passed albumen once during a short period of milk diet; thirdly, the chief precipitate occurred in the morning before he took any meat; fourthly, he was taking meat regularly during a period of nearly a month, when he had hardly any albuminuria; and fifthly, he on many days took meat while in bed and yet passed no albumen.

Between February 10th and March 4th there was a period for which I am quite unable to account, during which the albuminuria almost ceased. During this time the urine was examined 13 times on twelve days while he was up, and albumen was only found certainly thrice and doubtfully once. One, however, of the other nine specimens was of evening urine which would naturally be free from albumen. At the time we hoped that the immunity was due to treatment, but this fond hope was subsequently shattered. He was ill in bed with rheumatism for several days ending March 23rd, when he rose again. For the next three days he passed no albumen, but it began again on the 26th.

Nor did the microscope help me any more than other methods of inquiry. I was never able to see any casts, but there were plenty of leucocytes in the field, and more than once some oxalate crystals. Oxalates have been found in several cases, but their importance is a very doubtful matter.

As a result of these various observations, I came to the conclusion that position was the most important of all the conditions that I could consider, and though this may act in many unknown ways, I am at present inclined to believe that it produced its effect simply by increasing the weight of blood resting on the renal veins. One thing seems to be very common in this class of patients—a weak circulation and a general loss of tone. This would greatly predispose to a condition of venous stasis such as I suppose to be present, and to which the kidney is, by reason of its double capillary system, more liable than other parts. I have, at any rate, failed to find any other factor likely to have a bearing on the case.

The greater liability to albuminuria in the forenoon is, perhaps, due to the weaker state of the circulation at that time, and since exercise is the best stimulant, the instances in which we could postpone the albuminuria till the afternoon by keeping the patient at rest may probably be explained by the absence of exercise.

Lastly, I would add that this case is on all fours with the cases of Rooke,² Dukes,³ Pavy,⁴ and von Noorden,⁵ and that I should offer this explanation for this class alone. There are many other forms of intermittent albuminuria; indeed, in children, I believe slight albuminuria is generally intermittent, but I should think that they probably depend on quite other causes.

² BRITISH MEDICAL JOURNAL, 1878, ii, 596.

³ Lancet, 1878, ii, 794.

⁴ Lancet, 1885, ii, 706; 1886, i, 437.

⁵ Deutsches Archiv f. klin. Med., Leipzig, 1886, xxxviii, 20

Treatment was of little avail in this case:

From Jan. 6 to 15	...	Fish diet	...	No drugs	...	Albuminuria.
" " 16 to Feb. 7	...	Meat	...	Hydro-chloric acid	...	Albuminuria.
" Feb. 8 to 13	...	"	...	Atropine	...	Little albuminuria.
" " 14 to 18	...	"	...	Arsenic	...	"
" " 19 to 25	...	Fish	...	"	...	"
" " 26 to Mar. 11	...	Meat	...	"	...	Little albuminuria up to March 4; then increasing.
" Mar. 12 to 23	...	Fish	...	Salicin	...	For subacute rheumatism; in bed.
" " 24 to 26	...	"	...	Arsenic	...	No albuminuria
" " 27 to 28	...	"	...	"	...	Albuminuria.
" " 29 to Apr. 5	...	"	...	Iron	...	"
" Apr. 6 to 11	...	Meat	...	"	...	"
" " 12 to 14	...	"	...	Digitalis and arsenic	...	"
" " 15 to 20	...	Milk	...	"	...	Little albuminuria.

MULTIPLE MULTICULAR CYSTS OF THE MESENTERY.

By WILLIAM ROBINSON, M.S., M.D.,
Stanhope.

SIR SPENCER WELLS contributed to the BRITISH MEDICAL JOURNAL of June 14th, 1890, a note on mesenteric and omental cysts, and a description of the latter was given by Dr. Ransom in the JOURNAL for November 8th, 1890. These reports induce me to publish a few brief notes of a similar rare case.

A little girl, aged 24 months, was brought to me on September 7th, 1889, owing to the great and increasing size of her belly preventing her from walking and stooping. The enlargement was first noticed three months before. The child was anæmic and had signs of slight rickets (enlarged bone ends, lax joints, and history of sweating of the head); there were nebulae of the cornea and a history of severe bronchitis at 12 months of age. The belly was now 20½ inches in circumference; the enlargement was most marked on the left side, and was not quite uniform; on palpation the growth could be felt to extend from the left hypochondrium (where there was stomach resonance) to the brim of the true pelvis, into which it did not seem to dip, as the lower rounded end could be mapped out, and from the left loin to the median line, near which was a depression in its inner border; in fact, in shape and position it felt just like a uniformly enlarged kidney. The growth was highly elastic, freely movable, and dull on percussion, but the note varied in different parts of it; the left loin and groin were both dull. Sometimes on very slight percussion some resonance could be made out on the front of the tumour at the upper part. It was impossible to elicit a wave of fluctuation either in the whole or any part of the swelling. Alteration of the patient's position caused no alteration in the signs. Under chloroform all the above signs were verified, and an examination *per rectum* showed the pelvis was not invaded by the growth. On puncturing the growth with a hollow needle 2 or 3 drachms

of a straw-coloured limpid fluid, alkaline in reaction and highly albuminous, were obtained; the needle seemed to traverse a thick wall before the fluid came. On boiling the precipitated albumen with double the amount of strong acetic acid, the precipitate became gelatinous. On standing the fluid deposited a clear gelatinous material. Urine, 8 ounces in the twenty-four hours, acid, no albumen, no sugar, no casts.

On the following day I opened the abdomen, Mr. John Waldy assisting me and Mr. A. C. Waters giving chloroform. The incision was 3 inches in length in the median line. On dividing the peritoneum, an extremely thin, transparent cyst wall projected through the incision, and on sweeping the finger round the growth numerous soft adhesions were torn through; the growth was felt to be free of the pelvis and adherent behind. On tapping the cyst, 2 pints of fluid, similar to that obtained the day before, escaped. The mass was then gently drawn through the wound, and the descending colon throughout its entire length and the mesocolon separated from the growth, several double ligatures being used; the peritoneum was stripped off the tumour as this separation was made. Several smaller cysts were punctured, and one large multilocular cyst with thick white fibrous walls was also punctured; the pedicle contained a vessel which proved *post mortem* to be the inferior mesenteric vein. There were several small cysts the size of small marbles in the mesentery an inch or more away from the main growth, which I ruptured by squeezing. Both kidneys and the spleen were felt to be normal. The operation lasted over an hour; extremely little blood was lost, and the child was much collapsed after it.

At 9 P.M. the same night the child was crying as if in pain; there was rattling in the trachea; the lips were natural in colour; the pulse was almost uncountable and the temperature 102.5° F. Drop doses of tinct. opii had been given two or three times. At 2.15 A.M. the next day the child died.

The *post-mortem* examination, made the same day by Mr. Waters in my presence, showed the growth to have sprung from the upper part of the mesentery. Everything in the abdomen was found to be satisfactory; there had been no oozing of blood; the kidneys, spleen, liver, intestines, pelvic organs, and heart were normal. The lungs showed numerous depressed bright pink patches—to the extent of two-thirds of all the lung substance—which on section were carnified and contained hardly any air, and felt firm (atelectasis). The chest cavity was much encroached upon by the pressure from below, the lower ribs being bulged outwards.

A CASE IN WHICH SLOUGHS OF RENAL TISSUE WERE PASSED BY THE URETHRA.

By JOHN M. RATTRAY, M.B.,

Frome;

AND

J. GREIG SMITH, M.A., F.R.S.E.,

Bristol.

T. H., manufacturer, aged 45, a short, thick-set, plethoric man, was seen medically for the first time on April 29th, 1890. He had never been ill before, and the cause of the trouble he was treated for so many weeks he attributed to catching cold while attending a funeral three days before he sought advice.

His first symptoms were those of liver congestion. He looked very ill, and had flushed face, suffused eyes, thickly coated tongue, liver tenderness, and heavy bilious-looking urine. The first attack was a short one, for on May 8th he was so far improved as to be able to go downstairs and venture out of doors for a little fresh air in the verandah in front of his house. His temporary recovery was, however, of very short duration.

On May 11th he was taken seriously ill, and went to bed, where he remained for many weeks. His symptoms now were precisely those of acute Bright's disease. Fever ran high, urine scanty, dense, of smoky colour, with casts of blood and granular hyaline matter. Treated with hot bottles, milk diet,

jalap and saline medicines, he made but little improvement, and flatulence and vomiting caused him intense distress.

On May 16th, his pulse becoming intermittent, a consultation on his case was held by Dr. Rattray, his usual medical attendant, and Mr. R. L. Willecox, of Warminster. Mr. Willecox, on palpating the left kidney deep down in the abdomen, struck on a very tender spot, and he gave it as his opinion that there might ensue a pyelonephritis, consequent on the probable presence of a calculus in the pelvis of the kidney. During the fortnight subsequent to the consultation, active symptoms subsided, the temperature lowered to normal, the redness of the urine cleared away, and Mr. H. was in comparative ease. The albumen disappeared, and for a few weeks the urine was plentiful (65 to 90 ounces in twenty-four hours), pale coloured, low specific gravity (1010 to 1015), and deposited a heavy cloud of mucus on settling. Microscopically, clusters of pus corpuscles, granular matter, and epithelial squames were detectable daily. An offensive smell in the urine suggested to Dr. Rattray the expediency of washing out the bladder with boroglyceride and Sanitas fluid. Sounding for stone was tried on several occasions, but gave negative results. The case went on in a chronic state for many weeks. Recrudescence of active symptoms took place from May 24th to 27th, June 15th to 18th, and again on July 18th and onwards. On June 17th Dr. Rattray made a careful examination of the patient, and, thinking he detected a fullness of the left loin, he asked Mr. H. for the benefit of the opinion of a surgical authority. While waiting to decide as to this, the patient gradually recovered again, and for the following weeks so improved that he was able to leave his room and walk out of doors, one day going about a hundred yards from his house.

On July 18th Mr. H. had the final exacerbation of his symptoms. This was certainly his worst attack. Temperature rose to 104°, urine was loaded with albumen, pain in left loin set in, and flatulence and vomiting were continuous, intractable, and terribly distressing. Jalap, pilocarpine injections, hot bottles, etc., were tried all day, and at last, in despair and despite albuminuria, Dr. Rattray was impelled to administer morphine hypodermically. This was done with good result, and Mr. H., by means of repeated small doses of morphine, was tided over the worst of the attack.

Dr. Rattray had always cautioned Mr. H. to be careful to always pass his urine into a vessel, so that he might at all times see whether there passed from him a piece of grit or diseased substance.

On July 20th Mr. H. showed Dr. Rattray a piece of tissue which he passed the same morning after a considerable amount of pain. This was kept, and subsequently examined by Mr. Greig Smith, who saw Mr. H. in consultation on July 22nd. After consultation, expectant treatment being decided upon, there was a gradual and uninterrupted progress to recovery. Urine became quite normal. On August 25th the patient left for Weymouth, and in three weeks returned in perfect health. He has now resumed his business, and declares that he never felt better in all his life.

REMARKS BY MR. GREIG SMITH.—When I saw the patient the urgent symptoms had abated, but the patient was still seriously ill. I could detect no signs of renal enlargement, and no local evidence of perinephric inflammation. The case altogether was very puzzling, and no diagnosis was come to. The pieces of tissue which Dr. Rattray showed me threw a new light on the case. To the naked eye one of them, the size of a pea, was evidently the apex of a renal papilla, and this was afterwards confirmed by microscopic examination. I had seen pieces of renal tissue passed in a case of sarcoma of the kidney with large blood cysts, but was not aware that pieces of renal tissue were sometimes passed as sloughs in acute nephritis. This latter conclusion, or rather, working hypothesis, we adopted, and I returned, not having performed the operation I went prepared for. It was hoped that the passing of these sloughs signalled the termination of the complaint, but no definiteness in prognosis was ventured upon.

On referring to Ebstein's work¹ I find that he speaks of the occasional separation of small portions of renal substance in cases of suppurative disease of the kidneys, and their discharge through the urethra. He refers to two cases, both of

¹ Ziemssen, vol. xv, p. 357.