lupus and lupus erythematosus there are bonds of essential relationship, whether we regard their clinical features or their histological characters. Next, it has been made probable that all forms of lupus are in occasional relationship with the state of health which gives liability to tuberculosis and to diseases of the scrofulous class; but in limitation of this it has also been proved that many lupus patients are in excellent health, and that there appears to be some proneness to cancer as well as to tubercle. That lupus begins as an inflammation under the influence of various exciting causes and derives its peculiarities from the part affected, and the special structures in that part, as well as from the age and proclivities of the patient, has also been asserted. Thus it has been sought to discredit the supposition that lupus, or any of its peculiar forms, deserves such terms as *sui generis*, the implied creed being that they are closely related amongst themselves, and also to other various pathological processes. The doctrines of infective spreading and system-contagion have been asserted as being explanatory of the phenomena of lupus, and as finding in it some of their best illustrations. The laws of hybridity and those of partnership in disease have also been appealed to, as well as those of modification in hereditary transmission, under which latter it has been attempted to establish the pedigree of the very remarkable malady known as Kaposi's disease.

I much regret that I have not had time to say anything in reference to treatment, since I might have drawn from it very important evidence in support of the conclusions placed before you. This, however, must be reserved for another occasion.

In reference to the introduction of not a few new names, and the alteration of some old ones, I am well aware that I have laid myself open to criticism, and that I have taken on me grave responsibility. My innovations in these matters, however, I willingly leave to the judgment of others. If found useful they will live, if otherwise let them die. I can only assure you that I have done nothing from caprice, but that I have honestly endeavoured, as far as my ability admitted, to make clear a very intricate subject, and to lighten the labours of those who will follow me.

LETTSONIAN LECTURES

ON

SOME POINTS IN THE SURGERY OF THE URINARY ORGANS.

Delivered before the Medical Society of London, January, 1888.

By REGINALD HARRISON, F.R.C.S.,

Surgeon to the Liverpool Royal Infirmary; and Lecturer on Clinical Surgery in the Victoria University.

LECTURE II.

THE PATHOLOGY OF THE ENLARGED PROSTATE VIEWED IN RESPECT TO ITS CAUSATION AND PREVENTION, AND THE TREATMENT OF SOME COMPLICATIONS ARISING OUT OF IT.

MR. PRESIDENT AND FELLOWS,—I purpose this evening speaking of the pathology of the prostate relative to some points in connection with the prevention and treatment of certain disorders arising out of it when it becomes hypertrophied. The causes of hypertrophy of the prostate have formed the subject of much careful observation, and I feel some hesitation in again referring to them. Still, on the other hand, though my views may not prove generally acceptable, I am inclined to think that a narrative of the observations upon which they are to a large extent based may serve a useful purpose in indicating lines of thought which may result, sooner or later, through the investigations of others, in clearing up doubtful passages in the physiology and pathology of this part. In the first place I shall speak of the prostate as forming the retentive apparatus of the male bladder, under whose influence the urine is collected and held, irrespective of any glandular function which, by the nature of other structures it may contain, it is capable of exercising. And I think it will not be difficult to substantiate the proposition that, in the human species at all events, this part may with greater propriety be designated the "prostate muscle," rather than the "prostate gland," for whatever function it may exercise intermittently, relative to the process of generation in its latter sense, seems to be subservient to the physical part it is continuously playing as a portion of the retentive apparatus. In support of such a view I would draw attention to Mr. Ellis's important paper (*Royal Medical and Chirurgical Transactions*, vol. xxii), On the Muscular Arrangements of the Genito-Urinary Apparatus, wherein it is

remarked, "I would propose the name *orbicularis vel sphincter urethrae* for both the prostate and the prolongation around the membranous urethra, whilst I would confine the old term 'prostate' (with out the word gland) to the thickened and more powerful part near the neck of the bladder. This orbicularis may be considered as only an advanced portion of the circular layer of the bladder, though it must have the power of acting independently of the vesical fibres." But if its muscularity is admitted, we must conclude that it for the most part exercises its function in conjunction with the bladder in the form of a hollow muscle, for it would be against the nature of things for it permanently to exist in the shrivelled and contracted state it presents after death. We have been too much accustomed to regard the prostate from its *post-mortem* aspect, that is to say, as a mass of muscle of the size and form of a chestnut, in which is contained some secreting tissue. For concluding that it thus exists during life, I believe there are no substantial grounds, as it seems to me that under no circumstances, save the rare and momentary one when the bladder is absolutely empty, does it present such an appearance. On the contrary, the muscular fibres are spread out like a funnel, with the apex downwards so as to form a strong muscular support for the bladder and its varying amount of contents, the degree of expansion being naturally relative to what the viscus may contain. Hence the action of the prostate may be said to be just as continuous as that of the heart. In reference to this important point it will be necessary that I should furnish some reasons for such a conclusion.

In the first place, the clinical examination of a healthy person with varying amounts of urine in his bladder affords no evidence that the prostate presents the contracted appearance we are accustomed to see; on the contrary, when the finger is introduced into the rectum, in the natural condition, the parts are felt to be disposed in the manner I have indicated, and providing a muscular floor for the bladder and its varying amount of contents.

But the retentive function of the prostate is more strikingly shown when we proceed to what I would speak of as actual demonstration. And for this purpose we have only to observe what follows in connection with certain surgical operations on those parts with which most of us are familiar. Incisions may be made into the male urethra, in any part of its course, as far as that point which we are accustomed to call the apex of the prostate, without any incontinence of urine following. I have seen cases of lithotomy by the median operation retain full control over the bladder during the whole period of their convalescence, in spite of the dilatation to which the prostate has been subjected by the introduction of the finger and the extraction of the stone. And this remark applies equally to cases of external urethrotomy.

When, however, the knife impinges to any appreciable extent upon the prostate, as in the lateral operation for stone and the modified median operation which I have recently drawn attention to, where the prostate is divided, from that moment incontinence takes place; the patient has no command over his urine; he can neither collect nor expel it, and in this condition he remains until the healing process has made considerable advance. We have here striking evidence, not only as to the habitual function of the prostate relative to the contents of the bladder, but that the action of the part must be unceasing in its character, subject to the circumstances under which it has to distribute its force over the area it supports or brings into action. How completely the prostate forms the lower section of the bladder was demonstrated to me in a striking manner only a few weeks ago, and in a way that I had not previously noted. It was a case of lateral lithotomy in a young man where, ten days after the operation, there was some free hæmorrhage. As the bleeding did not proceed from any part of the perineal wound, I had the patient put under ether on the operating table, and, suspecting where the bleeding came from, I introduced the nozzle of a Higginson's syringe into the orifice of the urethra, expecting that the fluid would run out of the open urethra; this it certainly did, but not until it had first entered the bladder, out of which the fluid escaped, together with some old clots which had evidently collected in the viscus, just as water would do out of the side of a cask in which a hole had been made. This I was able to see, as the wound was held open with retractors under a strong light. In addition, however, to testimony of this kind, I am convinced, from a careful examination of the prostate, both before and after operations on this part, where the introduction of the finger into the bladder formed a detail of the proceeding, that the more usual condition of the healthy prostate is one of relaxation, and not of contraction, as we are accustomed to see when death has taken place.

But further, when there are no such physical functions to perform relative to the contents of the bladder, the prostate, as a muscle, so

far as I have been able to ascertain, only exists in a rudimentary form. I refer to those distressing abnormalities known as extroversion of the bladder, where, from the fissured nature of the structures, there is no receptacle for the urine which escapes from the ureters as it is excreted. Here, though the sexual sense may be normal, there is from the nature of the parts no necessity for any provision being made for the collection and retention of what the kidneys secrete. I have not been able to discover, though I have taken considerable pains to do so, that in these abnormalities hypertrophy of the prostate has ever been noticed. This seems to me to be a point of some significance in connection with what has been already urged. But though in an argument of this kind I would prefer to substantiate my position from positive evidence rather than by a deduction from the negative, I cannot help remarking that, if the office of the prostate was solely in relation to the sexual act, the pathology of the part would tend to indicate this more than I believe it does. It is not very common to meet in practice with cases of acute suppuration commencing in the follicles of this part, where one would think that irreparable damage must have been done, and yet we have evidence to show that the individual has not necessarily been unsexed by this process. Nor can I assert, though the opportunity for doing so has now for some years been abundantly provided me in my operations, involving varying degrees of mutilating the prostate, that I ever extinguished the procreative powers.

In the next place, I would point out that adult man seems to require some special provision such as the prostate affords for the purpose I have just indicated, by reason of the exceptional degree of pressure to which the most dependent portion of his bladder is constantly subjected. And in making use of the words "exceptional degree of pressure," I do so in contradistinction to what is found, so far as I have been able to observe, not only in the animal kingdom, but in other varieties of the human species which are not in exact correspondence with the type I have taken. As demand and supply in the disposition of muscular tissue are proportioned, it seemed to me that we had hitherto failed fully to recognise and appreciate all the circumstances which rendered it necessary that the adult male bladder should, as I have already pointed out, be provided with a muscular apparatus of considerable strength at its most dependent point.

But in what respects, it may fairly be asked, does the adult male bladder differ from other varieties of its own species, and from those animals which in their higher organisation more closely approach it? I maintain that in man the perpendicular axis of what I would speak of as urine-pressure falls directly upon the outlet of the bladder, whereas in the female, not only are there other means of supplying muscular support to the base of the bladder, but in consequence of the difference in the pelvic organs, a considerable portion of the weight of the viscus, especially in its more extreme degrees of distension, is borne by the pubic portion of the pelvis. Further, allowance must be made in the female sex for no provision being necessary for seminal ejaculation, in which function the muscular fibres of the prostate undoubtedly play an important but occasional part.

In the higher mammalia—excepting, perhaps, in the case of dogs—the prostate is not so largely developed as in man. In reference to this point, Mr. Bland Sutton kindly informs me that, although he has conducted many thousand *post-mortems*, yet, excepting in the doubtful case of dogs, he has never seen an enlarged prostate—at all events, such a one as would cause obstruction to the flow of urine. In animals, it will be observed that the perpendicular axis of urine pressure, when the bladder is tolerably full, does not fall, as in man, on the outlet, but on the parts which support the bladder, according to its varying degree of distension. In the dog, it seems to me not improbable that the exceptional degree of prostatic development is related to the somewhat peculiar condition under which, in this animal, micturition is performed.

As my opportunities of examining the urinary apparatus in the lower animals have been few, I feel much hesitation in making reference to any observations I may have rather casually made so far as they related to the point under discussion. It, however, appeared to me that where the habits of the animal, as observed in some apes and marsupials, necessitated the frequent adoption of an erect position, the males had a much greater development of the prostate muscularly than was usually found in other kinds of quadrupeds.

I regret very much that we have not at our disposal more abundant means for referring to dissections illustrative of the comparative anatomy of the body. I am sure they would prove of much assistance in investigations of this kind, having relation to normal and morbid human anatomy. But, apart from this, if time permitted, it

would not be difficult to adduce further evidence to show that in man the collection and storage of urine, if I may make use of such a term, implies a degree of muscular support to the most dependent portion of the viscus, for which no such complete provision has to be made under any other condition of life I am acquainted with. I should not, however, feel at all surprised if, when the "missing link" is discovered, he is found paying the penalty for his assumption of the erect posture by developing in some degree an enlarged prostate.

But to proceed; if the prostate represents the chief means of retention and support for the contents of the male bladder, is it necessary to pursue the argument for the purpose of explaining why it becomes hypertrophied? Does not one follow upon the other? Are not the circumstances of individual life sufficient to explain why it happens to one person and not to another?

Many of the facts connected with the natural history of hypertrophy which have been collected for us, or with which, in our daily practice, we are familiar, seem to me to support the view I have taken relative to the function of the part. In the first place, it is not met with during those periods of life which are most remarkable for muscular activity and development; on the contrary, it is called into existence where, pathologically speaking, quantity seems to supplement deficiencies in quality.

But as with similar changes in other parts of the body, so may prostatic hypertrophy be shown to be exactly compensatory; whilst in others, by an excess of the process, it may be proved to be detrimental. And in reference to this point I would remark that we have been too much accustomed to regard prostatic hypertrophy as necessarily a morbid and hurtful process. To such a conclusion I think exception may be taken on substantial grounds. It is stated by Sir Henry Thompson that actual hypertrophy of the prostate exists in about 34 per cent. of men at and above 60 years of age; that it produces manifest symptoms in about 15 or 16 per cent. From this it would appear that the larger proportion of persons are not injuriously affected by the change, a circumstance which seems to suggest that the majority of persons with large prostates are in some way or other benefited by them; and this deduction is considerably borne out by everyday experience. I am disposed to think, from my own examinations, that the frequency of some degree of prostatic hypertrophy is thus rather under- than over-estimated. Though it is not easy to form an absolutely correct estimate of the size of these parts relative to standard measurements by rectal examination, my conclusion is that a far greater number of males over 60 years of age have some enlargement of the prostate, and are never in any way conscious of the alteration, than we should conclude from the figures I have just referred to. How frequently, for instance, when we are examining elderly persons, say for some disorder of the rectum, we make the discovery that their prostates are enlarged, though they have no reason to be conscious of this change. I can at this moment recall many instances of this in my own practice where the persons are in every sense of the word in the enjoyment of the most perfect health, with an entire absence of any urinary ailment. Quite recently I was asked to examine a gentleman who, in the course of an acute bronchitis, had retention of urine for the first time in his life. This patient had an enormous prostate, which must have been in course of formation for several years. Surely, under these circumstances, in the first instance the hypertrophic change must have been called into existence for a distinctly conservative object, and had been merely disarranged by the violence of the cough from which he was suffering.

But though prostatic hypertrophy is, as I have shown, in the greater number of instances more or less compensatory, it may, as in others, be in excess of what is required, and thus prove detrimental. In the latter category, reference no doubt will occur to us of those examples where the prostatic mass is made up of projections having a somewhat lobulated or nipple-like form, where the degree of irritation and obstruction excited is often very considerable, and apparently out of proportion with the cause. These I would speak of as being the structural upheavings of a frequently-contracting muscular ring. In a muscle or part undergoing hypertrophic growth, and where the process is prompted by circumstances which are obviously liable to some degree of variation, tissue production may be excessive or become unnecessary. Such excess would naturally tend, under the contraction of the part, not only to protrude itself where the resistance was least, but to assume a more lowly-organised form than that which in the first instance was developed. In this way, I believe, these masses of more or less degenerated prostate tissue are formed, which subsequently cause so much vesical irritation and irritability. And this observation, whether a large prostate still remains almost entirely muscular or has become largely converted into fibrous tissue, has an important bearing in practice in those cases where we have to express

an opinion in reference to the probabilities as to whether the use of the catheter will be a temporary or a permanent expedient; for where there is evidence to the touch that fibrous tissue predominates largely over the muscular, the use of the catheter is generally perpetual; whilst, on the other hand, when the prostate is found soft and yielding to the touch, as when it remains muscular, a complete restoration of function may be anticipated. Thus, then, may the resistance which is constantly going on in the most dependent portion of the bladder against that downward pressure which is exercised by the collection and retention of urine in the bladder, prove the common cause of its hypertrophy.

But this is not the only condition under which hypertrophy involving the lower section of the bladder can be studied. In the examination of the bladder, both after death and in the course of surgical operations, it is impossible not to be struck with the altered relations which frequently exist between the body of the viscus and its neck or outlet. In early adult life the bladder may be regarded as an abdominal rather than a pelvic organ; as years advance it gradually sinks within the pelvis; whilst still later on it will often be found to have become further depressed within the pelvic cavity. In this way I have seen a prominence given to the floor of the prostate, which was really due not to the development of more prostatic tissue, but to the depression or partial prolapse of the posterior wall of the bladder. This artificial condition of obstruction I have endeavoured to show, in some instances, not only precedes prostatic obstruction, but, further, is followed by the development of a strong muscular buttress between the orifice of the ureters, which in many instances leads to the obstruction of what otherwise would prove, by the sinking of the posterior wall of the bladder, an inconvenient pouch. Thus hypertrophy of the prostate and the adjacent part may not only be due to the natural resistance of urine-pressure, but may also be called into existence for the evident purpose of supplementing a structural defect of more or less accidental origin.

It is impossible to observe the function of micturition in the child and the adult without recognising that in the two conditions there are wide differences in the influences that the will exercises upon the merely mechanical act. In the child the process almost approaches the involuntary; the calls of Nature are impulsive, and must be obeyed without hesitation. This may be illustrated in a variety of ways, in health as well as in disease. In adult man this is to a large extent changed, as we find the process becomes much more subservient to the will than perhaps at first sight we should feel disposed to admit. That there is a wide difference in our respective powers of continence in this respect I am willing to admit. In some individuals the function of micturition seems in health but very slightly removed from a purely voluntary act. Let me illustrate what I mean. A medical man, closely approaching 70 years of age, asked me to examine his rectum for some slight ailment. In doing so I found that he had a very large, soft prostate. I mentioned this to him, when he remarked, "I could hardly believe it, as I have always had such excellent waterworks. I have been in the habit of not passing urine for eight or ten hours." Here then is an illustration of a man who submits his bladder to the same degree of urine-pressure at 70 as he did when he was 27. Again, a captain of one of our ocean steamers, in consulting me in reference to some urinary disorder clearly connected with a large prostate, in comparing his present with his past powers of retaining urine, remarked that, in consequence of remaining on the bridge of his steamer for long periods of time, particularly in thick weather, without leaving it to pass water, he had acquired a power of retention which had gradually become constant. Is it remarkable that Nature had thus substituted quantity for quality, and had made provision against the effects of a muscular strain which could not have been inconsiderable? We might from our daily practice multiply examples of this kind where persons, so far as was known, were not deriving anything but good from that excess of muscular tissue which we are accustomed to speak of as the prostate gland. If we admit that the muscular fibres, of which the prostate is so largely composed, exercise the function I have urged relating to the contents of the bladder, I do not think we can be at a loss to understand how they become hypertrophied.

Let me, however, not be understood as implying that in the human species the prostate has no relation with the sexual act. Evidence is not wanting to show that, both as a muscle, as well as by reason of its glandular element, it exercises purely physical functions relative to this important but casual function, independently of the continuous action I have laid stress upon in relation to micturition. The less important function of the glandular portion of the prostate has been, I believe, correctly described by Dr. Handfield Jones (*The London Medical Gazette*, vol. v, 1847), in the following words:—"Its part in

the generative function is probably not to prepare any essential element of the fecundating fluid, but merely an appropriate viscid material, involved in which the seminal animalcules may be more securely transported on their destined route."

Passing to some considerations relative to the prevention of prostatic hypertrophy, and the complications arising out of it, I would observe that more care should be systematically taken to preserve the muscular power of the bladder, and to aid it artificially when the necessity obviously arises. If a man goes on using his bladder as he would a water-bottle, from adolescence to old age, he must not be surprised if Nature prevents it breaking down entirely by the process I have described. The timely use of the catheter has often, I am sure, provided the means for warding off changes which unfortunately do not always cease when they have become precisely compensatory. Nor is there anything exceptional in this, for as age advances we recognise the importance, and make use of artificial assistance in a variety of ways which I need hardly particularise. In the clinical history of most cases of hypertrophy of the prostate, we can generally trace a period when a weakness of the viscus indicated itself by a frequency in action, which was subsequently corrected by an overgrowth in the part. But, unfortunately, the process of hypertrophy, as observed in the prostate, is not infrequently in excess. In by far the larger proportion of these instances the individuals are able not only to keep themselves comfortable by the use of the catheter, but I question very much whether their lives are in any way shortened by the continued inconvenience they have thus to submit to. Many of us could record some remarkable example of longevity under these circumstances. But making due allowance for this class of cases, there are others whose lives, in spite of careful and efficient catheterism, are rendered miserable and useless by what they suffer in this way. For no sooner does urine collect in their bladders to any appreciable extent than they require to get rid of it. Many expedients have been resorted to, both in the way of drugs and operative procedures for the permanent relief of this condition, about which it would be impossible to speak in detail. Let me, however, refer to a case (*JOURNAL*, December 24th, 1881; April 8th, 1882) which I reported some years ago, where I tapped the bladder from the perineum through the enlarged prostate, quite independently of the course of the urethra, and where a cannula was retained in this position for two months with great comfort to the patient. He was a man about 80 years of age, and had suffered much from almost all the complications that an enlarged prostate is capable of giving rise to. But the future of the case was perhaps still more interesting, for he completely recovered and lived for several years, and in this I include the acquisition of the natural and unaided use of his bladder. Further, the explanation of this was found in the fact that during the time the cannula was retained in his bladder his prostate underwent a gradual diminution in size, which was noticed by all who had the opportunity of watching and examining him. His condition prior and subsequent to the operation proved a most remarkable and instructive contrast. I may add that this process has been repeated, not only by myself but by others, with very satisfactory results, and this tends in no slight measure to substantiate the views I have already advanced relative to the pathology of the disorder. And in connection with puncture through the prostate, I would like to say a few words in reference to cases where relief has been afforded under similar circumstances by an incision through the part similar in many respects to what is adopted in lateral lithotomy. There can be no doubt that prostatotomy, which affords a free and incontinent escape for urine from the bladder, has proved of very great temporary and permanent service in this class of cases.

At the meeting of the International Medical Congress at Copenhagen, in 1884, I took the opportunity of illustrating this method of treatment as an alternative in cases where the catheter was of no avail in affording relief. And I will again refer to this operation, as sufficient time has now elapsed to enable me fairly to judge of the results obtained. The kind of cases to which this treatment has been applied are those of difficult micturition, due to the large prostate, which are not adequately relieved by the use of the catheter. This class, therefore, includes instances where there is unusual difficulty in introducing the catheter; where bleeding almost always attends the use of this instrument; where the withdrawal of the urine is followed by no sense of relief; and where the bladder, by the constant presence within it of pus and tenacious mucus, is converted into little else than a chronic abscess through which urine percolates. These, as well as some other forms of prostatic disease which might be included, are practically unrelievable by the catheter, and soon terminate in a painful death. For the relief of such conditions various expedients for establishing a more or less permanent communi-

cation with the bladder, other than the urethra, have been practised, namely, puncture above the pubes, by the rectum, and from the perineum, with the retention of a cannula or tube for the discharge of urine at these several points. Excellent in design as these proceedings are, they appear to me to fall short in one important respect—in not dealing with the cause of the obstruction. Two of these measures are open to objection, on the ground that the artificial canals are inconveniently placed as permanent vents, for it would seem desirable that the urethra should, as far as possible, be utilised and the external opening for the escape of urine be dependent. On considering various plans of treatment which had been applied under the circumstances mentioned to the obstructing prostate, it appeared to me that, by combining Cock's well-known and safe operation for opening the membranous urethra with Mercier's for dividing the prostatic bar from within the urethra, as will be presently noticed, it would be possible to obtain precision with an increased freedom from risk. Such an operation was suggested by the late Mr. Guthrie, but I cannot find that he ever tested it in practice. The want of anaesthetics probably interfered with the progress of this as well as of other departments in surgery. Apart, however, from the operative procedure, it appeared to me, from some experience I have had in lithotomy and other operations on the parts constituting the neck of the adult male bladder, that the means which had been subsequently adopted for rendering the prostatic section thus made permanent were very inadequate. Further, I had been impressed with the advantage that followed the employment of suitable bougies as dilators in cases where the prostate threatened to obstruct micturition (*The Prevention of Prostatic Obstruction*. London: Churchill. 1881). Influenced by observations of this kind, I was led to attach considerable importance to the treatment immediately following section of the enlarged prostate with the view of rendering it more permanent than had hitherto been attempted. The necessity for prostatotomy having been determined by the symptoms presented in each case, as well as by physical examination, I will describe briefly the operation and after-treatment employed.

The former consists in opening the membranous urethra, from the perineum, on a guide, and introducing the finger within the prostatic urethra; the obstructing portion of the prostate is then divided a little to one side of the median line, partly by incision with a curved probe-pointed bistoury from within outwards, and partly by division with the finger when the latter is feasible; a large drainage-tube is then introduced into the bladder. I attach considerable importance to the prolonged use of the drainage apparatus, as the object is to render the section of the prostate not a temporary one as after a lithotomy, when no such provision is made, but permanent. Hence, I am in the habit of retaining these tubes for six, eight, or ten weeks. If, after such periods, on removing the tube, I find that a catheter can be made to enter the bladder easily along the natural route, or if, as it sometimes happens, urine forces its way, in spite of the drainage tube, along the natural passage, I regard these as indications that the object in view has been obtained. The perineal wound is then allowed to close. Let me briefly illustrate this by the narration of one of my earliest cases (*JOURNAL*, June 9th, 1833).

A man, aged 63, came under my care in February, 1833, suffering from a large prostate, which was a constant source of irritation to him. The straining and forcing to urinate were but very slightly relieved by catheterism, and the only way he could get any sleep was by tying the instrument in, and this was generally followed by some cystitis. A fortnight after I saw him, I opened the prostatic urethra by a median perineal incision, and incised the prostate laterally. In doing this I found that, though the prostate was not very large, the orifice of the bladder was much obstructed by one of the nipple-like enlargements, which are sometimes more effectual in rendering micturition difficult and catheterism uncertain than some larger masses. A drainage-tube was introduced and retained for four weeks, when it was removed, and the perineal opening allowed to heal. In this case, so perfect was the drainage that the whole of the urine escaped by the apparatus, and the patient was kept absolutely dry throughout. He remained under treatment for two months after the operation. Three months after he returned to report himself. There was a slight fistulous opening in the perineum at the lower angle of the wound, through which urine sometimes passed in drops during micturition. He could now hold his urine for several hours. Some slight enlargement of the prostate could be felt from the rectum, but this did not appear to have increased. A full-sized catheter was easily introduced without any hitch or obstruction in the prostatic urethra, or at the neck of the bladder, being felt. This patient was seen at intervals for two years after the operation, and remained perfectly well. He has since neglected to report himself. Though I advised him to pass a catheter for himself occasionally, I believe that he did not do so.

Taking the case thus operated on, I have noted the following results: 1. The operation has merely afforded just such a temporary relief as the retention of a catheter or cannula would do as a drain, the condition of the patients, so far as recovery was concerned, being more or less hopeless at the time of operation. 2. The establishment of a permanent vent for the urine through the perineum. In two instances where this occurred the patients exchanged a life of misery for one of comparative comfort. The shorter route here proved most acceptable. 3. The substitution of easy for difficult catheterism. In one patient I thus operated on, though he had to use his catheter afterwards, so long as I heard of him, which was for over two years, he never had any difficulty in passing the instrument, and the bladder continued to be far more tolerant of urine than it had previously been. 4. The complete recovery of the patient, as I have just illustrated.

In discussions relative to the method of operation adopted, it has been pointed out that in some cases of this kind, it must be absolutely impossible to explore the whole length of the prostatic urethra with the finger. This circumstance has never influenced me, or seriously interfered with my carrying out the object in view. I never met with a prostate of such a size or shape as to prevent a staff or guide being passed into the bladder, and where a director will go a knife may safely follow.

But, to proceed, though a more or less permanent vent or opening for the urine, either above or below the pubes, may prove of value in some instances of urethral obstruction due to prostatic hypertrophy, where ordinary catheterism is useless so far as the amount of relief that is afforded, it is not applicable to all cases of this kind. For we must remember that in the former class of cases it is merely a matter of urethral obstruction that we have to deal with, whilst in others the symptoms are for the most part due to the irritation that is provoked in the bladder by the protruding masses of more or less degenerated prostatic tissue. The one is to the other, as well as to the bladder, as urethral stricture is to vesical calculus. The examination of specimens of hypertrophy in connection with their clinical history clearly points in some to the mechanical irritation which protruding masses of more or less prostatic tissue excite in the interior of the bladder. In connection with the treatment of such cases, where the lives of the individuals are not only absolutely useless, but most miserable, I am glad to see there are signs that something now may be systematically done for their relief. Heretofore we have been drawing for any experience we may possess in reference to this point upon what I would speak of as the casualties of lithotomy. There is much, however, in favour of direct surgical interference in these exceptional cases of grave prostatic obstruction and irritation, as the nature of the growth, unlike cancer, renders a return of it improbable. I have now on three occasions deliberately removed by perineal section considerable portions of the hypertrophied prostate, with permanent relief to the patients; and I have no hesitation in recommending the adoption of this course in suitable cases, after, if possible, perineal exploration of the part with the finger. Now, three instances may seem very insufficient data for basing such a recommendation upon; but it must be remembered that this is an exceptional remedy for the purpose of meeting exceptional circumstances. No operation of this or any other kind would in my judgment be warrantable so long as a man can keep himself reasonably comfortable with the use of the catheter, and therefore the percentage of cases where more radical measures would seem to me to be indicated is really extremely small. But we must be provided for these, and though they are oftener met with in the aged and feeble, with reduced powers of repair, whose chances of recovery from any operation are reduced, such means as I am referring to must not be too hastily set aside in these days when the practice of operative surgery, especially in relation to the interior of the body, has been shorn of many of its dangers, and is altogether very different from what it was, even at the commencement of the present century. I am induced to make these observations partly because I think it would be a misfortune if operative procedures were extended to what I would speak of as ordinary forms of prostatic obstruction, and partly because I think the risks attending the removal of protruding portions of the prostate have been somewhat exaggerated, and thus surgeons have been deterred from undertaking operations which, in proper cases, have afforded both immediate and permanent relief. So far, I have been able to do all I desired in the way of removing more or less of the obstructing prostate by proceeding from the perineum, which I have always opened with considerable freedom. In connection with the subject of prostatectomy, or removal of portions of the prostate, under such circumstances as I have indicated, I would take this opportunity of expressing the interest with which I read the paper by Mr. McGill, and the no less instructive discussion which followed it, at a recent

meeting of the Clinical Society. In this paper the author demonstrates by instances how portions of the prostate might be removed by scissors through a suprapubic incision. I am disposed to think that this, on still further trial, will be found the best way to deal with such cases, for it places the operator in view of those projecting masses of degenerated prostate tissue which invariably protrude towards the cavity of the bladder, and which really are the cause of the distressing symptoms of irritation which accompany them. Care, however, will be necessary, as I have just urged, in the selection of cases for operations of this kind, it being distinctly understood that they are undertaken, not because the prostate is large, but because it has induced symptoms which cannot be alleviated in any other way. You may remove the prostate entirely, but this necessarily renders the individual incontinent for life. Lobulated masses proceeding from an hypertrophied prostate and isolated fibromas in the part, may thus be dealt with with permanent advantage; but this is a very different thing to extirpating the prostate in its entirety, or seriously mutilating the muscular ring, which under all circumstances is essential to the integrity of the bladder as a urine holder.

The following instance, which I will briefly relate, substantiates the proposition with which I opened this lecture; and, as it similarly bears upon the operative point now under consideration, I may consistently close my remarks with it.

In August, 1882, a patient, aged 64, came under my care with symptoms of primary carcinoma of the prostate, which caused bleeding, continuous pain, and irritability of the bladder. As no treatment afforded relief, I did a free median cystotomy, and proceeded with my finger and a blunt gouge to enucleate the prostate and the growth connected with it. In this I succeeded, with the exception of a small portion in front, which I could not get away. There was very little bleeding, and the patient made a good recovery, though he lost power of retaining his urine. Some degree of retentive power eventually returned, but it became necessary to provide him with an apparatus for controlling the incontinent escape of urine from the bladder. He lived for fourteen months after the operation, and was able to return to his work as a stevedore. He eventually died from a recurrence of the disease in the glands of the groin. I showed the specimen of what was removed at a meeting of the Royal Medical and Chirurgical Society, in connection with a discussion bearing upon the subject.

A FURTHER INVESTIGATION

INTO

THE SO-CALLED HENDON COW DISEASE,

AND ITS

RELATION TO SCARLET FEVER IN MAN.

Abstracted from a Report to the Agricultural Department of the Privy Council, read before the Pathological Society, January 17th, 1888.

By PROFESSOR EDGAR CROOKSHANK, M.B.,
Of the Bacteriological Laboratory, King's College, London.

Description of the Streptococcus isolated from the Ulcers of the Teats.—Determination of the Species: Streptococcus Pyogenes.—Acute Abscesses; Distribution of Streptococcus Pyogenes.—Nature of Contagium undetermined in the Exanthemata.—Post-mortem Appearances in the Hendon Cows.—Post-mortem Appearances of Cows from Wiltshire.—Post-mortem Appearances of Inoculated Calves.—Summary.

At a special meeting of this Society I laid before you the results of an investigation which, though admittedly incomplete in some details, enabled me to bring forward evidence to show that the theory that a certain eruptive disease of the teats of cows at Hendon was the very disease which we call scarlet fever in man could not be maintained. That was the object of my communication, while as a side issue—which I, nevertheless, regarded as of greater interest than the overthrow of the above-mentioned theory—I brought forward evidence to show that this disease was in reality cow-pox. I was anxious that these conclusions should be made known without delay, partly to allay the public anxiety, and partly because I wanted to give every opportunity for my statements to be examined. Thus I wished to afford anyone interested in the matter an opportunity for visiting the farms while the details were fresh in the memory of the milkers, and before the disease had totally disappeared from among the cows.

I pointed out that the study of the streptococcus which I had isolated was not yet completed. I had yet to study the effects of inoculation, and I was therefore only in a position to say that this streptococcus corresponded morphologically and on cultivation with the so-called scarlatina streptococcus. I inclined to the belief that these organisms were identical, not only with each other, but with certain other organisms which had been regarded originally as distinct species.

I promised further investigation on these points, and I also stated that I thought I should find in a calf at the Royal Veterinary College the *post-mortem* appearances which had been described by Dr. Klein. This animal showed symptoms of septicæmia, and has now been killed and carefully examined. I will therefore proceed to lay before you the details of these and of further inquiries in connection with this subject.

I desire to lay this further evidence before you, so that you may be concerned to-night in discussing the identity of the disease I have investigated with the disease which has been fully described clinically and pathologically by Dr. Cameron and Dr. Klein, and the micro-pathology of which has been so fully illustrated by Dr. Klein in a very extensive series of lithographed plates.

DESCRIPTION OF THE STREPTOCOCCUS ISOLATED FROM THE ULCERS OF THE TEATS.—I will first describe more in detail the chain-forming micrococcus which I found in the discharge from the ulcers.

The individual cocci are small spherical cells with a special tendency after fission, for the resulting elements to remain attached to each other, forming chains or rosaries. There may be a few, three or four, elements linked together, or a number of individuals forming straight or serpentine chains. The size of the individual elements is not always equal, here and there in a chain a larger element appearing.

In plate cultivations the appearances of the colonies are not very striking. They appear to the naked eye, after three or four days, as extremely minute, greyish-white, translucent dots, which, under the microscope, have a slightly yellowish-brown colour. They are finely granular and well-defined. They do not liquefy the gelatine, and after weeks do not exceed the size of a pin's head.

If the surface of nutrient gelatine solidified obliquely is traced over once or twice with a platinum needle, bent at the extremity into a little hook charged with the micrococci, a ribbon-shaped film develops in two or three days at 18° C. This film is composed of minute greyish-white translucent dots or droplets, which can be more easily recognised with the aid of a pocket lens.

According to the number of organisms seen on the jelly, the dots or colonies may be completely isolated or form a more or less continuous film. This film by reflected light has an iridescent appearance like mother-of-pearl, but has a bluish or bluish-grey tint by transmitted light, and with a pocket lens appears distinctly brownish.

The gelatine is not liquefied, and even after several weeks the cultivation is limited to the inoculated area and the individual colonies are not larger than pins' heads. Cultivated on the oblique surface of nutrient agar-agar at 37° C. the growth is very similar, forming a film composed of minute dot-like colonies like grains of sand. But the film appears less transparent, is whiter, and the colonies have a greater tendency to get irregular in form. If inoculated with one tracing of the needle the growth is scanty, but tends to get thicker in the centre than towards the margins, which may have a terraced appearance.

Inoculated in the depth of the gelatine there appears after a day or two, at 18° C., a thread-like growth along the track of the inoculating needle. This delicate thread is found on examination with a pocket lens to consist of a linear series of extremely minute granules. In a few days more the beads or granules become more marked, but even after weeks the cultivation only appears like a string of minute white compact globular masses or grains. In broth at 37° C. the cocci in twenty-four hours create a turbidity, and gradually develop chains varying in length according to the age of the cultivations. Even in forty-eight hours there may be chains of 8, 10 or 20 elements. After a few days the growth settles down at the bottom of the tube in the form of a white deposit, while the supernatant liquid becomes clear again.

Cultivated in milk at 37° C., they convert it in two or three days into a solid mass, while uninoculated control tubes remained liquid. Inoculated subcutaneously in the ear of rabbits, they produce in two days an inflammatory thickening, with erysipelatous redness, and without suppuration.

DETERMINATION OF THE SPECIES: STREPTOCOCCUS PYOGENES.—From its microscopical appearances, its characters on cultivation, and its effect on animals, it is obvious that this organism is identical with the *Streptococcus pyogenes* of other authors.