another condition to which dryness and furring have been attributed, namely, habitual openness of the mouth, especially during sleep; I find but three instances in which this attracted notice, and I am persuaded that other causes of drying and furring are far more important.

In short, the dry furred tongue is essentially due to two causes which are connected together—want of saliva and want of wear. Among the causes of want of saliva, the most important is a state of system which cannot be otherwise defined than as failure of nutrition and vital power. With how great a variety of disorders this failure is associated I need not recapitulate; I have expressed the greater degrees of it by the term "prostration."

7. Furred or shaggy; not dry.

Pneumonia, pleuro-pneumonia	...	...	2
Perityphilitis, perforation of appendix	...	...	1
Chronic constipation	...	...	1
Granular kidney; uræmia	...	...	1
Total	...	...	5

Observations relating to preceding cases.

Pyrexia (temperature 102° to 104°)	...	...	1
Hyperpyrexia	...	...	0
No observations as to temperature	...	...	2
Average temperature (of 3 cases)	...	...	99.8°
Much prostration	...	...	2
Chiefly on liquid diet	...	...	2
Strictly limited to liquid diet	...	...	1
Saliva abnormally deficient	...	...	2

Died, 2; recovered, 1; not improved, 1.

7 (D). Furred or shaggy; dry.

Disease of brain (coma)	...	2	Stricture of œsophagus	...	1
Hemiplegia (embolic)	...	1	Diarrhœa	...	2
Chorea	...	1	Perityphilitis	...	1
Valvular disease	...	1	Abscess (in axilla)	...	1
Aneurysm or dilatation of aorta (dry diet)	...	3	Acute rheumatism	...	1
Broncho-pneumonia	...	1	Typhoid	...	1
Pleurisy	...	1	Convalescence from typhoid	...	1
Cirrhosis of liver	...	2	Febricula	...	1
Enlargement of liver (cause uncertain)	...	1	Movable kidney and rheumatism	...	1
Total	...	...	Chyluria	...	1

Observations relating to preceding cases.

Pyrexia (temp. 102° to 104°)	...	...	3
Hyperpyrexia	...	...	0
No observation on temperature	...	...	2
Average temperature of 22 cases	...	...	99.1°
Much prostration	...	...	6
Prostration not severe	...	...	2
Chiefly on liquid diet	...	...	9
Strictly limited to liquid diet	...	...	5
No food by mouth (fed by rectum)	...	...	1
Dry diet	...	...	3
Saliva abnormally deficient	...	...	8?

Died, 9; recovered, 7; relieved, 8.

DONATIONS.—The annual report of the Leeds General Infirmary acknowledges the receipt of £150 from the Yorkshire Football Club, making a total of £1,000 in eight years.—The Trustees of Prison Charities have given 100 guineas, additional, to the National Hospital for Consumption at Ventnor, and 100 guineas to the Metropolitan Convalescent Institution.—The Sheffield Church Burgesses have given £120 to the Sheffield Public Hospital and Dispensary, and £68 to the Jessop Hospital.—The Mary Wardell Convalescent Home for Scarlet Fever has received £100 as "A Jubilee Offering."—Mr. Harold Smith has given £100 to Charing Cross Hospital.—University College Hospital has received £85 from the People's Contribution Fund.

THE General Court of the Governors of Guy's Hospital have sanctioned the erection of a residential medical college upon a site immediately adjoining the hospital. It is proposed that the sum required for this purpose, estimated at £20,000, should be raised by private subscriptions among the governors, the medical staff, the lecturers and teachers in the medical school of Guy's Hospital, and their friends. The special appeal fund lately raised, and which is only available for purely hospital purposes, now amounts to £85,000.

THE TREATMENT OF DIPHTHERIA AND TONSILLITIS.

By THOMAS F. RAVEN, L.R.C.P., M.R.C.S.ENG.

FROM time to time communications appear in the JOURNAL, and other medical periodicals, relating to the treatment of diphtheria. Various drugs, applications, and antiseptics are employed, and, when the result is satisfactory, a record is made of successful treatment, and sometimes even the specification of a remedy is announced. The clinical features of the disease, however, are often omitted, and when this is so, the value of the observation must suffer.

It is held by many authorities that sore throat, with membranous deposit, is in itself a sufficient evidence of the disease called diphtheria. The writer of the article on diphtheria, in Quain's *Dictionary of Medicine*, inclines to this view. Others, however, are driven to the conclusion that sore throat, with membranous deposit, occurs sometimes sporadically, sometimes in epidemics (with local appearances exactly resembling those of true diphtheria), which can be clearly distinguished, especially, "after the event," from diphtheria. My own experience compels me to take sides with the latter view. Recently I witnessed, in an institution containing eighty children, a severe epidemic of exudative tonsillitis. The disease was sthenic in type, and the local appearances of wash-leather-like patches of considerable extent upon the pharyngeal mucous membrane were indistinguishable from those of diphtheria. The cases were almost all severe, with temperatures, especially early in the disease, as high as 105°; relapses were common. Three cases of acute rheumatism (with cardiac mischief), one of erysipelas of the face, and one of diffused cellulitis with suppuration (in an adult) were associated with the epidemic. The disease was extremely infectious, and it undoubtedly arose from the poisonous effects of sewer gas. Had I treated the patient with sulphurous acid, or biniodide of mercury, or sulphite of magnesia, or turpentine, I should have been justified, by authority, in recording the successful treatment of sixty cases of diphtheria, by one or other of these remedies. For under careful management, the disease ran its course, and all got well; and, although chlorate of potassium was freely given, I never thought of attributing the recovery of the patients to its employment. During the epidemic albuminuria was never found in any case, nor swelling of the glands at the angles of the jaw; and afterwards there was not a single instance of paralysis or peripheral neuritis.

Within the last few weeks a similar but smaller epidemic has been under my observation. Almost exactly the same symptoms appeared, but the disease was less severe than in the epidemic which I have just given an account of. Thirteen cases of exudative tonsillitis occurred, some mild, some severe. In one case laryngeal symptoms appeared, but subsided in the course of a few days without operation being necessary. All the patients got well, and no remedy but chlorate of potassium was given, and in no case were the sequelæ of diphtheria manifested. The disease, which resulted from the escape of sewer gas into a lavatory and near a bedroom window, was decidedly communicable. This was shown by the fact that when the first case was removed to a lodging for supervision, a little girl in the house contracted the disease.

It is to be hoped that the Collective Investigation Committee, with the new schedules, clear and simple as they are, giving scope to every contributor to record his own experience in his own way, will do good work by collecting sufficient evidence by which a distinction can be definitely drawn between diphtheria and exudative tonsillitis. The subject is one that can hardly be overrated in importance and interest, and most practitioners have experience and notes bearing on the matter.

SCARLET fever has, we learn, broken out on board the Thames training ships *Shaftesbury* and *Exmouth*, lying off Grays. Sixteen cases from the former and one from the latter have been removed to the hospital.

FRENCH ACADEMY OF MEDICINE.—At a meeting of the Académie de Médecine on March 13th Dr. de Sabin, of Rio de Janeiro, and Dr. Lusk, of New York, were elected Corresponding Members in the Section of Surgery. In addition to these two gentlemen, the names of Professor Victor Horsley and Sir William Mac Cormac, of London, Dr. Macewen, of Glasgow, and Dr. Sayre, of New York, were submitted to the Academy.