circumstance that the articulation had already been the seat of a chronic inflammation. Even with this possibility before one, it may be questioned whether simple drainage and intermittent irrigation of an acutely inflamed joint would lead not only to the recovery of the limb, but also to the perfect restoration of the functions of the articulation.

In the second case, recorded below, the question of the influence of a previous chronic inflammation did not confuse the issue. The case was a simple one of acute suppuration following injury. The patient, a stoker, a vigorous, healthy man, aged 23, was admitted into the London Hospital on November 7th, 1887, with acute synovitis of the right knee-joint.

On November 1st he had fallen and had received a violent blow upon the knee. He was disabled and was taken home. The joint did not become visibly swollen until the evening, some hours after the accident. There was no wound. The patient had never had any previous joint trouble, nor, indeed, any notable illness.

When seen on admission the patient appeared ill, and was much worn out by pain and want of sleep. The right knee-joint was a little flexed, was extremely distended with fluid, and was the seat of great pain. The skin covering the articulation was red and oedematous, and the oedema had extended some little way beyond the joint district. The temperature was 100° F. The case was evidently one of suppurative synovitis. The limb was fixed upon a straight back splint, and the joint was opened by two lateral incisions, each about one inch in length; through the incisions escaped much synovial fluid mixed with pus and flakes of lymph. A drainage-tube was passed through the joint underneath the patella, and the irrigation apparatus was adjusted precisely as in the case already detailed. Cold water containing a little boracic acid was allowed to run through the joint without intermission night or day for one month. The temperature became normal the day after the operation, and never rose again above 99°. The patient was at once quite easy.

The tube was removed on December 8th, and was replaced by two short tubes for either incision. On December 17th all drains were left off, and the splint was removed. There had been no rise of temperature, no suppuration, and no sign of a relapse. The wounds were now (December 17th) quite healed, and passive movement was commenced. The patient left the hospital on January 8th.

In the following month I exhibited the patient at the Harveian Society. His recovery had been complete. The patella was quite movable, and he could execute a complete range of movements with the right leg; indeed, there was no appreciable difference between the right limb and the left, except that the former presented two cicatrices.

The treatment advocated is exceedingly simple, and would appear to be efficacious. In my previous experience of other methods of treatment I have never seen complete restoration of function to follow an acute suppuration of a large joint.

THE FORMATION AND EXCRETION OF URIC ACID, CONSIDERED WITH REFERENCE TO GOUT AND ALLIED DISEASES:

BEING A THESIS FOR THE DEGREE OF M.D. IN THE UNIVERSITY OF OXFORD.

By A. HAIG, M.A. and M.D. Oxon., M.R.C.P.,

Physician to the Royal Hospital for Children and Women, Waterloo Bridge Road; and Assistant Physician to the Metropolitan Hospital.

SOME researches on the excretion of uric acid, on which I have been engaged for several years, now seem likely to throw new light on the part played by this substance in the causation of disease.

The diseases to which my results more especially apply are gout a certain form of headache (migraine) (*Med. Chi. Trans.*, vol. lxx), some cases of epilepsy (*Neurologisches Centralblatt*, March 1st, 1888), and, lastly, rheumatism, acute and chronic, though the evidence of uric acid causation is not equally strong in all.

My researches show, (1) that the excretion of uric acid (*Journal of Physiology*, vol. viii, Nos. 3 and 4) can be controlled by drugs and foods in a way and to an extent not previously known; that, whatever the amount excreted to-day, it may, within certain limits, be made either greater or less to-morrow.

2. That, when uric acid excretion is diminished relatively to the urea, the diminution is due to retention in the liver and spleen, and possibly in some other organs and tissues, of part of the uric acid which was formed; and, conversely, when uric acid excretion is increased relatively to the urea, such increase is due to a washing out of a store previously accumulated in the above organs and tissues.

3. That the formation of uric acid bears an almost constant relation to that of urea, though its excretion can be varied by food and drugs to a considerable extent.

4. That uric acid is probably never in excess in the blood, except when alkalies or other solvents wash it out in considerable quantities from the stores which have gradually accumulated in the above-mentioned organs.

5. That excess of uric acid in the blood, brought about in this manner, is a sufficient cause of many of the symptoms of the diseases I have mentioned, and probably of others also.

6. That drugs and foods which diminish the solubility of uric acid diminish its excretion and lead to retention and accumulation; and, conversely, foods and drugs which increase its solubility facilitate its excretion and prevent retention.

7. That drugs which diminish the solubility and excretion of uric acid are harmful in the diseases connected with it, while drugs which increase its solubility and excretion are useful in these same diseases.

I began my research by investigating a form of migraine, or, as I now prefer to call it, uric acid headache, from which I had been a sufferer all my life; and my first observation of importance was that leaving off butcher meat (*Practitioner*, Aug., 1884, and March, 1886) produced gradually-increasing immunity from attacks. This and the clinical relationship of the headache to gout led to my examining the urine and finding a very marked rise in uric acid excretion, corresponding to the headache (*Med.-Chir. Trans.*, vol. lxx).

I then began to speculate as to the cause of this uric acid fluctuation; and, having noticed the apparently great gout-producing power of ales, and their strong acid reaction, I was led to test the effects of acids, alkalies, and, finally, of many other drugs, on the excretion of uric acid, obtaining results some of which form the subject of this paper.

The processes used in my experiments are as follows: for urea, the hypobromate test, with the apparatus of Dupré, variations of temperature and pressure being allowed for; for uric acid, Professor Haycraft's process (see *JOURNAL*, vol. ii, 1885, p. 1100); and for acidity, a solution of phenol-phthaleine with a graduated solution of soda.

I soon found that acids (*Journal of Physiology*, vol. viii, Nos. 3 and 4) always diminished the excretion of uric acid, and alkalies always increased it; and it thus became easy to see that butcher meat, beer, wine, etc., had made my headaches more frequent, because they caused a rise of acidity, prevented the free elimination of uric acid, and accumulated a store of it in the body, which on a future occasion would produce a headache. Alkalies, or a somewhat vegetarian diet, on the other hand, by facilitating the regular excretion of uric acid, would prevent any such troubles; and, in proof of this, I found that by first accumulating uric acid with acids, and then washing it out into the blood with an alkali, I was able to produce the headache in question.

Sir A. Garrod (*JOURNAL*, vol. i, 1883, p. 653 [Lumelian Lectures]) has discussed the gout-producing powers of wines and beers and puts it down entirely to the unfermented matters they contain. In discussing the matter he says the light wines which contain a considerable quantity of acid vegetable salts do least harm; while the heavy wines, as those of Spain, do most harm and are almost devoid of vegetable salts.

Speaking of acidity he says that fresh fruits diminish acidity, through conversion of their vegetable salts into carbonates, and this I think gives us the key to the whole matter.

Both the above kinds of wines are strongly acid, but in the light wines the acidity is due to vegetable salts which are converted in the body into carbonates and act as alkalies, so that they cause no retention of uric acid, and have, as observed, but little gout-producing power; the heavier wines, on the other hand, have an acidity due to more fixed acids, they act as a dose of acid does, and have great gout-producing power, and the same applies to ales and stout, which are as I have shown (*Med.-Chir. Trans.*, vol. lxx, p. 12 of paper) strongly acid and cause great retention of uric acid; so that the acidity of these beverages explains their action in gout production far better than their unfermented matters, the action

of which is not very clear; and in gouty patients who indulge in these beverages after an interval of care, the rise in acidity they occasion and the increase of arthritic pains can be watched.

Foods, again, influence acidity by the salts they contain (see Sir W. Roberts, *Urinary and Renal Diseases*, ed. iv. p. 56), fruits and vegetables, as we have seen diminishing acidity, and animal foods containing large amounts of phosphates, sulphates and chlorides increasing acidity. Some foods may also contain salts having a special solvent action on uric acid, for example, phosphate of soda.

Sir A. Garrod in the above-quoted lectures gives a very interesting table from Lehmann of the effects of different diets on the excretion of uric acid: thus

	Urea.	Uric Acid.
Mixed Diet.....	32.4	1.1
Animal Diet	53.1	1.4
Vegetable Diet	22.4	1.0
Non-nitrogenous Diet	15.4	0.7

and he draws the inference from it that uric acid is not so much influenced by the nature of the food as urea is; but if the uric acid excretion is regarded relatively to the urea, this conclusion will, I think, have to be altered; and I now give the table with the addition of my figures, showing the relation of uric acid to urea on each diet:

	Urea.	Uric Acid.	Relation.
Mixed Diet	32.4	1.1	1—29
Animal Diet	53.1	1.4	1—38
Vegetable Diet.....	22.4	1.0	1—22
Non-nitrogenous Diet...	15.4	0.7	1—22

A long series of observations on the relative excretion of uric acid to urea has convinced me that a relation of 1—33, that is, 1 grain of uric acid for 33 grains of urea is about the average or normal; and taking a long period a result somewhat near this is obtained; thus in four months I excreted 41,804 grains of urea and 1,281 grains of uric acid, relation 1—32.6; but daily variations in this period ranged as high as 1—19 or as low as 1—45; but over a long period these more or less balance each other.

To apply this to the above table it will be seen that on mixed diet there was slightly excessive excretion of uric acid; on animal diet considerably diminished excretion (retention), and on vegetable diet excessive excretion, some of the previous retention being washed out. And a continuance of the animal diet would obviously soon produce a store of uric acid sufficient to account for anything from a "bilious" to a gouty attack. The results in these tables are similar to those I have obtained with various diets, and the relation of uric acid to urea are just those I meet with in my every-day work.

I have been led to infer from all these facts, that while the formation of uric acid alters but little from its relation to urea of 1—33 (subject, I have no doubt, to some small variations) its excretion is subject to frequent and great fluctuations, which account for its action in the production of disease.

Iodide of Potassium has in five-grain doses but little action on the excretion of uric acid: in larger doses it acts apparently merely as an alkali. Iodine taken as the tincture has little or no direct effect.

Phosphate of Sodium is a well-known solvent of uric acid, and it does, just as one would expect, greatly increase its excretion: it is probably the chief solvent of uric acid, which is naturally present in the body.

Salicylates have the very important action of rendering the excretion of uric acid independent of acidity, that is, a rise in acidity does not cause retention while salicylates are present in the blood; and this fact may explain much of the value of these salts in diseases connected with uric acid (see discussion on my paper in the *Proceedings of Roy. Med.-Chir. Soc.*, January to March, 1888, p. 325).

Lead and Iron.—The urates of these metals are very insoluble, and they cause, as one would expect, retention of uric acid, which, as we shall see, may explain their action in several diseases (Paper read at Roy. Med. Ch. Soc., April 28th, 1888, reported in *JOURNAL*, April 28th; and also Sir A. Garrod, *JOURNAL*, vol. i, 1883, p. 495).

Colchicum.—This indirectly lowers acidity and facilitates the excretion of uric acid. It may also have a direct action on the nerves in relieving the pains of gouty arthritis.

Besides drugs and foods, there are other ways in which the acidity of the blood and tissue fluids may be raised and retention of uric acid brought about, as, for instance, the formation of acids which are known to occur in dyspepsia; but a more important source of acids is probably to be found in the skin, for a large

amount of acid is daily eliminated in sensible and insensible perspiration, and Sir A. Garrod (*Gout and Rheumatic Gout*, ed. iii, p. 258) has shown that when this function of the skin is checked by cold the acidity of the urine at once rises, and *vice versa*, and I have often repeated this experiment, both intentionally and unintentionally, on myself. The application of these facts to the causation of gout by exposure to weather or other sources of chill is sufficiently obvious, and is precisely similar to the case of beer, wine, meat, etc., previously spoken of.

And now to apply some of these results to the explanation of disease.

The causation of the headache of which I have spoken is, I think, fairly well worked out; it can be produced at any time in those subject to it by causing a *plus* excretion of uric acid with alkalies, the *plus* excretion and the headache being, as I believe, the signs of excess of uric acid in the blood. If there is no store of uric acid in the body alkalies will fail to produce either the *plus* excretion or the headache. I have also found an identical uric-acid fluctuation in some cases of epilepsy (*Neurolog. Cent.*, March 1st, 1888), and I think it probable that uric acid may be the cause of the fits in these cases just as it is of the headache of which I have spoken. Sir A. Garrod (*Gout and Rheumatic Gout*, pp. 459-61) pointed out many years ago that he had found blood drawn during fits of epilepsy to contain excess of uric acid, and this observation affords important support to my belief that we may safely argue from excess of uric acid in the urine to its excess in the blood.

Diet and drug action also affords many important parallels between the headache of uric acid and epilepsy. Thus, the value of a vegetarian diet in epilepsy is amply attested, and I believe that not a few of those engaged in its treatment habitually order their patients to abstain from meat.

Iron is believed by many to do harm in epilepsy, and lead poisoning sometimes causes epileptic fits. Iron, so far as I know, does harm in the uric acid headache; and both these metals cause, as we have seen, retention of the uric acid.

In a considerable number of cases the fits have been treated by diet and drugs, intended to influence the uric acid factor, and so far with some apparent success; but, as it is almost impossible to say when a fit has been caused and when it has been prevented, epilepsy presents a problem which is much more difficult to solve than that of the uric acid headache.

I have also been much struck by the large number of cases of epilepsy which have a strong family history of gout, rheumatism, or minor uric acid troubles.

Another result of my research is the conviction that a considerable amount of hypochondriasis, heaviness, depression, and irritability may be produced by causing a rush of uric acid into the blood and its excessive secretion, and that the same conditions may be cured or made to give place to the opposite feeling, by driving the uric acid out of the blood by an acid, or something that causes retention. There is here, again, an important parallel between epilepsy (*Neurologisches Centralblatt*, March 1st, 1888, p. 4) and the uric acid headache; for both are preceded by retention of uric acid, and, preceding both, certain bright and joyous feelings (*loc. cit.*) have been noticed; no doubt corresponding to this retention, just as they correspond to it when it is produced by acids.

And these mental effects of uric acid have led me to think that some forms (especially the paroxysmal ones) of melancholia and mania might yield some results to an investigation of their uric acid connections, though no doubt, in all such diseases, what may be at first functional and due to uric acid, is at last associated with vascular and organic change which is incurable.

In this connection, it is very interesting to note that Dr. Broadbent, in his Croonian Lectures of 1887, says: "An unbroken series of gradations may be traced from the irritability and depression of spirits attending functional disorder of the liver up to complete melancholia with delusions;" and he further says that this is accompanied by *plus* tension (of pulse), which, if not the cause, is at least the index of the condition of the system on which the mental state depends. These observations are of double interest to me—first, because of the unbroken series of gradations traced from conditions I can produce up to serious mental disorder; and, secondly, because they tend to confirm an opinion I have held for some time, that a high tension pulse is, to some extent and in the absence of other obvious causes, an index of excess of uric acid in the blood.

The relation of gout and uric acid is not a matter of doubt; but my researches give, I think, a better explanation of its intimate

pathology than was previously obtainable, by showing the way in which urates come to accumulate in the body, and that this accumulation is for the most part due to acids. We have also seen that acids may be introduced from without in food and drinks, may be formed internally in dyspepsia, or may be retained by deficient skin action; and that the result of the rise of acidity from any such cause is a retention of uric acid.

Sir A. Garrod has shown it to be probable (*Gout and Rheumatic Gout*, p. 292) that the cartilages and ligaments of joints are, not only to some extent extra-vascular, but are also less alkaline than the blood and tissue fluids; so that we can at once see why uric acid, which is brought to these parts in blood less than usually alkaline, should be liable to be thrown out of solution and retained in these parts, beginning generally in the smaller peripheral joints where the circulation is least active; and the same reasoning thus applies to retention of uric acid in the joints as was previously used in regard to its retention in the liver and spleen (*Journal of Physiology*, pp. 213, 214), and we now see how a dose of iron, causing retention, or a rise of acidity however produced, may precipitate an attack of gout. Also, why a uric acid headache, or an attack of melancholia or mania, may promptly clear up on the supervention of a gout attack; for the rise in acidity, which drives uric acid out of the blood and cures either of the first two diseases, drives it into the joint and produces the podagra.

Exercise rather tends to prevent gout, because it increases skin activity, which again reduces acidity, and promotes elimination of uric acid; and the same skin activity, no doubt, explains why those who suffer from the uric acid headache have less of it in summer than in winter. With rheumatism we stand on less firm ground, because uric acid has not been proved to be in excess in the blood in this disease; but I hope I shall be excused for believing, at least till careful investigation has proved the contrary, that with the help of the knowledge we now possess about the excretion of uric acid, it may yet be found to be in excess in the blood in this disease.

The marked effect of salicylates in acute rheumatism seems to me a very strong point in favour of uric acid causation, and in favour of the disease being due to retention of uric acid by acids, the vicious circle being at once cut through by the salicylate, which permits its free excretion. There are, besides, important connections in the matter of family history, and in effects of drugs, meat, beer, weather, and exercise between gout and rheumatism, which are not to be easily disposed of, except on the theory of a similar causation.

Why uric acid should produce in one patient gout, in another rheumatism, in a third headache, and in a fourth epilepsy, may depend to some extent on the time during which it is in excess in the blood, and the amount of such excess; but, beyond the uric acid, there are no doubt peculiarities in the structure and function of nerves or vessels in each individual which determine the point on which the poison shall act most forcibly, and so to some extent the result it produces; but, in all these diseases alike, such regulation of diet, exercise, clothing, and climate as shall prevent high acidity from bringing about continued retention and accumulation of uric acid will serve to keep off attacks, even though the other factors remain in operation.

"WHAT IS A STONE IN THE BLADDER?" A REJOINDER.

BY SURGEON-MAJOR P. J. FREYER, M.A., M.D.,
Bengal Medical Service.

IN a paper by Sir Henry Thompson published in the *JOURNAL* of February 18th, entitled "What is a Stone in the Bladder?" the discussion of a question is introduced, which, till the subtle influence of Sir Henry's pen was brought to bear on it, could scarcely be regarded as admitting but one simple reply. Hitherto the universally accepted definition of stone in the bladder was any calculus formation, no matter of what size, which failed to pass out of the bladder spontaneously, and for the removal of which an operation was necessary. Sir Henry now lays down certain proposals which would have the effect of materially altering the nomenclature.

As Sir Henry has done me the honour of selecting my paper on "A Recent Series of One Hundred Cases of Operation for Stone in the Bladder without a Death" to "point a moral and adorn a tale," and has paraded before the eyes of the profession, in not very

complimentary terms, twelve instances of small stone therein contained, I hope it will not be considered presumptuous in me if I venture to criticise the position which he now assumes.

In order to illustrate this change of position I will take the liberty of referring to former utterances of Sir Henry on the same subject. So recently as 1884, in his lectures at the College of Surgeons, I find that the chief portion of one whole lecture was devoted to the consideration of the selfsame question, "What is a stone in the bladder?" And that his views were unaltered in 1885 is apparent from a letter which accompanied a copy of these *Lectures* with which Sir Henry, with his usual courtesy, was kind enough to present me.

Sir Henry's views on this question, then, are set forth in pages 128 to 134 of the *Lectures*. He calls attention to the liability to abuse of the term "operation for stone" by surgeons setting down as independent operations the repeated removal of small concretions by the lithotrite and aspirator, and points to Sir Benjamin Brodie as one who has strongly erred in this direction, no fewer than 8 of his 115 operations by lithotritry having occurred in one individual. Sir Henry himself thinks it justifiable to regard these repeated operations, under certain conditions which he lays down, as individual operations for stone, the force of which statement may be gathered from the fact that his 812 operations by all methods occurred in 716 individuals; there being, then, no fewer than 96 "recurrent" operations.

We now come to a passage in the *Lectures* of singular significance, when taken in connection with Sir Henry's recent paper, to which I would call particular attention. "I think," he writes, "it will be generally agreed that when a bladder is found for the first time to contain a calculus, whether urate, oxalate, or phosphate, which is too large to be expelled by the natural efforts, the crushing and removal of it (together with others, if such be present), so as to empty the bladder, necessarily constitutes an 'operation for the stone,' whether it or they be large or small." And then he adds, as his reason for this opinion, "If a small stone only is present, the patient is fortunate in having it detected, because it would inevitably have become large had it not been discovered, and the operation is then attended with a corresponding increase of risk." I presume that the principles here so clearly enunciated are those by which Sir Henry has been guided in his practice, and that if he has no calculi below 20 grains in weight in his collection this is due to the fact that he has not met with such, or has failed to detect them with the methods of diagnosis used by him. At any rate I submit that there can be no stronger justification for my recording my cases of small calculi than that conveyed in Sir Henry's own words.

From the somewhat vague and indefinite manner in which Sir Henry writes in his recent paper, whilst dealing with my cases, about the possibility of his increasing the number of his operations by recording all his cases of recurrent concretion in diseased bladders, the impression may be conveyed that I have utilised this method of augmenting my statistics. The amount of foundation for this impression may be gathered from the fact that the 100 operations referred to were performed in 100 different individuals, and that there were amongst them only two cases of recurrent stone. One man had undergone lithotomy in his childhood, and the other had had litholapaxy performed by another surgeon one year before. I may further mention that the 239 litholapaxies performed by me up to the present time have been done in 237 different individuals, the two cases of recurrent operation being fully described in my work on litholapaxy (*The Modern Treatment of Stone by Litholapaxy*, J. and A. Churchill, London, p. 57)—figures which I venture to think do not compare unfavourably with Sir Henry Thompson's, as regards the augmentation of any numbers from recurrent stone.

In my paper in this *JOURNAL* I have recorded in detail three out of the twelve cases to which Sir Henry takes exception, two of them having occurred in children; and as my right to return these cases as operations for stone in the bladder has been called in question, I hope I may be permitted to give details of two others from amongst them which may be regarded as typical examples of these small calculi.

CASE IX.—Samaroo, a hill-man, aged 48, was admitted to the Mussoorie Hospital on November 8th, 1886, with retention of urine. The patient was in great agony, the outline of the bladder being visible as high as the navel. He was relieved by catheterism by my assistant, Mr. J. McHale, who felt a small calculus at the neck of the bladder. The usual symptoms of stone had existed one year. I performed litholapaxy the same day. The patient being