the smell was so bad that I came home, and vomited many times. The next day I went again to the open drain, and the smell immediately made me vomit. I have been ill ever since."

The chief symptoms at the commencement were those of disordered digestion, with headache and vomiting, which at first occurred at intervals of some weeks, but much more frequently of late. He had lost three stone weight since Christmas, and those who knew him were much struck by his greatly altered appearance. His illness had compelled him to resign his appointment as surveyor. There had been no oedema, and the urine had never been observed to be high-coloured. During the last few weeks micturition had been frequent and copious. The urine now was of light colour, specific gravity 1012, and contained four grammes per litre of albumen. It deposited a dense but scanty whitish sediment, containing large granular and large hyaline casts, a few epithelial casts, and a few oily cells, appearances indicating a large white kidney, now undergoing contraction.

The heart's impulse was excessive, with doubling of the first sound and accentuation of the aortic second sound. All the superficial arteries, in particular the radials and the temporals, were tortuous and very tense. There was severe headache, with restlessness and frequent vomiting. The evidence of advanced and incurable degeneration of the kidneys was complete.

On March 12th, when I had another consultation with the gentlemen before mentioned, the urine was slightly blood-tinged. During the last few days the sight had become very dim, no doubt the result of albuminuric retinitis, but he was too ill and restless to bear an ophthalmoscopic examination, and I had to abandon the attempt to make one. I heard from my friend Mr. Percy Reynolds that, except an attack of nose-bleeding of some hours' duration, no fresh symptoms occurred until, after a few hours of coma without convulsions, the patient died on March 20th.

That the renal disease in this case was a direct result of the sewer poison to which the patient had been exposed about eighteen months before his death can scarcely be doubted. Up to that time he had been in good health, he was of strictly temperate habits, and he had not been exposed to cold, or to any other known cause of renal disease; and, on the other hand, he emphatically stated he had never been well since the sudden onset of the toxæmic symptoms.

It is noteworthy that none of the men engaged in repairing the drain, the emanations from which caused the fatal disease in our patient, were made ill in the performance of their disagreeable work, although they must have been much longer exposed to the foul air. This affords an illustration of the common observation that a virus which is sufficiently potent to excite fatal disease in one man may have no appreciable ill effect upon others who have been equally or even more exposed to its influence.

The most rapidly fatal case of poisoning by sewage virus that I have seen came under my care many years ago when I was physician to the Public Dispensary. I regret that I cannot find my notes of the case, but the main facts are that a man having been engaged for an hour or more in unstopping a blocked drain by stirring with a stick was speedily seized with vomiting, and the urine became scanty and deeply blood-tinged. There was rapidly increasing cerebral oppression, and death within forty-eight hours after exposure to the sewer poison. The kidneys were found much enlarged and of a deep chocolate colour, presenting exactly the appearance which is admirably represented in Dr. Bright's Plate 5 (*Medical Reports*, vol. i). On microscopic examination every uriniferous tube was found injected with extravasated blood.

METHYLENE AS AN ANÆSTHETIC.

By WILLIAM H. DAY, M.D.,

Physician to the Samaritan Hospital for Women and Children.

ATTENTION is just now being drawn to the action of methylene, and its claim to certain advantages over other anæsthetics. Having had with Sir Spencer Wells some considerable experience in the administration of methylene, and its subsequent effects during the period of convalescence, I am tempted to offer a few practical notes.

It is a mystery to me that an anæsthetic so safe and effectual as methylene should not have been more generally employed by the profession. A few deaths have been imputed to it, it is true, and occasional deaths will occur from any anæsthetic, however

carefully administered, but notwithstanding, I think, methylene occupies in point of safety a first class position. Some years ago the late Mr. Peter Squire asked me whether I considered methylene possessed any advantages over chloroform, and I replied in the following words, which will be found in his *Companion to the British Pharmacopæia*:

"It is less likely to cause vomiting, it is more agreeable to inhale, and there is less excitement preparatory to the stage of anæsthesia; rarely more than three or four drachms are required for an operation lasting half an hour, and consciousness returns in a few seconds after inhalation is discontinued."

My conviction of these merits remains unshaken. We should always bear in mind that all patients do not behave alike under an anæsthetic, whether it be ether, chloroform, or methylene. The young bear anæsthetics better than the middle-aged. The strong and the vigorous, and those of sanguine temperament, resist the action more than those who are nervous and phlegmatic; they are longer in getting under the influence of it, and it sometimes requires much patience and tact before complete muscular relaxation and tranquil sleep are established. If such patients have been intemperate the difficulty is all the greater, and I have felt on these occasions some anxiety, though I have never yet failed to narcotise them. Men do not take anæsthetics so well as women. If young and healthy there is more struggling and resistance, and after the middle period of life fatty infiltration of the heart is more common in the former, and they indulge to a greater extent in alcoholic stimulants. Children are quickly narcotised, and danger is scarcely ever present. In patients who lead temperate lives, and whose heart and lungs are healthy, almost any anæsthetic may be safely administered; it is when these organs are damaged that death from asphyxia or syncope sometimes ensues. In abdominal tumours of long standing, with a large quantity of peritoneal fluid, the diaphragm is pressed upwards, and the lower lobes of the lungs are so compressed that very little breathing space is left, and hence the danger of suffocation if the anæsthetic is not given sparingly and slowly. Slow administration is in itself a safeguard. In such cases the shoulders of the patient require to be well elevated, and the mask of the apparatus not brought so near the mouth as to exclude the free admission of atmospheric air into the lungs. In these cases of pulmonary oppression the patient may become livid after a few inhalations, and be seized with a convulsive, choking cough, which embarrasses the operator, but as the ascitic fluid escapes, and the tumour is removed, freer respiration is established, and then the remedy can be pushed to full narcotism, more particularly when the abdominal distension is of recent date. It is in disregard of this warning that fatal consequences may result.

When the abdomen is distended with fifteen or sixteen quarts of fluid, and contains a multilocular cyst, weighing eight or nine pounds in addition, the umbilicus may reach a higher level than the nose of the patient as she lies on the operating table. In such a case slow and cautious administration is demanded, and as I have before said some of the fluid must be withdrawn before the patient can take a full inspiration. At this stage we cannot push the methylene beyond partial anæsthesia.

When mucus accumulates in the mouth, I am in the habit of twisting one end of a pocket-handkerchief round the forefinger of the right hand, and wiping it away. This simple practice prevents the mucus from collecting at the back of the pharynx, whilst it facilitates respiration, and prevents the choking cough.

When the pulmonary organs have been long pressed upon, or the heart displaced, bronchial irritation sometimes comes on a few days after operation, as the lungs undergo expansion, and the pressure is removed. This is all the more likely if a considerable quantity of the anæsthetic has been administered. Such complications are rare in recent abdominal distension, but in old standing cases where the lower ribs are bulged outwards like a fan this risk is frequent, and ought to be calculated upon. In these cases the heart becomes weak; it is often displaced upwards, and prone to take on degenerative change.

As to the existence of a cardiac murmur, undue importance may be attached to it. It is not always easy to diagnose a functional from an organic murmur, but where the pulmonary circulation is free, I never hesitate to recommend methylene as an anæsthetic. The pressure resulting from a large abdominal tumour may give rise to a soft cardiac murmur, which means nothing at all, if the muscular structure of the heart is vigorous enough to overcome any distension of its cavities, but we must not overlook the fact that long-standing upward pressure tends to invite feebleness of

the heart's action, thin walls, and degenerative changes. Sudden death at the time of operation, or soon after it, has to be reckoned with.

I have safely given bichloride of methylene to patients suffering from heart disease on many occasions; in dilatation and weakness of the organ, and in mitral change with hypertrophy. When there has been considerable distension of the right and left cavities of the heart from valvular defects I have never hesitated to use methylene if an operation has been urgent, even if the cardiac trouble has considerably impaired the pulmonary circulation. I am doubtful whether there is any other anæsthetic to supersede methylene in this condition. I have ascertained, before commencing the administration, that the heart was weak, or its valves damaged, and yet, in spite of this, the pulse has frequently improved in quality to the surprise of bystanders, and the patient has looked a far better colour at the close of a tedious operation than at the commencement. The bichloride has acted as a cardiac stimulant throughout. Exceptions are sometimes met with, and in these cases I invariably resort to brandy during the administration, never pushing the methylene to such an extent as to interfere with swallowing. I have given, during a long and desperate operation, as much as six ounces of brandy, and I am convinced that this precautionary measure has carried the patient safely through. I never trouble myself about the pulse; the countenance and the respiration are my guides. No hard-and-fast lines can be drawn. One patient can be sent into tranquil sleep with three drachms for half an hour, and a very exceptional case will require three times the quantity. Twelve drachms is the largest quantity I have ever given for an operation lasting one hour, and I have no remembrance of Sir Spencer Wells being a longer time than this over his worst operations.

I have known many cases of pleuritic effusion in ovarian disease, and when recent and serous I have advised the abdominal section before tapping the thorax, and in every such case where there has been enough breathing space left I have not been anxious. The effused fluid has disappeared from the chest without operative interference.

I have administered methylene with the best effects in cases of advanced phthisis with a large cavity in one lung, and softening going on in the other.

I do not think I have over-estimated the value of methylene, as in 1,230 cases I have not met with anything to alarm me. It is the most harmless of anæsthetics; it has never disappointed me, and I feel sure that those who will give it a fair trial will record a similar experience to my own.

It is not in healthy individuals that danger is to be apprehended from the employment of an anæsthetic. It is in the diseased and feeble, who have been reduced by pain, sleepless nights, and long suffering. In such cases as these Dr. Frederic Hewitt advocates the anhydrous ethylic ether, or the A. C. E. mixture. ("On the Production of Anæsthetic Sleep in Patients suffering from Grave Constitutional Disease," *The Lancet*, May 26th, 1888.)

Dr. Richardson introduced methylene bichloride in 1868-9, on a distinct principle—namely, that all the chloride anæsthetic series are excitants by reason of the chlorine. He therefore considered that by reducing the chlorine he reduced the danger. He used first methyl chloride (CH_3Cl); but, this being a gas, he found a difficulty in using it, because, unless it were given in the pure form from a large reservoir, when he found it a most perfect anæsthetic, it could only be given from a fluid in which it was condensed, and from which it escaped, producing extreme cold. He proceeded next to methylene bichloride (CH_2Cl_2), or one equivalent less of chlorine than chloroform (CHCl_3); and, after getting it in the pure state and inhaling it himself as a trial, he administered it successfully to a patient, Sir Spencer Wells being the operator. Dr. Richardson's argument has throughout been that methylene is safer than chloroform, because it is a lighter body, is more easily diffused, is more equable in action, and is less of an excitant. My experience confirms these conclusions to the letter.

ACUTE CIRCUMSCRIBED OEDEMA.—Dr. Falcone (*Monat. f. Prakt. Derm.*, 1888, p. 291) relates a case of acute circumscribed oedema in which the patient's grandfather had suffered from a similar affection. Dr. Riehl (*Ibid.*, p. 444) relates a case in which the acute oedema formed behind the eyeball. The eyeball protruded and there was oedematous swelling of the eyelid and conjunctiva. The swelling lasted for twelve hours. There were two relapses. Meynert stated that a similar process of acute circumscribed oedema sometimes occurred in the brain, leading to death.

POST-NASAL GROWTHS AND THEIR RELATION TO DISEASES OF THE MIDDLE EAR.¹

By ADOLF BRONNER, M.D.,

Surgeon to the Bradford Eye and Ear Hospital.

MEYER, of Copenhagen, was one of the first to draw attention to the frequent occurrence and great importance of post-nasal growths (*Hospitals Tidende*, November 4th and 11th, 1867, and *Archiv für Ohrenheilkunde*, 1873). Several authors divide post-nasal growths into two diseases, that is, adenoid vegetations and hypertrophy of the pharyngeal tonsil. There does, however, not seem to be any physiological difference between the two affections. They are simply two forms of one disease, that is, local hypertrophy of the normal pharyngeal tonsil or retinoid tissue, situated at the vault of the naso-pharynx. If the hypertrophy is regular and lobular, it is called hypertrophy of the pharyngeal tonsil; if irregular, adenoid vegetations or lymphoid papillomata. Of course in many cases the surrounding retinoid tissue is also hypertrophied, but rarely to such an extent as to be of much importance. Many authors still seem to believe that the growths grow from the posterior wall and sides of the naso-pharynx. That this is not the case has been proved by the *post-mortem* examinations of Trautman, of Berlin (*Anatomische Pathologische und Klinische Studien über Hypertrophie der Rachen tonsille*).

The symptoms of post-nasal growths are very characteristic. The child (as children are mostly affected) is generally pale and weakly, with in-drawn thorax and pinched nose. It keeps its mouth open, more or less, which gives it a peculiar and stupid expression. The lips are swollen and thick; often also the eyelids. The child sleeps with its mouth open, and invariably snores at night. There is a peculiar nasal twang, very similar to the speech in paresis of the soft palate. The child cannot breathe through its nose, and is subject to attacks of bronchitis. In nearly every case there is distinct history of recurrent disease of the middle ear. The growths can be easily felt by introducing the finger behind the soft palate. They can in most cases be distinctly seen by anterior and posterior rhinoscopy.

The treatment of post-nasal growths can of course only be a local one. I now invariably remove them with the ring-shaped curette, as recommended by Hartman, of Berlin. It removes the growth very readily, the operation is not very painful, there is no danger of wounding any important structures, and, what seems to me to be of very great importance, the operation can be performed without an anæsthetic. The curette is similar in shape to the curette of Lang or Dalby, but the upper edge of the ring is flat instead of round (*vide* diagram).

The head and arms of the patient are firmly held by an assistant; the tongue is depressed, and the bent end of the curette is passed behind the soft palate. The flat end (a) is then pressed well against the vault of the naso-pharynx, and firmly drawn from one side to the other, thus cutting, or rather slicing, off the growths.

In many hospitals it still seems to be customary to use an anæsthetic when the growths are removed. By operating in the manner I have suggested an anæsthetic is hardly ever necessary, and the removal of post-nasal growths is reduced to a minor operation which can be undertaken at any time or at any place, without the slightest risk and in a very short time.

Post-nasal growths are to be met with mostly in children, and are undoubtedly much more common than is generally supposed. Meyer examined 2,000 school-children at Copenhagen, and found that 20, or 1 per cent., showed distinct symptoms of the growths. He also examined 700 children of the Orphanage at Haverstock Hill, and found that 13, or nearly 2 per cent., had post-nasal growths. Professor Doyen saw 4,000 children at

Haverstock Hill, and found that 13, or nearly 2 per cent., had post-nasal growths. Professor Doyen saw 4,000 children at

¹ Abstract of Paper read before the Yorkshire Branch of the British Medical Association, at Leeds.