cases. Not a few of these were removed many years ago with Clover's aspirator, which I was in the habit of using to discover and remove these small bodies. Thus, among many other cases in my collection is one, in 1872, of 500 small concretions, weighing together a drachm or more, all thus removed, but never recorded as an operation for stone. It was an operation for gravel, a proceeding which Dr. Freyer appears to ignore; for one of the cases in which he has merely washed out a particle weighing 2 grains, is actually recorded as an "operation for stone!" He says, and rightly, that there is no hard-and-fast line in the matter of weight between stone and concretion. Then, if a 2-grain stone be admissible, why not a 1-grain, or a $\frac{1}{2}$ -grain stone, or less, if the patient fails to pass it naturally? Is there henceforth to be no operation for gravel or concretion, in order that stone operations may be multiplied?

But Dr. Freyer quotes a definition of stone I formerly made, and says I have recently modified it. That is precisely the course I desired to take, and intimated it in my lecture in autumn 1887. When making the first, I confess I never supposed any English surgeon was likely so to construe it as to regard a tiny piece of gravel as "a stone in the bladder." The intention which prompted the later remarks—and I had a very distinct one—pointed quite in a different direction. Hence it was a matter of astonishment and severe disappointment to me, who respected and admired Dr. Freyer's skill and energy, to learn that he condescended to record these trifles in terms which have never been used before. For I have the means of knowing that such a precedent is opening the door to practice outside our profession, of a very questionable kind.

To close a paper already too long, I think there is little doubt, judging from the opinions which have reached me here, and from abroad, that the common sense of the profession will exclude a trifling concretion of 2 or 3 grains from taking rank under the well-known and time-hallowed term a "stone in the bladder." A hard-and-fast line being impossible, most surgeons will probably agree that a product at least four or five times that weight might be held to constitute the smallest size to which that title should be applied. In my own collection there is nothing less than 20 grains.

To that common sense I leave the verdict, feeling more strongly than ever, after reading Dr. Freyer's second communication, how necessary it was to raise this question. I did so with great reluctance, and with a feeling of infinite regret that a paper of Dr. Freyer's should have furnished the occasion.

PERFORATION OF THE PERINEUM DURING LABOUR.

By JAMES FERGUSON, M.B., C.M.,
Perth.

CASE I.—On March 13th, 1887, I attended Mrs. J., aged 25, in her first confinement. On arrival I found a well-advanced second stage, with an occipito-anterior position of the head. The patient's condition was good, the pains frequent and strong, and everything seemed to favour a normal termination. After a short time, with my finger over the fourchette, I was conscious of a sudden considerable movement of the child's head, followed by a profuse rush of fluid, both felt and heard, all this coinciding with the height of a pain. Immediately exposing the maternal parts, I was impressed by the enormous distension of the sacral portion of the pelvic floor, including the anterior wall of the rectum, and just in front of the anus noted a nearly triangular space of dark colour, which on closer examination proved to be nothing else than a portion of the child's scalp applied to a tear in the perineum.

Finding the margin of the vulva lying quite flaccid and free from pressure, my first thought was to hook it backwards, so as to bring it into the axis of expulsion, meanwhile detaining the head in the pelvis by my own efforts, the patient being ordered to desist from hers. But the uterus seemed only aroused to increased energy by the measures adopted, and, in brief, within a few seconds a male child of average proportions passed wholly through the abnormal outlet. The placenta was suddenly extruded by the same opening in a few minutes. Examination showed the rupture to extend through the external sphincter ani, while anteriorly almost an inch of skin separated the perforation from

the vulvar orifice; the posterior vaginal wall was torn for an inch and a half upwards.

The injured parts were smeared with eucalypto-vaseline (1 to 9) as the only antiseptic at hand, and then appropriately brought together with carbolised catgut. Two sutures in the vaginal wound and half a dozen in the perineum were used. Opium was prescribed, to favour constipation rather than to soothe the pain, which was not complained of after the suturing, and the subsequent treatment fell to my friend, Dr. Sinclair Kennedy, in whose detention I had been temporarily acting in the case. Antiseptic injections and applications were employed, and the bowels not moved till the ninth day. By this time the wound had healed, for the most part as if by first intention, and the patient described herself as only uncomfortable by being kept in bed. The temperature never passed 99° F. By the twenty-first day the patient could walk, and alleged she felt no pain.

Meanwhile, Dr. Kennedy had ascertained an abnormality of the pelvis which quite accounted for the abnormal labour, and by his kind request I was allowed to confirm his observation. The chief facts were an unusual depth of the symphysis pubis and undue narrowness of the pubic arch. The vertical measurement of the former, allowing freely for its soft coverings, was fully three inches and a half, an excess of two inches over that commonly assigned to the female symphysis. Exact measurements of the arch were not possible, but the male type was very perceptibly presented, an acute angle taking the place of the rounded arch, and the pillars approximating in a corresponding manner.

CASE II.—Called to Mrs. P. in her third confinement on April 2nd, 1888, I found the head approaching closely to the perineum. The finger also detected a hard wedge of tissue extending upwards and backwards from the vulva, which suggested inquiry about former labours. My patient told me that at her previous delivery forceps were used by a doctor well-known in obstetric work in the town she then lived in, and that, though nothing was said about a tear at the time, for weeks after she began to move about "something like a piece of flesh hung down," and caused much pain, especially when sitting. As the child's head distended the perineum, the above-mentioned wedge persisted in passing doggedly forwards, refusing to share in the dilatation, and carrying the labia before it. Instructing the patient to use no voluntary efforts, and to cry out during the pains, I pressed the head backwards and upwards towards the sacrum, and attempted to dilate the vulva by stretching with two fingers over the fourchette (an old method recently revived, *vide* JOURNAL, 1886, vol. i, p. 12). The head was then allowed to come down with a minimum of pressure, the hand trying to guide it forward past the obstruction. Despite these methods, however, the hard body still advanced before the child, and I was alarmed to detect a sudden yielding of the perineum just in front of the anus during a pain. Recognising only danger in further delay, I pushed the head clear away from the pelvic floor, and with some difficulty—continuous uterine effort having been provoked—adjusted the forceps and delivered by the vulva, its margins yielding freely. The perineal wound was afterwards found to be almost three-quarters of an inch in length, communicating with the vagina, the sphincter escaping by a thin cord of skin. The wedge behind the vulva consisted of cicatricial tissue, passing about an inch upwards and obliquely backwards, and, strange to remark, bore trace of only the slightest laceration. A deep and superficial stitch having been inserted, and weak lotions of Condy used, the patient made an unretarded recovery, the wound healing apparently by first intention.

The treatment of perineal lacerations being a subject on which conflicting opinions prevail, the last-mentioned case may be regarded as evidence of some value on one side of the question. It is clear that a not inconsiderable tear at the time of the second confinement was ignored, and was the cause of much suffering for weeks afterwards. Not only so, but the closer apposition of the torn surfaces secured by timely stitching would have saved the patient suffering at her third confinement, and averted the risk of an "accident" of a serious nature, and one which would have certainly occurred had not professional assistance been at hand. May I not also ask, supposing more serious injury had occurred, despite attempts to avert it, on whom would any possible discredit have fallen, the first or the second practitioner?

As to the treatment appropriate to those cases of central rupture which are recognised before the birth of the child, Charpentier has advised that, to save the sphincter ani from danger, the tear should be carried forward by a cut into the vagina.

The above case proves, however, that, in certain instances at least, we have in the forceps an aid that obviates any extension of wounded surface.

AMAUROSIS AND STRABISMUS FROM ASCARIS LUMBRICOIDES.

By JABEZ HOGG, M.R.C.S.,

Consulting Surgeon, Royal Westminster Ophthalmic Hospital.

FROM a record of cases, carefully tabulated and extending over many years, I find that strabismus in young children is more frequently due to the irritation of intestinal worms than is generally stated in manuals of eye diseases. It is, however, quite an uncommon thing to find reflex amaurosis and strabismus associated in one and the same person, and arising from the same cause—the presence of worms. In the case I am about to narrate there is one other feature which renders it of more than ordinary interest. My little patient, barely three years old, was, when she came under my care, the intermediary hostess of three kinds of entozoa—namely: oxyurides (the little thread-worm), tænia (tape-worm), and ascaris lumbricoides (round worm). On searching through the medical journals, I have not met with a parallel case, and I have discovered only a very few cases of reflex amaurosis from worms, one of which, narrated by Dr. Burgers, was published in the BRITISH MEDICAL JOURNAL of 1862. A still smaller number of cases have also been reported in which, it is said, death has resulted from the perforation of the intestine by ascaris lumbricoides. It is said that the three kinds of worms named by me will not exist together. It is difficult enough to account for their presence in the intestine of so young a child, and belonging as she did to the middle class of society. Polluted drinking-water is the carrier of filarise, and this might have been unwittingly used by the nurse to mix with the milk which formed the staple food of the child. It is known, too, that the ova and embryos of parasitic worms will occasionally undergo a passive migration; that is, direct from intestine to intestine, and without passing through an intermediary host. It is quite possible, therefore, for embryos to be conveyed in the blood of the mother to the foetus *in utero*, and not reach their final stage until some months after the birth of the child. In my case, there was no history of the kind; the mother is quite certain she never was afflicted with worms, but, curiously enough, on going back another generation, the grandmother had passed several tape-worms after she was married. When Mrs. D. consulted me about her child's sight, the ordinary symptoms of worms were mostly absent. Her sleep was not disturbed; she had no nasal irritation; no craving for food of any kind; disliked meat, and had a small and capricious appetite; her daily food consisted for the most part of milk and bread and light sweet puddings. She was spare in body, weak in her legs, and her movements were consequently slow, passing from one chair to another by touch. This the mother attributed to the loss of sight, which had now persisted for more than a year. Her speech was very defective; her vocabulary consisting of a few unintelligible sounds, just sufficient to make her wants known to her nurse, and no more. When placed in front of a strong light, the eyeballs rapidly moved upwards, the irides being concealed by the upper lid. From this cause, and that of the persistent squint, I was quite unable to make any ophthalmoscopic examination. I, nevertheless, came to the conclusion, not shared in by anyone who had previously seen the child, that the strabismus at all events was symptomatic of worms. I accordingly prescribed anthelmintics, commencing with scammony and jalap, which, after the administration of a second powder, gave the first intimation of the correctness of my diagnosis, as it brought away a large number of ascarides. My next remedy, filix mas, dislodged a portion only of a tape-worm, eighteen inches long; other portions came away some days afterwards; and I next prescribed three-grain doses of santonin at bedtime, to be followed by a full dose of castor-oil the following morning. This treatment proved very effectual; the first powder brought away five or six large lumbrici. In the course of the following fortnight, five or six and twenty round worms, measuring from four to eight inches in length, were expelled; the relief obtained by the little patient was most gratifying to behold. Two months have elapsed. She is recovering her sight, and the strabismus has entirely disappeared. Her bodily health is quite changed for the better; she is strong on her feet, can walk a considerable distance, and has gained in intelligence and in speech

—is, in fact, an altered child. From the large size and strength of the female lumbrici, I should not have been surprised if perforation of the intestine, in one so young and delicate, had taken place before I succeeded in dislodging all the worms.

CLINICAL LECTURE ON CERTAIN OPERATIONS ON THE EYE.

By C. BELL TAYLOR, F.R.C.S. AND M.D. EDIN.,
Surgeon to the Nottingham and Midland Eye Infirmary.

GENTLEMEN,—Of the operations you have seen, there are three only to which I wish to call your attention. The first of these was the case of stenosis of the lachrymal duct, which was dilated without slitting the punctum. This is easily accomplished with the probe which I have had modelled by Messrs. Weiss, on the pattern of Weber's canaliculus knife (Fig. 1), and the results, as you may imagine, are much more satisfactory than when the sucker-like function of the punctum is permanently destroyed, as in the usual operation, by conversion into an open channel.

Fig. 1.

The second was an exceedingly obstinate case of trichiasis and entropion specially affecting the upper lid. The patient, aged 49, who had travelled a considerable distance in the hope of relief, has been twice operated on elsewhere without lasting benefit, and was so blind on admission that she could not get about alone. In these cases I make a perpendicular incision at each extremity of the lid, evert it, and connect the two incisions by a deep cut on the inside right down to the cuticle, just within the roots of the lashes, as in Burrow's operation. (Fig. 2.) I then take a piece of skin from the upper lid, and, retaining a broad pedicle, transplant it into the wound thus made. The approximation of the edges of the gap in the upper lid by sutures everts the lashes, and the transplanted skin effectually prevents any return to their abnormal situation.

Fig. 2.

I devised this operation some years ago, and have never known it fail to cure this obstinate and most distressing affection. The flap, prior to transplantation, is outlined in the accompanying woodcut.

The third case was one of cataract, in which the lens was, as you saw, removed without excision of any portion of the iris; and, in calling your attention to these patients, permit me to remind you that the ideal of success in cataract extraction, the success without restriction, the success which has nothing to fear for the future is in language which I quote from Suarez de Mendoza:—"Extraire tout le cristallin opaque sans perdre une goutte d'humeur vitrée, obtenir une cicatrisation régulière sans enclavement iridien ni capsulaire, une pupille nette et sans la moindre opacité, en conservant l'iris mobile et libre de toute synéchie."

Such, gentlemen, were the indications which have been fulfilled in every particular, under singularly adverse circumstances, in the patient you have just seen, who is 73 years of age, and who, as you have noticed, presents that peculiar complexion, the result of heart disease and impeded respiration, which is known as "blue skin." She is also very deaf, and strangely irritable. I operated upon her here on March 18th; and on the 20th, two days later, owing to some dispute with the nurse, she