rigid by means of a broom-handle or piece of iron tubing. Even that, however, is unnecessary, as with care the body can be placed at once without difficulty into the coffin.

The method is one which I believe will be much more frequently employed in the United Kingdom when it becomes more fully known than is the case at present.

The chisels and forceps are now made by Messrs. Weiss and Sons, Oxford Street, London, to whom I am indebted for the accompanying illustrations.

A CASE OF HYSTERO-EPILEPSY OF TWENTY YEARS' DURATION, TREATED BY REMOVAL OF THE UTERINE APPENDAGES.¹

By FRANCIS IMLACH, M.D.

OF the treatment of various neurotic pathological conditions by removal of the uterine appendages I have little knowledge and less belief. By a curious coincidence, two entirely different motives appear to have suggested this much discussed surgical procedure almost simultaneously to men of different bent of mind. One motive may be termed transcendental, hypothetical, doubtful; it is determined by the belief that certain untoward neurotic manifestations depend upon morbid ovarian activity, and may be cured, and can only be cured, by removal of the ovaries. The other motive is commonplace, obvious, sure; namely, that disturbing, disabling and often more dangerous and even fatal conditions, due to acute or chronic inflammatory diseases of the uterine appendages can only be relieved, and ought to be relieved, by removal of the diseased parts.

It is by the latter motive I have been influenced, and the greater my experience becomes, the more convinced I grow of its soundness.

While in Tenby, however, at the new year, Mr. Beamish Hamilton asked me to see a hystero-epileptic patient, who might, he thought, be benefited by surgical treatment, and whose record may be useful at the present time. She was aged 40, and had been bedridden for the past nine years. At 20 or 21 years of age she became subject, she said, to epileptic fits, which had continued to the time I saw her. Indeed she appeared to have become bedridden almost as much from the increasing severity and frequency of the fits as from pain; her fits generally recurred three or four times in the day, and it was rare to pass a week with so few as four or five fits. She was always unconscious for ten or twenty minutes during a fit, but did not sleep heavily after it. Having been in five different hospitals, special and general, for periods varying from six weeks to four months, without relief, she had given up expectation of cure, and Mr. Hamilton informed me that bromides and other nervines appeared entirely useless.

It was plain, upon examination, that the Fallopian tubes were greatly distended, and I therefore advised their removal. Menstruation was very profuse, and there was great pain; my opinion was formed regardless of the neurotic condition which now, however, induces me to record the case in detail. At 28 years of age she married, and had a child, which died two years later. I have no record of the condition of the uterine appendages previous to this.

As I was unable to remain at Tenby, she was brought to Liverpool, entered my private hospital on January 10th, and four days later I removed her uterine appendages. It was a case of double hydrosalpinx, with extremely cirrhotic and shrunken ovaries. The right tube contained half a pint of serous, shreddy fluid, and the left about half an ounce. The left ovary was no larger than a dried split pea, and the right ovary about double that size (specimens shown). While being anaesthetised she had a prolonged and rather alarming fit: she became suddenly unconscious, her lips turned blue, respiration was arrested, and her back became strongly arched. An hour after the operation, which lasted only a few minutes, she had a similar attack, and, to our disappointment, a week later she had another, but no more.

On February 15th, being able to walk alone, she returned to Tenby, and the following letter was received on March 18th from her medical attendant, Mr. Hamilton:

"Tenby, March 17th, 1888. Dear Dr. Imlach,—In reply to your letter of queries about Mrs. B., she has been practically bed-

ridden for the past twelve years, since her 29th year, and has passed through the hands and treatment of many medical men, without getting the smallest benefit. Eight years ago she was kept upon her back for three months in the — Infirmary, and treated with pessaries, etc.; twelve months ago she was treated in the — Hospital with stem pessaries, etc., but without relief. For the past twenty-one years she has suffered from epileptoid seizures, accompanied with marked arching of the spine at the commencement of the attack, insensible for periods varying from ten to twenty-five minutes, for which she came under my treatment about eight months back; but nothing I could do seemed to mitigate them, and they were increasing in frequency and duration, when she had the good fortune to fall into your hands. Since her return home there have been only two fits of any severity, and they seem now to be amenable to bromides. Seven weeks from the day of operation she had a spurious menstrual flow, which lasted for five days. This made her nervous, and I think was the cause of the fits mentioned. She is eating well, her back-pain is gone, and I am only waiting for the fine weather to send her out of doors.—In great haste, yours very faithfully, BEAMISH HAMILTON."

As a primary result of the operation this is very encouraging. Had this treatment been carried out years ago, the patient would probably have been a useful housewife, free from pain, and a fit companion to her husband, in place of a hopeless and expensive invalid.

PRACTICAL ANTHROPOMETRY.

By CHARLES ROBERTS, F.R.C.S.

I HOPE that the suggestion thrown out by Dr. Stone in his Harveian Oration that "medical physics" shall form part of the science to be taught in the proposed laboratory for "scientific purposes" to be attached to the new Examination Hall of the Royal Colleges, will be acted on, and that practical anthropometry will form part of that important branch of the medical man's education. Anthropometry, or "man-measurement," strictly speaking, is a branch of human anatomy, and deals with the body as a living organism, while the anatomy taught in our medical schools deals only with the structural arrangement of its several parts; and it is obvious that if our students were properly taught, their first lesson should be to acquire a knowledge of the physical characters of the whole body before proceeding to study its elementary constituents. It is necessary, moreover, that, in order to practise his profession with satisfaction to himself, his patients, and, in public medicine, to the public, that the medical man should be acquainted with the normal condition of the body, and the variations in its conformation which result from race, sex, age, and sanitary surroundings. Besides being a branch of anatomy, anthropometry is a valuable means of diagnosis, not only for defining diseased conditions, but the influence of external agents on the growth and development of the whole body, or some of its organs. Indeed, in its widest sense diagnosis is anthropometry, for all our attempts at differentiation of diseases are measurements of either time or space, and all our instruments, whether for pathological or physiological investigation, are anthropometric instruments when applied to the investigation of the anatomy and functions of the human body. "The indispensable part of the experimental observation of physical facts is the measurement of quantities," says M. Henri Martin in his *Physiques*. "All physical objects are extended, and consequently measurable, though they may not always be measurable by the methods and instruments we possess; and all physical phenomena take place in periods of time susceptible of measurement, although some of the measurements are imperceptible by our senses. The right method as regards all these phenomena," he goes on to say, "is that taught by Galileo, to measure all that is measurable, and to endeavour to render measurable all that is not directly so." But while claiming for anthropometry this relation to medical physics, and thereby securing its recognition as an essential element of that much neglected branch of medical study, it is desirable in these days of the division of labour to confine the sphere of anthropometry to the study of the physical characters or qualities of the body, and to leave the subjects which have hitherto been included in the provinces of physiology and pathology, such as the functions of the senses and other obscure physico-psychological phenomena, which, though measurable by suitable instruments, can only be properly studied by persons

¹ Communicated to the North Wales Branch, March 20th, 1888.

who are acquainted with the minute anatomy of the body and of the various imperfectly, as yet, understood functions of the nervous system.

I mention this here because I fear that practical anthropometry, as generally understood on the Continent and America, as well as in this country, is likely to suffer from a series of ingenious inventions manufactured and sold by the Cambridge Scientific Instrument Company, and advertised in the columns of the JOURNAL as anthropometric apparatus. Now the instruments required for recording the purely physical qualities of the body, that is to say, the stature, the length, breadth, circumference, and thickness of different parts of the trunk and limbs; the weight, strength, and breathing capacity; and of the complexion, as distinguished by the colour of the hair, eyes, and skin, are few in number and inexpensive, and consist of a pair of ordinary callipers, two or three slight modifications of the shoemaker's rule, a measuring rod and tape, a weighing machine, one or two dynamometers, and a spirometer. Not only does the catalogue of the Cambridge Company contain instruments of this description, but many others for testing the keenness of eyesight and appreciation of colour (not the same as have been perfected by ophthalmic practitioners); keenness of hearing, and appreciation of musical pitch; and the numerous qualities of the senses and the mind which are largely, if not entirely, the result of education and individual experience, such as "judgment of the eye [mind?] as regards squareness and angular division; judgment of the eye [mind?] in estimating the division of a line; reaction time to sound and sight; appreciation of slight differences of weight." These are doubtless interesting subjects of inquiry, but they are not anthropometric inquiries in the sense in which the word is generally understood, and to which I think it is desirable to restrict it.

I am inclined to think, moreover, that Mr. F. Galton, who has designed these instruments, does not see what complicated problems he is asking us to investigate under such an apparently simple question, as, for instance, "the judgment of the eye as regards squareness." An instrument for this purpose might serve in a rough way to distinguish between an artist and a barrister, an artisan and an agricultural labourer; but between two persons of equal training and experience the test would be one of the different amount of astigmatism in the two persons under examination. Similar objections occur to me with regard to some of the other instruments in the catalogue; but my object is not to criticise them, but rather to point out to the members of my own profession the proper sphere and the practical utility of anthropometry as generally understood and as was understood by Quételet, who invented the word to cover the series of inquiries which he carried out on the human body, and which are recorded in his work entitled *Anthropométrie ou Mesure des Différentes Facultés de l'Homme* (1870).

When I published my *Manual of Anthropometry*, in 1879, I was soundly rated by the *Lancet* for making such free use of Quételet's observations, but it would be just as reasonable to rate Quain and Gray for appropriating the labours of previous anatomists. Although anthropometry has existed in a slipshod manner for many centuries among artists and sculptors, it was Quételet who first reduced it to a scientific form and applied it to scientific uses; and it is desirable, if any progress is to be made in this as in other sciences—and especially in those which, like anthropometry, so largely depend on collective investigation—that changes in its methods of procedure should not be made without very grave reasons.

Unfortunately for the stability of the science of anthropometry, Quételet was a mathematician and an artist, and not an anatomist, and he gives us in his book only the results of his anthropometric observations, and not the method by which he obtained them. He dealt with the subject as if he had exhausted it, as indeed, for artistic purposes, he very nearly has done so. It is when we come to try to apply his method to the distinctions of the races of man, and to medical, sanitary, and other scientific uses, that it fails us, simply because he has not defined the conditions under which his observations (mostly made by his medical friends) were made, and it was for the purpose of defining these conditions that my essay was published, and this only after a translation of Quételet's work, with the necessary notes, was refused by the Sydenham Society. I mention all this because I wish to impress on everyone who takes up the subject of anthropometry that it is better to get some results by methods which have been found to work fairly well, than to be constantly tinkering the methods and obtaining no results and disheartening others of a less critical turn of mind. *Ars longa,*

vita brevis is a motto especially applicable to this subject. While we are inventing instruments and illustrating mathematical problems by our miserably small collection of anthropometric observations, the French are applying the science in its simplicity to the identification of criminals, the Germans to the differentiation of races, and the Americans to the results of gymnastic and athletic exercises, and to all of them anthropometry has become serious everyday duty and study—as serious, to compare large things with small ones, as the study of micro-organisms is with ourselves. Sufficient work has, however, been done in this country to form a basis for more and better work in future, and sufficient has been done, especially in differentiating sanitary conditions such as the influence of occupations, food, social status, etc., and of the racial elements of the population, to promise great results if the methods of anthropometry we already possess were recognised, and their practice properly encouraged by the Colleges of Physicians and Surgeons and the medical schools.

Simple and easy as anthropometry is in practice, its application to the solution of racial, physiological, and sanitary questions is so wide and comprehensive, that it is hardly within the power of individuals to carry it out. Quételet first described the method but he shrank from the labour it involved, and instead of measuring men, women, and children by thousands he measured them by tens, and so missed seeing some important facts in relation to sexual and racial differences which have since been discovered. Dr. Beddoe has spent a lifetime in collecting anthropometric data for differentiating the racial elements of our British population. Professor Bowditch has with much labour discovered by anthropometric observation the difference in the rate of growth of the two sexes.

Mr. F. Galton has illustrated in a similar way some important facts relating to the hereditary transmission of physical characters, and I have endeavoured to establish standards for comparison, and to define some of the results of the "conditions of life" on the development of the body, while in another field Hutchinson's investigations have become classical.

But not only does all this work want revising and extending, we want also a profession which can understand and appreciate its value, which we have not got at present, and which, I fear, we never shall have till the heads of the profession, as represented by the colleges and schools, recognises its importance as equal at least to those of comparative anatomy, chemistry, and botany, and, indeed, as superior to them in its educational value, the mathematical and statistical accuracy which anthropometry demands being of supreme importance as a check on the imagination of the student, and a suitable training for other investigations necessary to the progress of medicine and surgery in our day.

HOW TO REFORM THE DENTAL DEPARTMENTS OF OUR HOSPITALS.

By F. NEWLAND PEDLEY, F.R.C.S. AND L.D.S.,
Dental Surgeon and Lecturer on Dental Surgery to Guy's Hospital.

THIS is a problem I took in hand at Guy's Hospital, and now that the experiment has resulted well and nearly two years have been allowed to lapse as a time-test, I bring the matter forward in the hope of inducing other hospitals and other dentists to adopt necessary changes.

I do not advocate that each medical school should be a school for dental students; at present this is impossible. Let dental students seek special training at special hospitals, but those who hold the dental surgeoncies to hospitals should show their operations and the scope of their subject as the other specialists do. The status of dental surgery, though improved, is far from what we would see it, and from what we think it deserves. What can be more damaging to our profession, which, above all, rightly claims to be a branch of surgery, than the lamentable fact that medical men so frequently feel obliged to admit utter ignorance of the subject? Hence it becomes a frequent source of complaint amongst dentists that the importance and value of their work is inadequately appreciated, and the medical fraternity is blamed for possessing little or no knowledge about teeth. The work done at the dental hospitals is admirable, but busy medical practices cannot give the time to attend there, and they go into practice with the idea that the extraction of teeth and their replacement artificially form the limit of our practice and science.

The fault lies with those dentists connected with hospitals who