changes which, I think, take place when the abdominal cavity has been filled with fluid or an ovarian tumour.

The result of my treatment in this case will encourage me to attempt, at some future date, a repetition of the operation. I may mention that I have under my care several patients who are troubled with symptoms of displaced cartilage or growth in the joint, which are constantly slipping or getting pinched between the ends of the bone, thus causing pain and crippling them. I intend, therefore, if I find no improvement after a fair trial, say six months, to open their joints, and remove a fringe or fix a cartilage; and, by so doing, I shall hope to rid them of pain, discomfort, and inability to use the limb.

P.S.—The patient wrote to me on April 16th, 1888 (seven months after the operation), saying he continued quite well.

AN UNUSUAL CASE OF HÆMATURIA.

By THOMAS OLIVER, M.D., M.R.C.P.,

Physician to the Royal Infirmary, and Lecturer on Physiology, University of Durham College of Medicine, Newcastle-upon-Tyne.

THE following case is so unique in my experience that I venture to place it before the profession.

On September 11th, 1887, I was asked to see, in consultation with Dr. Campbell, of Newcastle-upon-Tyne, Miss L. B., aged 11, a pale, emaciated young lady, who for a month had been suffering from hæmaturia, accompanied by high temperature. I was told that when an infant she had suffered from some illness, which had been followed by general desquamation, but whether it was scarlet fever the parents could not say. A few years after this she had measles, and soon after this whooping-cough. A year ago she had a slight attack of hæmaturia, the cause of which was never properly understood. Her recovery was slow, and although the temperature was not taken by the medical gentleman then in attendance, the mother states that the patient was decidedly feverish, particularly towards evening. Ultimately she made a good recovery.

Whilst spending last summer at Cullercoats along with her family, her brother took ill and was brought home only to develop and pass through a very severe attack of typhoid fever. When Dr. Campbell and I met in consultation the boy was convalescent; it was the thirty-second day of the fever. Five days after the illness of this brother was distinctly recognised as typhoid fever our patient had become feverish, and hæmaturia suddenly made its appearance. The temperature ranged from 101° to 103°. Hæmaturia had persisted without any intermission from the commencement of the feverish attack. At times the urine was deep black in colour; it was always plentiful, had a specific gravity about the normal, and was acid in reaction. It was passed without pain, never contained clot, never pus cells or tube casts, but numerous blood cells. A few days before I saw her, vomiting had set in, always coming on immediately after eating, and without pain. Bowels were natural; there was slight cough, no expectoration.

When I saw her she was very ill, much reduced from the fever and vomiting; her tongue was moist and creamy; she was sleepy and seemed heavy and apathetic; pulse from 120 to 124, and it had remained at this all through her illness. Small moist musical *râles* were heard all over the lower part of the chest; the base of the right lung was rather dull on percussion. The heart's sounds were healthy; liver dullness normal; splenic dullness slightly increased. There was no tumour or increased dullness detected over the region of either kidney, but a degree of pain was experienced when the right kidney was tilted forward. The abdomen was flat, no spots were seen in the skin, no fluid was detected in the peritoneal cavity, no œdema of face, feet, or hands existed. A record of the temperature was unfortunately not kept at first, but on the evening before I saw her it was 104°. From September 2nd to the 18th, the evening temperature generally registered 1° to 3° higher than the morning; the evening temperature being 103° or 104°, whilst the morning was 101° or 102°.

On September 19th the morning temperature suddenly fell from 103° to 99.2°, and whilst on this evening the temperature rose to 100.2°, it never again did so. The morning and evening temperatures were for the future pretty nearly equal, never being higher than 99°.

All sorts of medicines had been tried, iron, gallic acid, quinine, ergot, ergotin, in the view that the hæmaturia might be in some

way or other associated with a specific fever. Sulphur, carbonate of soda, and quinine were given, but all without avail.

After the sudden fall of the temperature on September 19th, blood never appeared in the urine, and the patient had a convalescence which was quite uninterrupted.

Now here was a case of hæmaturia which began with high temperature quite suddenly; both continued for thirty-five days, the high temperature and the hæmaturia each being quite uninfluenced by drugs; then a sudden fall of the temperature occurred, also a cessation of the hæmorrhage, both followed by cure. What was the hæmaturia symptomatic of? There was nothing to suggest stone in the kidney or in the bladder, no enlargement of the kidney pointing to cancer, no history of hæmophilia, no purpura hæmorrhagica, no Bright's disease. The diagnosis, in my opinion, lay between tubercular disease and some peculiar blood condition; and we eliminated the former partly by the good family history we obtained, and the absence of any diseased condition of the lung which we could definitely regard as tubercular; the few *râles* which we heard in the chest had not been of two days' duration. From the sudden onset of the illness and the high temperature, the continuation of the high temperature and hæmaturia as associated conditions of thirty-five days, the distinct evening exacerbations, their refusal to be influenced by medicines, the enlargement of the spleen, the contemporaneous illness of the brother, and that illness unmistakably typhoid fever, and the fact that brother and sister had been living under identical conditions at Cullercoats, the one taking ill five days before the other; all these led us to regard the hæmaturia as specific, or, in other words, that our patient was also suffering from typhoid fever. Am I right in even suggesting this? All our textbooks speak of the relationship. In scarlet fever we have an illustration of how a poison known to act principally upon the skin and the tonsils may yet attack the glands in the wall of the intestine and the kidney; for as early as the second or third day of the fever the kidney *post mortem* has been found to be the seat of an interstitial or a glomerulo-nephritis. Besides, in typhoid fever itself, is it not the case that while the glands in the lower part of the wall of the ileum are the seats usually selected by the poison, there yet occurs every now and then an inflammation of the lungs with all the physical signs of a pneumonia, a pneumonia difficult to explain as regards its causation and relationship to the typhoid fever, and yet at other times so acute and so severe as to mask the other signs of typhoid fever, of which it is after all but the local expression? I am inclined to think that in our case the kidneys had to bear the brunt of the poison, or as the result of the poisoned condition of the blood, the walls of the renal vessels allowed the blood to escape. If so, then I trust I have made good my claim to this case of hæmaturia being regarded as quite an unusual one.

WOLFFBERG'S COLOUR-TEST.

By SYDNEY H. A. STEPHENSON, M.B. EDIN.,

Clinical Assistant to the Royal Westminster Ophthalmic Hospital.

By the employment of this test Dr. Wolffberg (*Klin. Monatsbl.*, p. 359, 1886) claims to have discovered a simple method of telling in a given case whether defective vision is caused by refractive errors or by defect of the light-sense. He employs two discs placed on a background of black velvet; the one is red, 2 millimètres in diameter (R²), and the other blue, and 7 millimètres in diameter (Bl⁷); if the vision is less than $\frac{1}{2}$, larger, red and blue discs are used. By comparing the distances at which these discs are recognised by the eye under examination, with the degree of visual defect, as ascertained by Snellen's types, Wolffberg believes that he is able to tell whether the defective vision is due to an error of refraction. An eye which reads $\frac{1}{2}$ should see the discs at 5.5 metres; if this be not the case, he concludes that it is suffering from some affection, other than a refractive error, which is likely to affect the light-sense.

If vision be impaired by simple ametropia, the discs should be seen with each degree of vision at a certain constant distance from the eye. Wolffberg believes that he has ascertained these distances by experiment, and he has constructed a table from which can be seen at a glance the distance at which the discs ought to be recognised with each degree of vision. If the defective vision be caused by defect of the light-sense—which is supposed to influence colour-vision more than form-sense—the distances at which the discs will be seen will correspond to constant distances,

shown in a second table, and ascertained by discovering at what distance emmetropic eyes were able to recognise the discs when the illumination was reduced by gradually diminishing the amount of solar light.

Dr. Boehm (*Klin. Monatsbl.*, November, 1887), from a large number of experiments made in Wolffberg's *clinique* and at his suggestion, believes that he has not only confirmed the above conclusions, but also that, by means of this test, he is able to detect the presence of astigmatism; since this impairs the form-sense in a much higher degree than does spherical ametropia, but has not a proportionate effect on the colour-sense.

An able and lucid review of Wolffberg's and Boehm's papers was published in the *London Medical Record* by Mr. W. Adams Frost (*London Medical Record*, Art. 7583, December 15th, 1887), at whose suggestion I made the experiments about to be described.

It is obvious that if Wolffberg's data were correct, we should have a valuable addition to our methods of diagnosis, but as my investigations on the subject do not confirm those data, I have thought it advisable to place them on record.

The observations were carried out on a fairly large number of intelligent children and adults, and the apparatus used was obtained from the source recommended by Dr. Wolffberg. The patients were first made to understand what was expected of them by being shown discs similar in every respect to those actually employed in the test, except that they were of different colours. The real test-discs were then placed at such a distance from the patient that they were invisible to him, and they were gradually approximated until they were recognised. Wolffberg adopts the opposite course, gradually increasing the distance until they become invisible. Although this difference in our respective methods of employing the test might explain trivial discrepancies, it would certainly not account for the totally different results obtained.

My experiments were limited to finding out the distances at which the discs were seen; (1) in eyes whose vision = $\frac{5}{5}$; (2) in spherical ametropia of known amount; (3) in spherical ametropia of artificial production. My original intention was to endeavour to verify all Wolffberg's conclusions, but when I found that the results obtained from the above three classes of cases did not coincide with his table, I thought it useless to proceed further, since the trustworthiness of Wolffberg's whole scheme depends on the integrity of his facts with regard to emmetropia and simple myopia and hypermetropia.

The conclusions which I have reached are as follows:

1. *As to the Distances at which the Discs could be seen by Eyes with V. = $\frac{5}{5}$.*—Many eyes were unable to recognise the coloured discs at 5.5 mètres. The majority of those examined could not distinguish the Bl⁷ disc beyond 5.0 mètres, and in some cases not beyond 3.0 mètres. According to Wolffberg the two coloured discs should be visible at the same distance from the eye, but in most of the cases examined the R² disc had to be held 0.5 to 1.5 mètres nearer the eye than had the Bl⁷ disc before it could be recognised. Many patients had great difficulty in saying the exact distance at which the discs could be seen. So far as I could see there appeared to be no definite constant distance at which different eyes whose V. was $\frac{5}{5}$ could recognise the discs.

2. *Myopia and Hypermetropia in which Corrected Vision = $\frac{5}{5}$.*—The distance at which, with each degree of vision-defect, the discs could be recognised did not correspond with Wolffberg's table. In different experiments persons who had the same degree of defective vision did not see the discs at the same distances; for example, A. (myope) had V. = $\frac{5}{5}$, and saw R² at 0.5 mètre, and Bl⁷ at 1.0 mètre; whilst B. (myope), who also had V. = $\frac{5}{5}$, saw R² at 1.25 mètres, and Bl⁷ at 1.75 mètres.

3. *Myopia and Hypermetropia produced by Convex and Concave Glasses.*—In this class of cases, also, the distances at which the discs were seen in the different degrees of visual defect did not agree with Wolffberg's table; often the results obtained from the two eyes of the same person with the same amount of defect in each gave different results; no two cases ever coincided exactly one with the other; and frequently a person when re-examined gave widely different answers from those obtained at the first examination.

In practically all cases it was found (a) that R² had to be held nearer to the eye than had Bl⁷ before it could be recognised; (b) that the worse the vision the nearer had both the coloured discs to be held to the eye, in order that they might be clearly seen; for example, V. $\frac{3}{5}$ sees R² at 3.5 and Bl⁷ at 5.0 mètres; V. $\frac{4}{5}$ sees R² at 2.5 and Bl⁷ at 4.0 mètres; V. $\frac{1}{5}$ sees R² at 1.0 and Bl⁷ at 2.0

mètres; V. $\frac{2}{5}$ sees R² at 0.75 and Bl⁷ at 2.5 mètres; V. $\frac{4}{5}$ sees R² at 0.5 and Bl⁷ at 1.75 mètres; V. $\frac{3}{5}$ sees R² at 0.33 and Bl⁷ at 1.0 mètre.

Although this test possesses some scientific interest, I fear that it is of no practical value as a help to differential diagnosis: and although there may be—and in all probability is—some ratio between form and colour-sense, I am driven to the conclusion that Wolffberg has not correctly ascertained it, and that his test is open to many sources of error from its purely subjective nature.

THERAPEUTIC MEMORANDA.

SALINE PURGATIVES IN THE TREATMENT OF TYPHLITIS AND PERITONITIS.

At a recent meeting of the Midland Medical Society I showed a patient who had recovered from an undoubted attack of acute peritonitis secondary to typhlitis. In this case opium and belladonna failed to give relief, while the administration of sulphate of magnesium and sulphate of sodium in half-drachm doses with ten minims of tincture of belladonna every four hours was quickly followed by improvement, the motions, at first liquid, becoming more and more solid till normal stools were passed. Two or three slight relapses in this case were at once checked by the mixture, and the man rapidly recovered, there remaining a small induration in the right iliac fossa.

Since the above case was recorded, I have had under my care at the Workhouse Infirmary a severe case of typhlitis. I gave the same mixture as in the first case, with great relief; in fact, enemata of soap and water and of glycerine failed to evacuate. After continuing the medicine for a week the bowels failed to act, and in a few days the abdomen was distended, there being dullness in each flank, with a distinct thrill on percussion, all the signs, in fact, of fluid in the peritoneal cavity being present. The patient was very prostrate, having been allowed only a pint of peptonised milk and a pint of beef-tea a day. I gave him three ounces of whisky, and the next morning he passed an enormous liquid motion containing scybala. I continued the stimulant, and allowed him another pint of milk. He continued to pass large motions with scybala, the enlargement of the abdomen and other signs of fluid in the peritoneal cavity completely disappearing. Evidently the saline aperient had caused a large flow of fluid into the intestine, but the bowel was not sufficiently powerful to evacuate it; restoration of tone by stimulants at once enabled the bowel to empty itself. At this time another complication appeared in the form of a painful swelling of the left parotid gland, which, however, subsided without suppuration. Finally, the patient completely recovered, and was discharged six weeks from the time of his admission.

It seems to me that in typhlitis due to faecal retention, and in peritonitis from the same cause, saline purgatives are of great value, especially if enemata fail to act. In moderate doses they do not cause peristalsis, their action is quite painless, and they are exceedingly useful in washing away hardened scybala. During their administration the abdomen should be frequently examined, and any accumulation of fluid in the intestines treated by stimulants.

C. W. SUCKLING, M.D., M.R.C.P.

Birmingham.

JAMBUL IN DIABETES.

A WOMAN suffering from diabetes was admitted into the Newbury District Hospital on September 24th, and during the whole of her stay there she was dieted. For the first nine days she was given a tonic mixture of quinine and iron. The average daily amount of urine was 98 ounces, with an average specific gravity of 1041.5.

During the next ten days she was given half a grain of opium, at first three times, and afterwards four times, a day. The average daily amount of urine was 91.3 ounces, with an average specific gravity of 1040.

During the next eleven days she took jambul, at first two grains and a half, and afterwards five grains three times a day. The average daily amount of urine was 138.3 ounces, with an average specific gravity of 1041.5.

I tested the urine frequently, and always found that sugar was present. The jambul was given in parts containing two grains and a half each, obtained from Thomas Christy and Co. It is as