so pathologically complete, to coin a phrase, that I have not felt disposed to try any other remedy. What I wish to insist upon is this, that in frequently repeated small doses of antimony we have our best known remedy at present; it not only eases the distress of the acute stage, but one which is superior to any other in inducing the greatest amount of resolution as tested by the condition of the lung afterwards.

The dyspepsia and general want of tone of the system, as well as the subsequent consolidation itself in imperfect recoveries, are best treated by mineral acids in addition to the most careful attention to every hygienic detail. Some I know prefer hydrochloric acid in an aromatic bitter, but nitric acid has appeared to me to be of greater benefit. The acids have always seemed to me to have as great an effect in improving crippled lungs as they have in benefiting atonic dyspepsia. Indeed, I suspect the indigestion they prove most useful in has its origin in some pulmonary cause.

ON THE VARIETIES OF HEPATIC CIRRHOSIS.

By GEORGE MUNRO SMITH, L.R.C.P.Lond., M.R.C.S.,

Lecturer on Physiology, Bristol Medical School; Demonstrator of Morbid Anatomy, Bristol Royal Infirmary.

VARIOUS attempts have been made from time to time to classify the different kinds of general interstitial hepatitis, either in accordance with the microscopic appearances of the organ after death, or according to the supposed causation of the disease. Ziegler doubts the usefulness of such an undertaking, inasmuch as the types pass one into another. He says: "It is enough for the present to distinguish the hypertrophic conditions from the atrophic, and even then it must be borne in mind that these conditions merely represent different degrees or different stages of what is essentially one and the same process." Dr. Saundby, on the other hand,¹ enumerates seven varieties—namely, alcoholic, cardiac or cyanotic, biliary, syphilitic, tubercular, malarial, and scarlatinal. Charcot and Hanot have, moreover, described a "hypertrophic cirrhosis of diabetes," and Cornil and Ranvier a "fatty hypertrophic cirrhosis." My object in this paper is to suggest a grouping of these morbid processes that may, perhaps, serve for a working classification.

After studying the chief literature on the subject, and comparing various sections of diseased liver, it appears to me that confusion has arisen, in the first place, from confounding the cause with the morbid anatomy; secondly, from describing partial cirrhosis² as general; and, thirdly, from considering *post-mortem* appearances in early and late epochs of the same disease as two distinct varieties.

For example, Charcot described a condition characterised by considerable enlargement of the liver, jaundice, not usually much ascites, and a rapid course. After death was found a general and uniform increase of connective tissue in the organ, with proliferation of the smallest bile-ducts and capillaries. To this disease he gave the name "biliary cirrhosis," indicating thereby its mode of origin from blocking of the bile-ducts. This term "biliary" has, however, been transferred from its etiological sense, and made to mean increased formation of the small ducts, a condition not in the least characteristic of this form of cirrhosis, but existing in quite as marked a degree in alcoholic and in cyanotic, as the examination of a few sections will satisfy anyone.

As an example of the second source of error, namely, the confounding together of partial and general fibrosis, two so-called "varieties" may be given, "tubercular" and "hypertrophic diabetic" cirrhosis. Tubercles, of course, frequently form in the liver, generally in or near the capsule; but occasionally there is found an infiltration in the interior of the gland in the interlobular spaces. If the patient lives long enough, this is sure to lead to patches of sclerosed tissue and new formation, probably, of bile-ducts. Yet this certainly should not be looked upon as a variety of cirrhosis; it is a localised inflammation round the irritating masses of tubercle.

As a rule, no lesion of the liver is found in diabetes unless the subject also happens to be an alcoholic; and although Charcot and Hanot have described two cases of enlarged liver with sclerosis in non-alcoholic patients, yet in both these cases distinct

tuberculosis existed in the lungs; hence the appearances found in the liver may, perhaps, have been also due to this cause, and the evidence in favour of a distinct variety is not, I think, clear. "Tubercular cirrhosis" may, I think, be struck off from the list, inasmuch as it is partial and not universal, and, for practical purposes, "hypertrophic cirrhosis of diabetes" may also be omitted.

Again, the period in the course of the disease affecting the liver at which the patient dies must always be taken into consideration. So-called "monolobular cirrhosis" becomes indistinguishable microscopically from "multilobular," and an enlarged liver may shrink until it is exceedingly small. Moreover, morbid anatomy cannot be divorced from clinical history, and a description of tissue change alone is sure to lead to confusion. The classification I suggest is as follows:—(a) *Obstructive cirrhosis* due to either (1) some impediment to the flow of bile, "biliary," or (2) to the flow of blood from the liver (as in pulmonary or heart disease), "cyanotic." (b) *Irritative cirrhosis* caused by the irritation (and chronic interstitial inflammation which follows) of some poison brought to the liver by the portal vein or hepatic artery. Under this heading come (1) "alcoholic," (2) "malarial," and (3) "syphilitic," all undoubtedly well-marked conditions. "Scarlatinal cirrhosis," if such exist, would come in this group, but the evidence of its existence is not very abundant.

The *obstructive* are always "monolobular," that is, the connective tissue grows uniformly, not necessarily round the lobules, but round the hepatic zone as well. In the *irritative* forms the connective tissue grows irregularly; some lobules are divided up by septa, whilst others escape altogether. The "syphilitic" is found only in children; acquired syphilis appears never to produce general cirrhosis. The "malarial" is also chiefly seen in children. In "alcoholic" the disease probably begins (Sabourin) both round the portal and hepatic systems of vessels; it is more distinctly multilobular than any other variety.

In the irritative lesions there is often thickening of the small arteries (peri-arteritis and endo-arteritis). I do not think this is common in the obstructive varieties.

As to the distinction into "hypertrophic" and "atrophic" types, it must be remembered that the liver very readily enlarges; a rich and copious dinner, especially when combined with wine or spirits, may increase its size to one-third greater than the normal.⁴ We should, therefore, expect enlargement in the early stages of interstitial hepatitis; but, in the irritative forms, this subsides much more rapidly than in the obstructive, and the latter terminate more swiftly than the former, otherwise the gland would probably dwindle and atrophy. Both forms may be complicated by adenomatous growths, and, when these are multiple, an otherwise atrophied liver appears to be abnormally large.

I offer the above classification in the hope that it may simplify a subject that has, I believe, hitherto been somewhat in confusion.

SURGICAL MEMORANDA.

OSTEOPLASTIC OPERATIONS ON THE FOOT.

It was with the greatest interest that I perused in the JOURNAL of May 5th the report by Sir William MacCormac of an osteoplastic operation on the foot, which generally bears the name of Mikulicz, of Prague, although in the above-mentioned report the credit of first devising and performing the operation is given to Wladimiroff, of Kazan. In Russia it has also been performed by Skleffasowsky, and in this country Sir William MacCormac looks upon himself as the first to do it.

My own interest in it is due partly to the fact that, in 1883, in the presence of Dr. Merson and Mr. Macnab of this town, I went through all the steps of the operation in a case of disease of the tarsus, but unfortunately I found, after removing the posterior half of the tarsus, that the other bones were involved, and I was obliged to amputate in the position of an ordinary Syme, taking a flap from the dorsum of the foot. In addition to this, however, I have for the past few weeks been trying to fix the date of operation in a case of the late Professor Syme's, which, as far as I can see, must have been performed even before Wladimiroff's first case, and which has every appearance of being an identical operation, the foot and leg exactly resembling that figured by Sir William MacCormac in the JOURNAL of May 5th.

The patient, a young woman, frequently appeared in the outpatient room at Huddersfield Infirmary when I was house-sur-

¹ *A Text Book of Pathological Anatomy*. By Ernest Ziegler. Translated by D. MacAlister, M.D.

² BRITISH MEDICAL JOURNAL, June 24th, 1886.

³ The term cirrhosis (*κίρρος*, yellow) suggests a condition not always present, but long use has sanctioned it. Sclerosis is, of course, a better word.

⁴ Cornil and Ranvier, vol. ii, p. 325.