and a piece of bread is a sufficient meal in itself, and should be available for nothing to every man who applies for it. The soup can be made from the large bones of the rations; the sale of the refuse will generally be found to cover the cost of the vegetables, split peas, and flour required to complete the soup. This has been adopted in some regiments, and it is found that the applicants for the soup are usually those whose finances are low, and on the days immediately before the pay-day, there is often a difficulty in supplying the numbers. There is naturally a greater demand in the winter. The universal adoption of this plan cannot be too strongly recommended.

A few words as to the times spirit rations should be issued; they are best taken with meals—at dinner time preferably, never before. This is not the place to discuss the merits or the action of alcohol; in small quantities it increases the secretion of the gastric juice and the movements of the stomach, and thus aids digestion. Alcohol should be taken, largely diluted with water, with, or rather at the close of, a meal. In Queen's Regulations, Sect. xv, Para. 103, it is laid down that malt liquor may be taken from the canteen to the barrack rooms for the men's consumption at dinner; this privilege is, however, not acknowledged by all commanding officers and its observance should be universally insisted on.

AFTER-HISTORY OF PNEUMONIA THE GUIDE TO ITS TREATMENT.

By ARTHUR JAMISON, M.D.,
London.

I WISH in this paper to give my experience of the imperfect as well as the complete recoveries of my cases of pneumonia during a period of over twenty years. I have never seen its therapeutics discussed from the consideration of the after-history of cases of this disease. I will not, therefore, in this paper discuss the lessons taught me by my fatal cases, but confine my remarks to those that recovered. These observations are founded on 213 cases of all grades of severity of single and double pneumonia, at ages varying from 15 to 73; and by pneumonia I mean croupous pneumonia, characterised by rusty sputa and consolidation afterwards. I was able to trace the after-history of 155 of my cases. Most of them were of the artisan class; some came under my care subsequently for other ailments, some for entrance examination into benefit societies, and I always made it a routine rule to examine the chests of all my cases whenever I could afterwards. Of these 155 cases, I found rather above half—namely, 81—free from any traces of their pneumonia on examination at periods not under two years afterwards. The other 74 had traces left discoverable by finding dulness of the affected region of greater or less extent and degree, indicating impaired lung, and were very frequently ill; 13 of them died of phthisis afterwards.

During my first years of practice my treatment was mainly expectant. I thought and had been taught that pneumonia had a definite course; that drugs could not alter its natural history, and, beyond stimulating the old, rest in bed, a warm poultice, and any sort of a placebo mixture was treatment enough. Symptoms, on an average, subsided on the eleventh day; convalescence seemed established, and expectancy triumphant. Any prolonged debility afterwards was put down to insufficient means, imperfect hygienic surroundings, or to the generally below-par state of health of many operatives living in towns. But I began to wonder if expectancy were the best treatment when I found so many whom I had so treated so frequently ill afterwards. The chief symptoms were slight cough and expectoration, inability to work as well as formerly, a constant sense of slight weariness, and in nearly every case a feeling of flatulent distension and weight after meals, and a varying amount of anorexia. Though the body might be fairly well nourished, yet the face was too thin proportionately to the body, and wore an expression of considerable depression; the bowels usually torpid; the tongue furred, and especially indented with teethmarks. On closer inquiry I found that for three or four months following the attack of pneumonia these patients got on fairly well, but after then began to droop and never felt up to the mark; and with this chain of symptoms I always found evidences of unresolved pneumonia. I then began to try other measures to prevent these after-consequences if I could. With this view some were treated by quinine in large doses—now we would say in antipyretic doses—by aconite, by ammonia, and by antimony.

Thus, of the 74 cases of unresolved consolidation, 29 were treated by pure expectancy, 20 by ammonia, 10 by quinine in full doses, 7 by aconite, and 8 by antimony in largish doses of its tartrate.

I wish to say a few words on each of these methods of treatment. My experience of aconite in doses to produce any effect is against its use at all in pneumonia, as I feel sure it increases the danger of, and often actually causes, cardiac collapse. The ammonia treatment seemed much the same as the expectant, except that I thought there was more pain in the side during its use. Quinine appeared not so risky as aconite, but just as inefficient to control the disease, even in doses large enough to lower the temperature for a time. My limited experience of it only corroborates the American estimate of its value. On the other side of the Atlantic it has been most extensively used, and last year there was a discussion in the New York Academy of Medicine on the subject. There was a general consensus of opinion upon the inutility of quinine as an antipyretic in the treatment of pneumonia. It was considered useless in small doses, and dangerous in large ones, producing cardiac and nervous depression. My impression very strongly is that any treatment directed to lower the temperature only is bad. But if a remedy is to be had that is able to lower the temperature for a time in a marked degree, surely some will say it must do good. Well, kairin is able to do this; and in the JOURNAL for January 28th there is a record of the antipyretic treatment *in excelsis* so instructive that I wish to quote it before leaving this portion of my subject:

"At first kairin was used in 10-grain doses, and it was always given when the temperature was 104° and over. After each dose the temperature fell rapidly; observations made every quarter of an hour showed that the fall was continuous for about two hours, the total fall being from 3° to 5°. There was a striking constancy in the effects produced by a given dose of kairin, whether the dose was 5, 10, or 15 grains. A dose of 10 grains always gave the best results; 15 grains did no more than 5 grains, and neither did so much good as 10 grains. The stationary period varied, but at most it lasted only a few hours; the longest time recorded was ten hours. The subsequent rise of temperature was as rapid usually as the decline. Each dose of kairin was followed within half an hour by profuse sweating and symptoms of depression, and when two doses were given in close succession (the interval being three hours and a half) marked cyanosis and collapse occurred. Kairin was exhibited twice daily for ten days; on the fourteenth day of the disease the maximum temperature was still 105°."

I wonder how many of us on reading this report of a case of pneumonia rubbed our eyes and thought what blows the pyrexia got even till the patient was black in the face. I wonder also how many were surprised to read further on that the patient died of phthisis ten months afterwards. My opinion is that he treats pneumonia best who, after having the lowest percentage of deaths, has the fewest cases of consolidation afterwards. First, because it diminishes the risk of tuberculosis. Observers like von Brunn, Siegel, and many others make it apparent that the tubercle bacillus finds its best nidus in old inflammatory products. Secondly, the condition of impaired health that too often follows even apparently good recoveries from pneumonia and the dyspepsia which, I believe, is constantly associated with and evidence of incompletely resolved pneumonias, press home this view of the aim of treatment.

My first cases treated with antimony gave better results proportionately than by the other plans of treatment, so that I was led to continue its use further. After a while many reasons led me to prefer small doses frequently repeated to the larger, more knock-down doses recommended by the older writers. These small doses are well borne even by the old. I found ʒi grain given every hour for young adults, less frequently for older people, a dose quite large enough. When pain, cough, and chest oppression are relieved, the remedy may be given less often. I continued the use of the drug for several days, often a week, after the temperature had fallen to normal, to ensure removal of the inflammatory products remaining behind. There were no ill effects from the drug used in this dosage, no sickness, no diarrhoea; its action seemed to be simply that of a respiratory sedative. Generally a little tincture of camphor was given with it to ease pain.

Of the 81 cases in which no traces of mischief were to be detected afterwards, no fewer than 65 were treated in this manner. At first I had a dread of cardiac failure, but this, as time went on, grew less and less, as I found these small doses, by easing the strain of respiration, seemed to diminish the danger of sudden heart failure. Moreover, the recoveries seemed, as it were, to be

so pathologically complete, to coin a phrase, that I have not felt disposed to try any other remedy. What I wish to insist upon is this, that in frequently repeated small doses of antimony we have our best known remedy at present; it not only eases the distress of the acute stage, but one which is superior to any other in inducing the greatest amount of resolution as tested by the condition of the lung afterwards.

The dyspepsia and general want of tone of the system, as well as the subsequent consolidation itself in imperfect recoveries, are best treated by mineral acids in addition to the most careful attention to every hygienic detail. Some I know prefer hydrochloric acid in an aromatic bitter, but nitric acid has appeared to me to be of greater benefit. The acids have always seemed to me to have as great an effect in improving crippled lungs as they have in benefiting atonic dyspepsia. Indeed, I suspect the indigestion they prove most useful in has its origin in some pulmonary cause.

ON THE VARIETIES OF HEPATIC CIRRHOSIS.

By GEORGE MUNRO SMITH, L.R.C.P.Lond., M.R.C.S.,

Lecturer on Physiology, Bristol Medical School; Demonstrator of Morbid Anatomy, Bristol Royal Infirmary.

VARIOUS attempts have been made from time to time to classify the different kinds of general interstitial hepatitis, either in accordance with the microscopic appearances of the organ after death, or according to the supposed causation of the disease. Ziegler doubts the usefulness of such an undertaking, inasmuch as the types pass one into another. He says: "It is enough for the present to distinguish the hypertrophic conditions from the atrophic, and even then it must be borne in mind that these conditions merely represent different degrees or different stages of what is essentially one and the same process." Dr. Saundby, on the other hand,¹ enumerates seven varieties—namely, alcoholic, cardiac or cyanotic, biliary, syphilitic, tubercular, malarial, and scarlatinal. Charcot and Hanot have, moreover, described a "hypertrophic cirrhosis of diabetes," and Cornil and Ranvier a "fatty hypertrophic cirrhosis." My object in this paper is to suggest a grouping of these morbid processes that may, perhaps, serve for a working classification.

After studying the chief literature on the subject, and comparing various sections of diseased liver, it appears to me that confusion has arisen, in the first place, from confounding the cause with the morbid anatomy; secondly, from describing partial cirrhosis² as general; and, thirdly, from considering *post-mortem* appearances in early and late epochs of the same disease as two distinct varieties.

For example, Charcot described a condition characterised by considerable enlargement of the liver, jaundice, not usually much ascites, and a rapid course. After death was found a general and uniform increase of connective tissue in the organ, with proliferation of the smallest bile-ducts and capillaries. To this disease he gave the name "biliary cirrhosis," indicating thereby its mode of origin from blocking of the bile-ducts. This term "biliary" has, however, been transferred from its etiological sense, and made to mean increased formation of the small ducts, a condition not in the least characteristic of this form of cirrhosis, but existing in quite as marked a degree in alcoholic and in cyanotic, as the examination of a few sections will satisfy anyone.

As an example of the second source of error, namely, the confounding together of partial and general fibrosis, two so-called "varieties" may be given, "tubercular" and "hypertrophic diabetic" cirrhosis. Tubercles, of course, frequently form in the liver, generally in or near the capsule; but occasionally there is found an infiltration in the interior of the gland in the interlobular spaces. If the patient lives long enough, this is sure to lead to patches of sclerosed tissue and new formation, probably, of bile-ducts. Yet this certainly should not be looked upon as a variety of cirrhosis; it is a localised inflammation round the irritating masses of tubercle.

As a rule, no lesion of the liver is found in diabetes unless the subject also happens to be an alcoholic; and although Charcot and Hanot have described two cases of enlarged liver with sclerosis in non-alcoholic patients, yet in both these cases distinct

tuberculosis existed in the lungs; hence the appearances found in the liver may, perhaps, have been also due to this cause, and the evidence in favour of a distinct variety is not, I think, clear. "Tubercular cirrhosis" may, I think, be struck off from the list, inasmuch as it is partial and not universal, and, for practical purposes, "hypertrophic cirrhosis of diabetes" may also be omitted.

Again, the period in the course of the disease affecting the liver at which the patient dies must always be taken into consideration. So-called "monolobular cirrhosis" becomes indistinguishable microscopically from "multilobular," and an enlarged liver may shrink until it is exceedingly small. Moreover, morbid anatomy cannot be divorced from clinical history, and a description of tissue change alone is sure to lead to confusion. The classification I suggest is as follows:—(a) *Obstructive cirrhosis* due to either (1) some impediment to the flow of bile, "biliary," or (2) to the flow of blood from the liver (as in pulmonary or heart disease), "cyanotic." (b) *Irritative cirrhosis* caused by the irritation (and chronic interstitial inflammation which follows) of some poison brought to the liver by the portal vein or hepatic artery. Under this heading come (1) "alcoholic," (2) "malarial," and (3) "syphilitic," all undoubtedly well-marked conditions. "Scarlatinal cirrhosis," if such exist, would come in this group, but the evidence of its existence is not very abundant.

The *obstructive* are always "monolobular," that is, the connective tissue grows uniformly, not necessarily round the lobules, but round the hepatic zone as well. In the *irritative* forms the connective tissue grows irregularly; some lobules are divided up by septa, whilst others escape altogether. The "syphilitic" is found only in children; acquired syphilis appears never to produce general cirrhosis. The "malarial" is also chiefly seen in children. In "alcoholic" the disease probably begins (Sabourin) both round the portal and hepatic systems of vessels; it is more distinctly multilobular than any other variety.

In the irritative lesions there is often thickening of the small arteries (peri-arteritis and endo-arteritis). I do not think this is common in the obstructive varieties.

As to the distinction into "hypertrophic" and "atrophic" types, it must be remembered that the liver very readily enlarges; a rich and copious dinner, especially when combined with wine or spirits, may increase its size to one-third greater than the normal.⁴ We should, therefore, expect enlargement in the early stages of interstitial hepatitis; but, in the irritative forms, this subsides much more rapidly than in the obstructive, and the latter terminate more swiftly than the former, otherwise the gland would probably dwindle and atrophy. Both forms may be complicated by adenomatous growths, and, when these are multiple, an otherwise atrophied liver appears to be abnormally large.

I offer the above classification in the hope that it may simplify a subject that has, I believe, hitherto been somewhat in confusion.

SURGICAL MEMORANDA.

OSTEOPLASTIC OPERATIONS ON THE FOOT.

It was with the greatest interest that I perused in the JOURNAL of May 5th the report by Sir William MacCormac of an osteoplastic operation on the foot, which generally bears the name of Mikulicz, of Prague, although in the above-mentioned report the credit of first devising and performing the operation is given to Wladimiroff, of Kazan. In Russia it has also been performed by Skleffasowsky, and in this country Sir William MacCormac looks upon himself as the first to do it.

My own interest in it is due partly to the fact that, in 1883, in the presence of Dr. Merson and Mr. Macnab of this town, I went through all the steps of the operation in a case of disease of the tarsus, but unfortunately I found, after removing the posterior half of the tarsus, that the other bones were involved, and I was obliged to amputate in the position of an ordinary Syme, taking a flap from the dorsum of the foot. In addition to this, however, I have for the past few weeks been trying to fix the date of operation in a case of the late Professor Syme's, which, as far as I can see, must have been performed even before Wladimiroff's first case, and which has every appearance of being an identical operation, the foot and leg exactly resembling that figured by Sir William MacCormac in the JOURNAL of May 5th.

The patient, a young woman, frequently appeared in the outpatient room at Huddersfield Infirmary when I was house-sur-

¹ *A Text Book of Pathological Anatomy*. By Ernest Ziegler. Translated by D. MacAlister, M.D.

² BRITISH MEDICAL JOURNAL, June 24th, 1886.

³ The term cirrhosis (*κίρρος*, yellow) suggests a condition not always present, but long use has sanctioned it. Sclerosis is, of course, a better word.

⁴ Cornil and Ranvier, vol. ii, p. 325.