After this Mr. C. gained steadily in strength, and before he left the country, about the end of December, was able to walk two or three miles at a stretch, his pulse being generally about 70.

He spent the winter in Edinburgh, and about a month of spring in North Wales. He is now at his residence in Perthshire, and pretty well out of the doctor's hands. He has lately spent the greater part of three days in the Glasgow exhibition, which I take to be a pretty severe trial of strength. He still continues to take the liq. arsenicalis for every alternate month.

Dr. AFFLECK thought there was now sufficient clinical evidence to justify an alteration in the nomenclature of this disease. Many of the cases, having all the characteristics of pernicious anæmia, recovering completely under treatment by arsenic, in conjunction with a suitable diet. Dr. Affleck had repeatedly had cases of this disease under treatment in the Royal Infirmary, Edinburgh, where hæmacytometric observation has shown corpuscles reduced considerably below 1,000,000 per cubic millimetre completely recover under treatment by arsenic and iron.—Dr. M. A. BOYD said that, in a great many cases of chronic dilatation of the stomach, supervening on chronic gastric catarrh and without organic disease, a condition of anæmia and exhaustion supervenes, which was often mistaken for pernicious anæmia. The history of this case would closely correspond. The persistence of chronic gastro-enteric derangement for so many years would produce all the symptoms enumerated, more especially as the patient improved ultimately with arsenic.

ON THE VALUE OF ELECTROLYSIS IN ANGIOMA AND GOITRE.¹

By JOHN DUNCAN, LL.D., M.A., F.R.C.S.E., etc.,

Surgeon and Lecturer on Clinical Surgery, Royal Infirmary, Edinburgh.

I THINK it may be interesting to put on record a few of the outstanding cases of angioma which have come under my observation during the last year or two, with the view of illustrating and defining the varieties in which electrolysis is the best treatment at our command. I shall select the cases because they present individual peculiarities which fit them for this purpose, and because they are in themselves unusual. Nævi, or the congenital forms of capillary angioma, of course present themselves in large numbers. I choose three.

CASE I.—A remarkable example of the cutaneous variety was brought under my notice by my colleague, Mr. Miller, not because it was possible to treat it, but because he knew I was interested in the subject; and I cite it simply as typical of a class for which as yet we have no reliable means of cure. In this child the whole of one thigh as high as the buttocks and other extensive areas of the body were of a most delicate pink tint, which, indeed, it would have been difficult to recognise as abnormal but for the sharpish edge where it met the normal skin, and the possibility of demonstrating the dilated vessels by gently stretching the skin. Besides this, there were dotted irregularly over the body a multitude of black specks about the size of a pin's head, more and less, which looked like minute extravasations or pigmentations, but which careful examination showed to consist of closely-packed vessels. I have never seen the cutaneous nævus or port-wine stains at once so delicate in colour and so extensive, nor have I seen so extraordinary a number and blackness of these tiny vascular dots.

CASE II.—A case of nævus interesting for its size in a puny child three months old was brought to the infirmary in May, 1887. The tumour, gigantic when compared with the patient, was the size and shape of a medium-sized Jaffa orange. It was situated behind the ear. The whole surface was of a bluish pink colour not uncommon. It was exceedingly soft, and by pressure the blood could be so entirely forced out of it that it became completely wrinkled and flaccid. It was also remarkably slow in refilling. An absolute cure in every respect, not a mark or trace being left, was attained by three electrolytic operations, one of them performed in my absence by Dr. Hodsdon.

CASE III.—On Dr. Taylor's recommendation a nævus was put under my care, which presented this striking peculiarity, that a portion of it occurred in bone. It occupied the anterior and

chiefly the outer part of the foot in a child about eight months old. It did not involve the skin in the slightest degree, but reached completely through and through the foot from dorsum to sole. By pressure four metatarsal bones could be distinguished, of normal size and shape, although all except the first were imbedded in the growth. But the fourth metacarpal bone was very greatly enlarged, and, on introducing the needles into it, they plainly entered first the vascular tissue, then passed through a thin osseous and cartilaginous shell, next the nævoid medullary cavity of large size, and lastly the opposite circumference of the bone. As the child has advanced in age this toe has not developed with the same rapidity as the others, and is now somewhat recessed.

There was at first considerable difficulty in diagnosis. The total absence of any affection of the skin, except as to one or two dilated veins which any tumour might cause, and the expansion of the metatarsal bone, were the sources of this difficulty. But the compressibility, extent, and uniform consistence of the portion external to the bone formed the grounds for believing the tumour to be a nævus and not a sarcoma, an opinion which has been confirmed by the lapse of two years, and the behaviour under treatment.

It is plain that such a nævus was amenable to no other method of treatment than amputation or electrolysis. To leave it alone while it was rapidly growing would have been unjustifiable, if there were any possibility of safely arresting its progress. The whole anterior part of the foot was so involved as to make excision impossible. The injection of coagulants is to be looked upon as acting in a somewhat similar manner to electrolysis, but is always less manageable, more dangerous, and not so effective. Amputation was not to be thought of in a condition not dangerous to life, and in which a cure natural or artificial was still possible. In short, the case was one of those in which the advantages of electrolysis stand out in the clearest manner, because it was really the only available method, was free from danger, and with time and perseverance was absolutely certain of success.

The applications were somewhat difficult and uncertain because the child lived two hundred miles away, but six operations have had this effect: that the tumour has become greatly consolidated, is shrivelled, and is steadily diminishing in size.

CASE IV.—From several examples of extensive cavernous angioma I select one which was sent to me by Mr. Lawson Tait. As I have elsewhere pointed out, these forms are generally adult varieties, a few commencing late but most being extensions and alterations of congenital examples. In this young lady the tumour occupied the bend of the elbow and extended over a large part of the anterior and inner aspect of the upper arm. It had been treated in various ways, and injection of perchloride of iron had caused much sloughing and great danger from septic poisoning. It showed extremely well the large caverns which exist in these cases. Some could be felt with the needles to be as large as walnuts, and one or two had a covering including skin and venous wall not thicker than a piece of parchment. The operations were performed at intervals varying from six weeks to three months. The difficulty consisted in producing sufficient effect on the walls of the angiomatous caverns without involving the skin in destruction. Now, however, after working at it for more than a year and a half I have the satisfaction, thanks to the indomitable perseverance of the patient, of knowing that it is at length cured. Here again no other method of treatment was possible. The question of amputation had indeed been seriously entertained in despair of success by other remedies.

The other cases which I wish to record are examples of pulsatile angiomata.

CASE V.—The first is the case of a medical student, sent to me by Dr. Frew, of Kilmarnock. It belongs to a common class, the ordinary pulsatile nævus. A considerable portion of the side of the neck and face was occupied by a lightly tinted port-wine stain, but of late an extraordinarily rapid development of a pulsating swelling had occurred in the ear, so that the auricle resembled a huge pear, and the pulsations could be seen at any distance. One operation sufficed to cure the disease.

CASE VI.—Cirsoid aneurysm is comparatively a rare disease. I had the satisfaction of seeing a short time ago a girl, aged 19, in whom an exceedingly severe form had been entirely cured by electrolysis in 1886. She was sent to me by Dr. John Moir, of St. Andrews. The tumour chiefly affected the left temporal artery, but extended to the occipital on the same side and the temporal on the other, forming an irregularly rounded tumour more than three inches in diameter. Two operations consolidated the greater

¹ Read in the Section of Surgery at the Annual Meeting of the British Medical Association, held in Glasgow, August, 1888.

portion of the mass, but four subsequent sittings were required before I succeeded in obliterating some tortuous branches of the anterior temporal which curved backwards into the tumour, a long and very twisted portion of the occipital, and the neck of the tumour in front of the ear. In the end the success was most gratifying, for not a trace of the disease remains except in a slight flattening of that side of the skull.

CASE VII.—About the same time a woman, aged 48, was sent to me by Dr. Robert Moir, also suffering from cirroid aneurysm. This case was peculiar in that it consisted of one central branch of the temporal, which was dilated to the thickness of the little finger, and was folded and refolded on itself in its course from the ear nearly to the vertex. I have never seen a case in which the dilatation was so perfectly confined to a single vascular tube. This condition made it more amenable to treatment. One operation sufficed for the cure.

Coincidences of occurrence such as presented themselves in these two cases are curiously frequent, but this was rendered still more striking by the fact that at the same time, in June, 1886, I had under treatment two cases of traumatic arterio-venous aneurysm of the orbit, concerning which I was consulted by my colleague, Dr. Argyll Robertson.

CASE VIII.—J. M., aged 11. Five months before I saw her she wounded her right lower eyelid with the broken rib of an umbrella, which caused free bleeding and extravasation. After a few weeks the eye began to protrude, and diplopia occurred. On admission to the infirmary thrilling pulsation could be felt over the eye, most markedly towards the inner side. At each pulse the eyeball was protruded, and pulsating veins could be seen in the upper lid. A purring murmur could be heard all over the head from forehead to occiput.

Operation, June 25th, 1886.—The needles were inserted and the current passed for about fifteen minutes. Considerable ecchymosis followed their withdrawal, and on its subsidence great improvement was manifest. A second operation of seventeen minutes' duration was performed on July 15th, and on July 28th the patient went home with the exophthalmos nearly gone and the pulsation imperceptible. She was to return if any recurrence of the disease manifested itself, but has not done so, and has unfortunately been lost sight of. Practically the disease was cured when she left, and I have no doubt continues so.

CASE IX.—A. F., aged 3. Two years before admission to the infirmary, she ran against a knitting needle, which punctured the lower eyelid near the inner angle. There was no hæmorrhage, but quickly much swelling and extravasation ensued. On its subsidence, a divergent squint became manifest, and the eyeball began to protrude. On June 10th, 1886, I saw her, with Dr. Argyll Robertson and Dr. Joseph Bell. Considerable exophthalmos was found, with strong pulsation and thrill of the orbital contents, especially towards the inner angle. A murmur was heard over a wide area of the usual rough to-and-fro character. The symptoms were decidedly on the increase. It was determined to electrolyse the growth, which was accordingly done next day, the operation lasting about half an hour. This was followed by considerable swelling and ecchymosis. As this subsided, the child's complaints of the eye trouble quickly ceased. The *bruit* and pulsation steadily diminished, and, with the exophthalmos, had almost disappeared when the patient left a fortnight afterwards. On January 17th, 1887, the patient was brought back for examination. There were a few slightly dilated subconjunctival veins, and, on careful comparison, it was thought that the left eye protruded somewhat more than the right; but there was no *bruit* or pulsation, and, ophthalmoscopically, the eye was found to be quite healthy and the sight good.

These cases may serve to illustrate the value of electrolysis as a means of treatment for vascular tumours. In looking over the records of angiomas which have come under my observation of late years, I have been struck by the fact that I operate by electrolysis in more than two-thirds. This, however, is doubtless chiefly due to the fact that the medical friends who are good enough to send me cases choose those which are either unusual and serious, or specially fitted for cure by electricity; and it is therefore important to define accurately the precise position of electrolysis in relation to other modes of treatment.

I have already pointed out, but I think it well to reiterate, that electrolysis is the best treatment we possess, and, in some cases, the only one for nævi, in which the three following conditions are coincident—namely: 1, that the growth is extending; 2, that it is important to avoid a scar; 3, that the subcutaneous bears a con-

siderable proportion to the cutaneous part of the tumour. If the growth be not extending, it is unnecessary to interfere, because more than half the total number of nævi of the mixed and subcutaneous type spontaneously disappear. If it be not of importance to avoid a scar, there are other methods of treatment, such as excision and ligature, which are equally certain and more speedy. If the nævus be a pure port-wine stain, it will not spontaneously disappear; but, on the other hand, neither electrolysis nor any other method can be relied on to cure it without also destroying the skin in which it lies. Certain exceptions, however, must be made to these general propositions. When a mixed nævus has prolonged its existence into adolescent life, it ought to be treated, because there is no further hope of its disappearance. When a mixed nævus extends over a wide area, even although it be of little importance to avoid a scar, the perfect certainty and safety of electrolysis gives it the preference over every other method. In the pure port-wine stain success may sometimes be attained by the long maintenance of an artificial dermatitis.

The first case I have related is a good illustration of the class which is beyond the reach of treatment of any kind. The second is a typical example of the nævi in which electrolysis is our best treatment, because it is safer and more certain than any of the methods which avoid a scar. The third is a case in which the only possible alternative to electrolysis was amputation. The fourth, as occurring in an adult, is an example of the nævus in which it is no longer possible to hope for spontaneous cure, and in which the only treatment short of amputation is electrolysis. The fifth case is one of those which, continuing into adult life, sometimes then become truly dangerous from rapid pulsatile spread, and the possibility of hæmorrhage from injury or ulceration. Indeed, in this case ulceration had actually begun to manifest itself. The more dangerous and less effective method by the injection of coagulants was the only possible alternative.

The two cases of cirroid aneurysm illustrate the value of electrolysis in this form of disease. I have now operated in eleven cases without failure or accident. There is, indeed, no alternative but expectancy. I find, however, that this statement is not yet generally accepted by surgeons. In a case sent to me by Dr. Beatson of Glasgow, and cured three weeks ago by one operation, the patient had received many varieties of advice, of which the most reasonable was to leave it alone, the most dangerous was to ligature all the affluents and excise. The most reasonable plan, I say, is to leave it alone if the surgeon is not acquainted with the effects of electrolysis, because the disease is slow in its progress, and, except from traumatism, little dangerous, while all other methods of treatment are either very formidable, very ineffective, or both.

I have no hesitation in saying that the only justifiable method of treatment for cirroid aneurysm is electrolysis. Its perfect safety and reliability ought to determine its use, even though the tumour be not enlarging.

I would here distinguish true cirroid aneurysm from what may be called aneurysm by anastomosis, or that form of angioma which consists of large cavernous tissue, communicating on the one side with enlarged arteries, on the other with dilated veins. A considerable number of them may likewise be cured by electrolysis, but I confess to having failed in one or two large and exceptional cases which I have elsewhere recorded.

The eighth and ninth cases are examples of traumatic arterio-venous aneurysm, and the only two which I have had an opportunity of treating by electrolysis. The situation prevented the application of any other remedy, and the success was gratifying, although not more than was to be expected.

A word only remains to be said as to the *modus operandi*. I find that I habitually work with a current of between 40 and 80 milliamperes; but that is really a matter of small importance. A galvanometer is of no value in operations where the duration is determined by the palpable and visible effect produced. It is very different in cases where you cannot see and feel the gradually increasing swelling, tension, and hardness. There a means of measuring the current in operation is essential.

Four to eight cells of large size and good electro-motor force are most convenient to work with. It is not necessary to use a large number of cells if both poles be introduced as they ought to be. The body is as good a conductor as acidulated water if the resistance of the skin be avoided.

I introduce both electrodes (insulated, of course, so that the operation may be truly subcutaneous), but I work chiefly with the negative pole. It is to be remembered that the effect is pro-

duced mainly by shrivelling up and destroying the vascular walls' and that coagulation of the blood is a matter of very secondary importance; therefore, I keep moving the negative pole about and penetrating as many vessels with it as I can, because its destructive effect is more powerful and diffuse than that of the positive. I maintain it in one place just long enough to bring about a decided effect, and then change to another.

In these operations the process is essentially distinct from that by which Apostoli and others have produced their effects upon uterine tumours. However his results may be brought about, it is not by electrolysis proper, it is not by the decomposition of fluids and solids at the poles of the battery. It may be neurotic or vascular or trophic or osmotic, it is not electrolytic, and to call it so is no doubt a misnomer. I have for several years, however, been applying electrolysis in the way I describe to various forms of goitre with very striking results, and I have been impressed by the fact that on several occasions the desired effect has been produced by one operation. This was the case, for example, in two out of three cases sent to me by Dr. Watson Campbell, of Duns. There has just left my wards a woman, aged 25, on whom I operated two years ago for a goitre of considerable size, chiefly on the right side. It shrivelled up very quickly after one operation, leaving only a tiny knot low down on the left side. She came back recently on Dr. McBride's recommendation, because this lump was giving her some uneasiness, and she was anxious to get rid of it entirely. I have thus had a rare opportunity of noting the result. Having ascertained by puncture that the swelling was partially cystic, I cut down upon and shelled out a beautiful little cysto-adenoid growth about an inch and a quarter in diameter. I found the gland itself normal in size. I took away a small portion of the left lobe, to which the tumour was attached. I could detect nothing unusual in its structure, but it must be remembered that the needles had been chiefly worked in the right lobe.

I do not think that this rapid atrophy after one operation is likely to be the rule. In the larger number I have found more are required. Moreover, this speedy cure is not without its parallel among the angiomas, in which such success certainly depends on the thoroughness of the operation in reaching all parts of the tumour. It may be, then, that they were simply further examples of absorption of destroyed parts and cicatricial contraction of the rest. Still, however, these thyroid operations were largely tentative, and fell far short of what I should have considered necessary in angioma; and since Apostoli has described his method by the passage of strong currents for short times, without appreciable electrolysis at the poles, I have sometimes thought that, perhaps, there might be something analogous in those thyroid operations.

I have not yet had sufficient experience to be able to define with precision the value of electrolysis in goitre. I can only give the results so far as they have gone.

Fourteen cases have been operated upon. Three are at present under treatment, having been recently operated upon, and already showing beneficial results. One is a fibro-cystic, another a strongly marked exophthalmic, the third a vascular goitre, with considerable dyspnoea. Eleven remain, two or three having slight vascular and exophthalmic symptoms, others with slight dysphagia or dyspnoea, all of the vascular variety. Four of them I have found it impossible to trace. These cases are in the infirmary only for a day or two, and the records of their addresses have been imperfectly kept. In two cases, however, they returned for a second operation, having been benefited by the first. Of the others, six have been absolutely cured, and the seventh, I hear, has the tumour still in no way changed by one operation. I think the results are very encouraging, and that, with greater experience, one will be able to operate with more confidence. The chief difficulties have hitherto lain in the manipulation of the needles, and our want of knowledge of the effects which might be produced on surrounding structures in so important a locality.

SUCCESSFUL VACCINATION.—Mr. T. Pickthorn Thomson, M.B., of Half-Way House, Shrewsbury, has been awarded, for the second time, the Local Government grant for efficient vaccination.

PRESENTATION.—Mr. A. R. Anderson, F.R.C.S. Eng., was presented on Monday, October 29th, at the General Hospital, Nottingham, by the president, staff, and numerous other friends, with a walnut library table and chair, a dining-room clock, and a brass newspaper stand, on his resigning the post of Senior Resident Medical Officer.

INTRODUCTORY ADDRESS ON THE REFORM OF MEDICAL EDUCATION.

Delivered at Glasgow University at the Opening of the Session 1888-89.

By GEORGE BUCHANAN, M.D.,
Professor of Clinical Surgery, University of Glasgow.

PROFESSOR BUCHANAN said he was going to speak a few minutes, not in dictating to his audience, but in suggesting certain topics for their consideration and certain topics upon which he thought they might take some action, because they were, perhaps, aware that probably before the end of this session, for the first time, the students would be recognised by the State as a corporate body in the University. He was going to take up two topics of importance. The first was free trade as against protection; and, secondly, some considerations with regard to the curriculum and examination. What had free trade and protection or monopoly to do with the University, as applied to the study, the teaching, and the practice of medicine? Should the study of medicine be regulated or influenced simply by the principles of free trade; and the same with teaching, and the same with practice? He had not the least doubt that all young, generous minds who heard this stated for the first time would say, "Down with monopoly! Free trade for all connected with medicine." When he was a student, the principles of free trade regulated the entrance into the profession. All that a student had to do was to pay his fee, get his name enrolled, and sit on the benches. No questions were asked. He continued to do so till the end of four years, and then he was examined in one day from botany to medicine. He either came out a doctor or not a doctor. In 1860 free trade was abolished, and now they were under protection; and the protection consisted in demanding that the student should pass a preliminary examination before he began his studies. Was this preliminary examination serving the purposes intended? The original idea of a preliminary examination was to secure that a young man should be engaged in some studies preparing him for the special study of medicine up to the very time he enrolled as a student. But now they passed the preliminary subjects piecemeal, and in many cases left off the really educative subjects as soon as they could pass them—perhaps two or three years before registering as students of medicine. In the preliminary examination there was a great slaughter in mechanics; 204 entries for mechanics, with 101 passes. Why was that? It was easily explained. There was no means in country schools for a young man learning mechanics, and he had to get it up by cramming out of a class book. He thought that with regard to entrance to professional study, mechanics ought to be taken out of the registration subjects. What he held was that a student ought to pass in everything else required for registration; and every student should be compelled to begin his studies in May, not October, and during the first summer he could devote himself to elementary mechanics and physics, and to botany and zoology, and pass in these before the first winter session. In the first winter the student should devote his time solely to anatomy and chemistry, which he could do if the ordeal of the examination in botany and zoology were removed out of the way; while at present during the first winter the examination in botany and zoology was weighing on his mind, which should be free for chemistry and anatomy, especially the latter. Or if the student could afford the time, one whole year might be most usefully spent in mechanics and demonstrations in physics—as well as botany, zoology, and chemistry—as the first medical academic year in Cambridge was arranged. Unquestionably the time had arrived when the curriculum must be extended to five years—and in the case of those who had had no opportunity to study mechanics and physics, probably to six years—but in order to utilise their first year for either botany, zoology, or chemistry, or all, the subject of mechanics might well be included in the first professional examination of a six years' course, instead of regis-

¹ The address was delivered without the use of MS. It appears here in an abridged form, in a report by a shorthand writer. Hence the colloquial and sometimes fragmentary style.