CLINICAL LECTURE ON RETENTION OF URINE.

Delivered in University College Hospital.

By CHRISTOPHER HEATH, F.R.C.S.,

Holme Professor of Clinical Surgery in University College, London.

I PROPOSE to-day to speak of retention of urine, and it will be convenient to consider it at different ages. In the case of a newborn child there is sometimes a difficulty in passing water, which always depends upon a contracted orifice of the prepuce. When the child tries to make water it "balloons" its foreskin, swells it out, that is, into something like a child's balloon. Of course, the treatment in such a case is obvious and simple. You have only to dilate the orifice of the prepuce with a pair of forceps, or if it is at all a severe case, when the child is a little older, you had perhaps better circumcise it, and there is an end of the matter. But you sometimes meet with retention in the case of children who are a little older, and I believe that retention in children, not dependent upon a tight foreskin, is almost always the result of calculus in the urethra. A child may have a small calculus in the kidney without any obvious symptoms; he has never made any complaint, until one day you are sent for because he does not pass his water; perhaps for some hours the bladder has been distended, and there is obvious retention of urine. Now in a case of that kind always suspect a stone and look for it; usually it is easily found just within the meatus urinarius. You take a silver probe and pass it down the urethra, and you feel it click against the stone, and if the calculus is just within the meatus it is an exceedingly simple matter. You have only to get it out with a little scoop, or if you have not one, the loop of a hairpin is a convenient thing. If the orifice of the urethra is at all tight you had better nick it in order to get the stone out; but remember when you are doing this that you should always cut downwards and not upwards into the glans penis. But sometimes the stone is not at the anterior part of the urethra, but behind; and these are more difficult cases, and are apt to be overlooked. The stone does not produce complete retention perhaps, although it makes a difficulty, and then ulceration is sometimes set up, with abscess, and possibly extravasation of urine; but where there is real retention of course you can make it out by passing a probe or catheter down and touching the stone, and you may be able to feel it through the perineum. In that case it is best, if you can, to extract the stone, but it is not easy to do it. You will have to put the patient under chloroform, and with a very fine slender urethral forceps you may be able to extract the stone. More frequently the stone has become fixed, and you cannot get it out. Then the best thing is to cut upon it at once, for it is not a serious thing in children to open the urethra. You have the stone to guide you, and it is much better that you should at once cut upon it and take the stone out than leave it there to set up, as it certainly will do, irritation, and possibly, as I have said, abscess and extravasation of urine. Besides, if it is a case of *bona fide* retention, the child cannot be left many hours without being relieved.

Then there is one other thing in the case of children that should always be borne in mind; that the child may have tied a string or thread round the penis. Such cases are met with every now and then. Two or three years ago a child was brought one morning to my consulting room with the glans penis hanging merely by a thread of cellular tissue. It was as nearly cut off as it could be, but not absolutely. I simply took a pair of scissors, cut the remaining thread, and took it off. Round the glans penis just where it had been divided there was a long hair; and whether that hair had cut it through, I cannot tell. The child utterly denied ever having had a thread tied round it, but it is just conceivable that a long hair might have been twisted round accidentally, and have cut its way through.

Whenever a child has a swollen penis which is not clearly a case of paraphimosis, you should ascertain whether there is a thread deeply embedded in the groove, which is formed by the swollen tissues all round.

[1452]

Now let me pass on to a much more important matter, the retention of urine in adult age. With young adults, I suppose the most common cause of retention is some mischief supervening upon gonorrhoea. People who have perfectly healthy urethras do not get retention of urine. It is only after they have had gonorrhoea or some mischief about the urethra that they are liable to attacks of this sort. Sometimes you find a patient with retention of urine during gonorrhoea, or at all events in the later stage of it; and if that is the case, it is more usually due to some inflammation travelling to the prostate. The prostate becomes swollen and tender; if you put your finger into the rectum, you will feel it there; and it is because of that swelling of the prostate that the man cannot pass water. The obvious thing in such a case is to pass a catheter for him very gently through the prostate, and draw the water off. You proceed to treat the prostatic inflammation by a leech or two to the perineum, and then the swelling generally subsides, but it may, of course, run on to an abscess. And remember that these cases of acute prostatitis in young men are more apt to run on into abscess than any other form of prostatic disease. Then the later stage, after the gonorrhoea, is perhaps the more critical time for the urethra as regards retention. What happens is this. The man during the gonorrhoea has been debarred from all liquor, and then, assuming that he is well, he perhaps takes more than he should do. In hospital practice and amongst the lower classes you will meet with that kind of case much more frequently than in private practice. The better class of people do not, as a rule, drink more than is good for them; they do not indulge in that sort of debauch, but a labouring man is apt to take more liquor than is good for him after abstaining for a few weeks, and then he gets retention of urine.

Now that form of retention is by some surgeons thought to be purely spasmodic, others call it inflammatory. I believe it is really a little of both. If it is not inflammatory, it is congestive certainly, and there is a certain amount of spasm mixed with it. Therefore, whether we call it congestive retention or spasmodic retention, we mean the same thing. At all events, clinically, you find a patient who has had gonorrhoea, and apparently recovered from it, who is unable to pass his urine. Sometimes such patients go on for many hours before they knock up the house-surgeon, so that at one or two o'clock in the morning you are called upon to relieve such a patient, and, of course, it is important that relief should be given at once. Now it is quite true that these cases will mostly get well of themselves, if you can have time. I mean, if you give a patient of that class a good dose of opium, put him into a warm bed, and put fomentations about his perineum, in the majority of cases the retention will pass off, and the patient will make water. But you must bear in mind the circumstances of the case. If it is a hospital patient, the house-surgeon is called up in the night, and naturally he wants to get to bed again. There are no great conveniences for putting the patient into a hot bed, or even into a hot bath, in the middle of the night, and you have to pass a catheter for him; and, as Sir Henry Thompson says, in one of his lectures, no doubt the patient much more appreciates your skill, if you pass a catheter without hurting him, and draw his urine off, than if you merely gave him a dose of medicine, put him into a warm bath, and let Nature do the rest. It is better, then, on the whole, to draw off the urine at once. The great thing is to use a good-sized catheter. These patients have no stricture, and you want a good-sized catheter; and of all catheters, commend me to the old-fashioned English gum-elastic catheter. If you take a catheter of the size I hold in my hand (No. 9), with a good blunt end to it, and if you warm it and oil it well, and pass it, with a stilette if you like or without, until you get to the obstruction point, you will find it generally somewhere about the bulb of the urethra; and then, if with a little gentle pressure you push against it, you will be conscious of a sort of yielding; and so, as you steadily push on, the catheter goes into the bladder, and you draw the water off. That, of course, is an extremely satisfactory thing for all parties; but the affair does not always come off so smoothly as that. You may find a difficulty in the way. Possibly there may be more congestion than you think, more spasm perhaps, and the obstruction does not yield quite so readily. Now the greatest mistake that you can make, I believe, is to use very small catheters. No. 7 is a very good average size, and, if you cannot get that in, the patient having no history of previous stricture, you had better put the patient under chloroform, and you will be able to pass the catheter with the greatest ease. It is much better for the patient that you should stick to that catheter or thereabouts than that you should poke holes in his urethra

with a small instrument; and, if you give the patient chloroform, you overcome the difficulty at once. I am aware that, when you get into practice in the country, you may have to modify your proceeding. When a medical man is on his rounds and is stopped somewhere because a patient has retention of urine, he may not have chloroform with him, and perhaps not a catheter. Those are the cases in which you may fall back upon opium, with warm fomentations. If you give a patient of that kind a good dose of opium, and have fomentations put about the perineum, the probability is that in a couple of hours he will be able to make water. If he does not, then of course he will send for you, and you must be prepared to pass a catheter and draw the water off. I want you to understand that the case may be treated in two ways, and that it is really a matter of convenience which method you should adopt.

These are cases of spasmodic or congestive retention; it is a much more serious matter when you have a patient who has a tight organic stricture. The patient has perhaps had a stricture for years past, and he has been treated two or three times by different surgeons. People get very callous and careless about their urethras, and neglect themselves, and in that way the stricture gets tighter and tighter until at last some day, or more probably some night, the patient finds that he cannot make water at all, and he then sends for you to give him relief. These are confessedly very difficult cases to treat. You do not know anything previously about the man, you do not know the condition of the urethra, you have merely his story to guide you; do not therefore be deceived by what he tells you; do not begin at once with very small catheters, but always with a fair average size. For that purpose I do not know that you can do better than use the French instruments, the olivary catheters, beginning with a medium-sized catheter just to find out how the land lies. You pass that steadily down the urethra until it is stopped; then you carefully make a little pressure and try and get it through. If you find that it is of no use you abandon the attempt, and you steadily try one instrument after another until you get to the very smallest. I quite allow that with care and skill—it requires a very considerable amount of skill—you may insinuate through the stricture such a very small catheter as this which I hold in my hand, a catheter equal to about No. 0 on the English scale (it is No. 1 on the French scale), the smallest catheter I think in use. You may not succeed, and then you must have recourse to silver instruments. I know that silver instruments are not in fashion at the present time. Some surgeons hardly ever use them, but I think you ought to be able to use silver instruments, and that you may use them quite as safely as the soft ones. I presume, therefore, that having failed with these soft instruments, you try a silver one, Nos. 3, 2, or 1, or No. 0 if you have it, and do your very best to get it in. And let me say, without wishing to urge any violence whatever, that you must bear in mind that in these hard strictures you must use a little steady pressure, but no force, and you must be careful that the catheter does not suddenly slip from your fingers, and make a long false passage.

Then having tried all you know in the way of catheters, you put the patient under chloroform, and that for this reason: in all these cases of retention there is a certain amount of spasm; and if you overcome that spasm by chloroform, you get the catheter in more easily than you can do without it. Of course there is the mechanical obstruction, but there is a certain amount of spasm with it. When the patient is under chloroform he is necessarily recumbent, and then you have the advantage of putting your finger into the rectum, and seeing that your catheter is not in a false passage; you can feel it going in the proper direction, and perhaps assist it a little in passing in. Now if you have a case of that kind, and everything has been done in the way of catheterising that is possible, and yet the patient is not relieved, the bladder is nearly or half way up to the umbilicus, and the patient is in considerable distress. It is clear that the patient cannot be left, because, if he is, there will be a risk of the urethra giving way at some point, and possibly of the bladder giving way, but that is, of course, an extremely rare occurrence. Therefore it resolves itself into what is the safest thing to be done in the case of retention of urine with an impassable stricture. I have no hesitation in saying that the safest thing is to aspirate the bladder above the pubes, and you cannot well go wrong if the aspirator is in proper order, with a clean needle, which you take every precaution to make aseptic, carefully carbolicising the whole machinery, and washing the skin of the abdomen with carbolic acid, then puncturing a little way above the pubes. That operation seems to be

attended with as little risk as possible. Yet, a year or two ago, there was a paper read at one of the medical societies by an eminent surgeon, in which he endeavoured to show that this operation was not free from risk. The point was this: that if you let out putrid urine in that way, it was possible that you might get some of the urine into the cellular tissue about the pelvis, and mischief might occur. I think he was quite right; it is quite conceivable; but fortunately in these cases of retention, the urine is not putrid; it is usually a case in which the urine is fairly healthy, and, therefore, you may hope that, in the majority of cases, you will not run much risk in aspirating the bladder above the pubes. At all events, I recommend you to do that rather than any serious operation.

But, naturally, aspiration is only a temporary arrangement. You would not think of keeping the aspirator needle in. You merely draw off the urine, and then hope that the difficulty may be overcome in other ways. When you have aspirated once or twice, it might be better to puncture above the pubes with an ordinary trocar and cannula somewhat curved, leaving the cannula in (there is a cannula made on purpose). Patients may no doubt go on for years making water in that way through a permanent opening above the pubes. On the other hand, if you do not choose to adopt that method, there is the much older plan of puncturing through the rectum. There are disadvantages in that, and I think that surgeons generally have given it up. It is still used at Guy's Hospital, I believe; because Mr. Cock, formerly surgeon there, specially advocated it, but I do not think that that method is often adopted elsewhere. The objection to puncturing by the rectum is this. In the first place, it is inconvenient to keep a tube in the rectum, and in the next place there is a risk of wounding the peritoneum. The peritoneum is not always so high as it ought to be. Sometimes it comes lower down to the prostate than you would expect, and, under those circumstances, if you puncture it, you are sure to get peritonitis. Then, again, it is possible that a little urine may get out, and then you may get cellulitis about the pelvis. So that there are several drawbacks, and altogether the operation is, I think, practically superseded by the other proceeding.

If a man has a tight stricture in the perineum which is impassable, and if he has also retention of urine, of course it becomes a question whether you would not be wise, under some circumstances, to operate in the perineum and divide the stricture, and, at the same time, relieve the retention. I quite allow that, under favourable circumstances, that is, perhaps, the best proceeding. I mean, of course, where you have plenty of assistance, plenty of good light, and every convenience for an operation such as we have in hospital practice. In such a case an operation in the perineum is perfectly justifiable, and is sometimes better than anything else. But under the circumstances that many of you will have to deal with, where the patient is entirely isolated, and where you have no conveniences and no assistance, I should strongly advise you not to attempt any perineal operation. It is always a difficult proceeding, and sometimes it is an extremely arduous operation to perform.

Now, if you elect to do any perineal operation in case of retention of the urine, I should strongly recommend you to do Wheelhouse's perineal section, which is, in fact, the only operation that would be performed at the present day. Let me point out shortly what the operation is before I go on to other matters. You take a staff, like that which I hold in my hand, with a little knob at the end, and you pass it down to the obstruction, which is, of course, usually about the bulb. The patient is then tied up in the lithotomy position, and the surgeon opens the urethra quite in front of the stricture, presumably a healthy portion of the urethra. Having opened it upon the groove, the assistant turns the instrument round, and hooks it up so as to draw up the urethra a little, and with two pairs of forceps the urethra is held open on each side. Having exposed the interior of the urethra the operator has to find the orifice, which is, of course, exceedingly small. He takes a probe-pointed director, and, having passed the probe through the stricture, he pushes the director portion a little on into it, and with a bistoury he divides it. Then, still keeping the groove there, he passes a gorget into the stricture and along the urethra, and so guides the catheter through the whole length of the urethra, and in that way into the bladder.

And now I come to the last type of retention, at the other extremity of life—in old age; and let me say that this is a class of case that you are likely to meet with in private practice. Old people get retention of urine, not because they have

enlarged prostates merely, for many men have enlarged prostates after sixty, and plenty of them go through life afterwards without having retention of urine. The reason why they have retention is because they get congestion of a large prostate, arising generally from exposure to cold. What happens is this: an elderly gentleman, who has been wearing thick clothes all the day, dresses in the evening to go out to dinner, and puts on a very much thinner pair of trousers. Without the slightest reflection upon his appetite in any sense whatever, he is exposed to cold and gets congestion of the prostate. It is very uncharitable to say that all these elderly people have taken more wine than is good for them, or anything of the kind; it is not the fact. Perfectly reasonable and sober persons get retention simply from exposure to cold. They find, when they go to bed, that they are not able to make water; they go on trying, perhaps, to make water at intervals during the night, and late at night or early in the morning you are sent for, to find an elderly gentleman in considerable distress because he cannot relieve his bladder. Such a case as that of course requires a little careful management; but, after all, it is not a very desperate case to treat, unless, unfortunately, the old gentleman has been trying to pass a catheter for himself. The cases that give trouble are where you find that the patient has been making false passages with his own catheter. Some time or other, probably, he has been provided with a catheter, and then, getting retention of urine, he tries to use it, and he makes himself bleed. In such a case you may have a good deal of trouble; but, fortunately, in the great majority of cases no catheter has been used when you are first called in. In going to a case of that kind, of course you provide yourself with suitable instruments. You want an ordinary English catheter of good size, No. 9 or 10. A soft india-rubber catheter is also a very useful thing; and then there are the elbowed or *coudé* French catheters, such as the one you see here. Usually, with one or other of those instruments you can overcome the difficulty if you only go to work quietly and steadily, giving your catheter a good curve, and slipping it down until you come to some obstruction, then gently pressing against it till the catheter goes in. I think that the old-fashioned English catheter without a wire is the best instrument, but the fashion rather runs on these *coudés*.

Occasionally there are great difficulties. The third lobe of the prostate rises up in the floor of the urethra, and makes it exceedingly difficult to get into the bladder, and in these cases you may sometimes advantageously push the catheter in as far as it will go with the wire in it, and then withdraw the wire and push the catheter on a little further, and softened as it is by the heat of the urethra, it finds its way in. Of course you must bear in mind that the projection of the prostate may be on one side, and that is one great reason why you cannot put the catheter straight in; but you must let it fit itself as it were to the urethra, and gradually find its way into the bladder. However, with care I believe that all these cases are to be successfully treated. You may be obliged to have recourse to a silver instrument; it is better not to do so if you can help it, but anything is better than leaving the patient unrelieved; and the large, well-curved silver catheter is a very useful instrument in such cases. The size I prefer is about No. 10, and you want a good long curved catheter in these cases. Sometimes where the other catheters have been tried and failed you can pass a catheter like that without any difficulty, if you will only keep along the roof of the urethra.

Now in such a case of retention, supposing that you have drawn the water, what kind of urine will you expect to see coming through the catheter? It is always more or less bloody from the congestion of the prostate. Patients are apt to be alarmed by this, but you need not be, for it is just one of the ways by which Nature relieves the congestion of the parts, and the urine will be bloody for some few days, and then gradually will clear off, and the patient will recover his usual habits. It is also well to bear in mind that these patients do not at once recover the use of the bladder, they cannot micturate for some days. The time varies in different cases, but it is not at all uncommon to find a patient who has had a bout of prostatic retention, going on for a fortnight or three weeks before he can thoroughly make water for himself. It is always well to know that, so as to be able to warn the patient's friends that you will have to come and draw the water for him for a few days until he recovers entirely the power over the bladder.

I have now given you all the varieties of retention in the male,

from the baby to the old man. Let me say a word about retention in the female, which does occur from time to time as the result of some displacement of the uterus—the tilting backwards of the unimpregnated uterus, or possibly of the impregnated uterus, or may result from the natural changes of pregnancy. In the later weeks or days of pregnancy, when the uterus is pressing considerably upon the neck of the bladder, one can easily understand how it produces retention of urine. I simply mention it here because it is just one of those things that are apt to be overlooked. A woman makes a certain amount of water, but the bladder is more or less distended, and therefore, when you are called in to attend a labour, always ascertain whether the bladder is empty or not. I suppose there are no more unfortunate cases than those where a child's head has been allowed to press upon a full bladder, for those are the miserable cases where you have ulceration and consequent vesico-vaginal fistula. If you always pass a catheter or see that the bladder is empty in every case of labour, you will hardly ever have such a thing as vesico-vaginal fistula.

There is another class of cases of female retention which should not be treated with the catheter: that is the hysterical. Young women sometimes cannot or will not micturate, and it is a mistake to use the catheter if you can avoid it. Every now and then you are obliged to use the catheter, but it should not be used if it can be avoided, because you can really treat those cases more satisfactorily without it. The great thing is to put the treatment in the hands of a judicious matron. Let her put the patient into a bath of warm water, and then, without giving her notice, throw a bucket of cold water over her back and shoulders. The sudden shock from the cold water over the trunk will produce a violent expiratory effort, which will overcome the already relaxed muscular fibres of the sphincter vesicæ, so that the urine passes, and there is an end of the difficulty.

A DISCUSSION

ON

THE OPERATIVE TREATMENT OF CLUB-FOOT.

In the Section of Surgery at the Annual Meeting of the British Medical Association, held in Glasgow, August, 1888.

I.—MR. E. LUND, F.R.C.S.

Consulting Surgeon, Manchester Royal Infirmary.

At the request of the President of the Section, Mr. E. Lund (Manchester), opened the discussion on the subject. He thought that in guiding the discussion it would be desirable to limit it to those methods of treating talipes in its more confirmed and aggravated forms other than by merely dividing the tendons and then resorting to mechanical appliances, but rather to discuss the merits of certain recent methods of making sections of the tarsus, or by removing some of the separate tarsal bones. By way of illustrating one of these methods, Mr. Lund said it was a pleasure to him to call to mind that, just sixteen years ago, in 1872, at the Birmingham meeting of the Association, Sir William Ferguson, Bart., being President of the Section of Surgery, he (Mr. Lund) related a case in which he had then recently removed both astragali from a patient who recovered perfectly, and to-day he was able to show the same patient, now grown up to manhood, and able to walk fairly well without any perceptible deformity. Mr. Lund attached great interest to the exhibition of this case so many years after the operation, for when he was President of the Section of Surgery at the Manchester meeting he had taken for the subject of his address "The After-History of Surgical Cases," and it was by this way alone that we were able to form a right judgment of the advances made by such new modes of procedure.

II.—MR. ROBERT WILLIAM PARKER,

Surgeon to the East London Hospital for Children.

1. *Introductory.*—I consider myself honoured in being invited by the officers of this Section to take part in so important a discussion as this, and very fortunate in having a surgeon, so able as Professor Buchanan, to direct the discussion in the first place, and then to