THE BOWMAN LECTURE ON THE VALUE OF EYE SYMPTOMS IN THE LOCALISATION OF CEREBRAL DISEASE.

Delivered before the Ophthalmological Society, November 9th, 1888.

By HENRY R. SWANZY, A.M., M.B., F.R.C.S.I.,

Surgeon to the National Eye and Ear Infirmary, Dublin; Ophthalmic Surgeon to the Adelaide Hospitals, Dublin.

MR. PRESIDENT AND GENTLEMEN,—Recent advances in cerebral surgery have brought with them, in some respects, increased responsibility, and increased anxiety for the surgeon, in cases of focal cerebral disease. In these cases perhaps the gravest responsibility and anxiety is encountered at the very outset, when the regional diagnosis has to be made. It is true that our present knowledge of the anatomy, physiology, and pathology of the brain frequently enables us to diagnose the position of focal lesions; but it must be confessed there remains much to be learned before we shall be able to say that in all of the cases met with localisation can be successfully effected. Yet we may confidently hope that the difficulties which still beset this subject will gradually disappear; and we are encouraged to think so by contemplation of the marvellous progress which has been made in cerebral physiology within the last eighteen years.

In order that the desired end may the sooner be attained, it is of the utmost importance that the symptoms of each case of focal cerebral disease should be carefully noted, and that each necropsy should be conducted with the greatest precision; and in looking over the records of such cases, one cannot help observing that it is just in respect of the necropsy that most of them are defective. I do not mean merely that in many of the cases which ended fatally no necropsy was, or probably could be, made, but also that in many instances where a *post-mortem* examination was obtained the record of it affords only scant or inaccurate information as to the seat and extent of the disease.

I feel sure we shall see quicker and better progress in cerebral localisation when many physicians and surgeons recognise more fully than they yet have done that the interests of science, as well as their own reputations, are best served by entrusting the examination of the brain in all these cases to skilled pathologists and microscopists, rather than by undertaking the necropsies themselves; for it must, I think, be admitted that the successful examination of a diseased brain requires special training. The mere fact that marked focal symptoms may be caused by minute cerebral lesions is sufficient to demonstrate the importance of this view, without dwelling upon it further.

In the meantime, while our knowledge of the physiology and pathology of the brain is gradually increasing, we must be careful in our daily practice to employ all the facts which have hitherto been acquired concerning this interesting organ; and it has occurred to me that it might serve a useful purpose were I to occupy the time at my disposal this evening in reviewing the symptoms which are derivable from the eye in cases of focal cerebral disease, and in considering how far they may be utilised for the localisation of cerebral lesions. I incline to the opinion that these eye symptoms are not as much valued as they should be, perhaps because their often subtle and sometimes subjective nature renders them less readily studied than are other focal brain symptoms.

I also incline to think that eye symptoms are too often not looked for at first, but utilised rather as a *dernier ressort*. This is a serious error, for many of these symptoms are of the highest localising value; while their very nature may render their discovery in a late stage of the disease impossible. How, for example, can a field of vision be examined in a late stage of tumour of the brain when the patient is racked with pain, delirious, or half unconscious?

And, if it be remembered what a prominent rôle in cerebral regional diagnosis the process of exclusion must necessarily play, and what a very large extent of the brain is placed in relation to

the eye, it will be admitted that when a search for eye symptoms in a case of focal brain disease gives only a negative result, yet the significance of this for the diagnosis falls little short of that which belongs to a positive result.

In endeavouring to accomplish the task I have set before me, I invite your attention to the facts provided for us by clinical pathology, rather than to those of experimental comparative physiology. For, important as are experiments upon animals for our knowledge of the physiology of the human brain, yet no one will deny that the evidence of the bedside and of the *post-mortem* table, when it can be obtained, is of even greater importance. You will understand, then, why it is that I refer but little in the following to the epoch-making researches of such distinguished men as Ferrier, Schäfer, Victor Horsley, Hitzig, Munk, Flechzig, Goltz, and many others.

Gentlemen, in the details of this lecture I can hardly hope to offer anything that is new to you, yet I trust that in this respect you will extend to me your indulgence, and that you may be able to find at least some interest in the presentation of the subject as a whole.

In estimating the localising value of the focal eye symptoms which may be afforded by a given case of recent brain disease we are immediately confronted with a difficulty which is common to all focal symptoms, the difficulty, namely, of distinguishing between direct symptoms and the so-called indirect symptoms; the former, as you are aware, being those which depend upon the loss of function of the part in which the lesion is situated, and which, consequently, are the symptoms it is desirable to point out, while the indirect symptoms are not the result of the local disorganisation caused by the lesion, but of its pressure, of disturbances of circulation to which it gives rise, and, it is also thought, of inhibition effects, all of these being liable to interfere with the function of parts of the brain more or less distant from the lesion.

I agree with Dr. Gowers in thinking that the term "indirect" conveys a false conception of the nature of the symptoms to which it is applied, and I venture to suggest to you in its place the term "distant symptom" as being more suitable. It is an adaptation of the *Fernwirkung* of some German writers, and implies, I think, the one fact of which we are certain in connection with these symptoms, and no more.

We may at once say that there is nothing in the manner in which focal eye symptoms present themselves which can enable us to distinguish between the direct and distant symptoms. It is only in cases of stationary lesions, which remain under observation sufficiently long—some six weeks or more—after the first onset, that we can feel pretty sure that the symptoms before us are direct symptoms; for, after that interval, it is probable that the distant symptoms, if any have been present, will have passed away. But, in the case of a tumour, which constantly and slowly increases in size, distant symptoms may come and go, and then we can often only recognise the direct symptoms by their greater constancy. No doubt some symptoms are more often distant than others, and it will be my duty, as I go along, to endeavour to point out those which seem to be, respectively, the most and the least prone to be distant.

The opposite of those cases, in which not only direct but also distant focal symptoms are present, is afforded by cases where the lesion is latent, where it gives rise to no focal symptoms whatever. There is probably no part of the brain in which a destructive lesion may not be present without producing any focal symptoms, and this is one of the most remarkable facts connected with cerebral disease. Here the presence of diffuse symptoms renders the existence of focal brain disease indubitable, while the absence of focal symptoms renders localisation of the disease impossible. Lesions in those parts of the brain which, as a rule, give rise to eye symptoms, may sometimes be latent like lesions elsewhere. Marked disease, for example, in the occipital lobe, in the internal capsule, in the cerebral peduncle, and so on, may fail to produce the accustomed eye symptoms, or any others. It was necessary that I should refer to this interesting point, but its further consideration does not come within the scope of this lecture.

Focal eye symptoms may be divided naturally into those which depend upon disturbances in the motor apparatus of the eyeball, including the intraocular muscles, and those which depend upon disturbances in the special visual apparatus. We have also to consider symptoms due to lesion of the nerve of ordinary sensation of the surface of the eyeball.

And, first, as regards the symptoms derivable from the motor apparatus. In the cerebral cortex, centres exist for the motions of the face, arm, and leg, respectively, but no centre, in precisely the same sense, for the motions of the eyeball is present; that is to say, there is no centre in the cortex of one hemisphere a lesion of which will produce ophthalmoplegia, partial or complete, of the opposite eyeball alone, although attempts have been made to describe some such centres. All are agreed that the reason for this is, that the two eyes being associated in their motions, it is only those associated motions which are represented in the cortex.

By far the most common derangement of these associated ocular movements, as the result of cortical lesions, is the conjugate lateral deviation of the eyes to one side—that is, abnormal function of the internal rectus of the one eye, and of the external rectus of the other eye. In paralysis, the deviation being, of course, towards the side of the lesion, the eyes look at the cerebral lesion, as Prevost has expressed it; and in spasm, from the side of the lesion. In many of these cases, if the patient's intelligence be not impaired, the eyes can be moved as far as the middle line, or even further, by an effort of the will. We are not able, as yet, to say where the cortical centre for these motions resides in man, for Grasset's statement¹ that it is situated in the supra-marginal and angular gyri has not been verified.

But, did we know its position, it is not likely that much would be gained, so far as clinical localisation is concerned, for this centre seems to be extremely sensitive, and to be easily thrown out of gear by lesions of many different parts of the cortex. Conjugate deviation is, in short, very apt to be a distant symptom, especially in cerebral hæmorrhage, when it is often accompanied by a rotation of the head in the corresponding direction, and lasts only a brief time. Moreover, it is thought that, when this centre may happen to be actually involved in the lesion, its function, being largely bilateral, is rapidly taken up by the opposite hemisphere; and hence, even when conjugate lateral deviation plays the part of a direct cortical symptom, it can never be recognised as such, owing to its evanescent character.

Again, conjugate lateral deviation similar in character to that caused by a cortical lesion may proceed from a lesion of the internal capsule.

This same symptom, finally, may be caused by a lesion of the pons, involving the special nucleus for the associated motion, which is common to the sixth and third nerves, although distinct from their proper nuclei, and which probably resides in the superior olivary body. But conjugate deviation from a lesion in this locality differs from the same symptom in a case of lesion of the cortex, or of the internal capsule, inasmuch as the eyes are here turned away from the side of the lesion, towards the hemiplegic side of the body, in paralysis, and towards the side of the lesion, away from the convulsed side of the body, in irritating lesions. Gowers, in his recent exhaustive work on *Diseases of the Brain*,² points out that, in these pontine lesions the symptoms vary according as the disease does or does not involve the nucleus of the sixth nerve. In the first place, if the disease is above the nucleus of the sixth nerve, that is at or in the neighbourhood of the superior olivary body, the eyes cannot be moved towards the side of the lesion beyond the middle line, but in some cases, although the associated movement is lost, yet convergence can be effected. But, if the very nucleus of the sixth nerve be involved, there is complete loss of power of the external rectus, so that the eye deviates inwards, and cannot be rolled outwards; while the other eye can be moved by its internal rectus as far as the middle line and no further. Attention was first drawn to this form, I think, by Broadbent.³ In it the facial nerve, on the side of the paralysed sixth nerve, is often also paralysed, owing to the proximity of its fibres to the nucleus of the sixth nerve; or, because the sixth and portio dura have a common nucleus. Finally, if the fibres of the sixth nerve in the pons, after they have left their nucleus, be injured, the external rectus is paralysed, but the associated internal rectus of the other eye is not impaired in its function. Conjugate deviations due to pontine lesions differ again from those due to cortical or capsular lesions, in that they are almost always direct symptoms. Doubtless the reason for this is, that the close proximity of the two nuclear centres for the associated lateral motions of the eyes to each side, respectively, lie so close together that a distant effect can hardly be produced upon one of them alone. Yet if both nuclear centres should

happen to be paralysed, we must not at once conclude that the resulting symptom is a distant one, for, according to Wernicke,⁴ cases of simultaneous focal lesion of both of these centres have been observed. The clinical appearance is then very peculiar, the eyes being directed straight forwards, and being absolutely immovable to the right or to the left, while their upward and downward motions are preserved, as well as the motions of the upper eyelids.

Conjugate lateral deviation of the eyes, then, may assist us in the diagnosis of a cortical or of a capsular lesion from one in the pons. It may aid us in deciding in which side of the brain a lesion is situated when other symptoms are not readily observed, as in coma. And the varieties of the symptom, when it is due to disease in the pons, may enable us to form a refined diagnosis as to the precise seat of the lesion there.

As regards loss of other associated motions of the eyeball, an interesting case of loss of motion of the eyes upwards has been recorded by Gowers,⁵ in which a small tumour was found in the middle line behind the posterior quadrigeminal bodies, damaging them slightly, as well as the velum and the adjacent part of the inferior vermiciform process of the cerebellum. In a case of tubercle of the corpora quadrigemina, Hennoch⁶ observed this same defect as the first focal symptom to make its appearance.

Paralysis of the upward and downward motions of both eyeballs, sometimes with ptosis, while the lateral motions are unimpaired, may be the result of a focal lesion involving the third nerve nuclei in the floor of the Sylvian aqueduct; and, if attended by hemiplegia, the lesion involves the pyramidal tracts, probably at the level of the anterior quadrigeminal bodies, the posterior commissure, and the neighbouring part of the optic thalamus. Lang and W. A. Fitzgerald reported a case to this Society⁷ in which this symptom and hemianopsia were the two focal signs. The case rapidly recovered, leaving only homonymous insular scotomata. In their communication the authors do not profess to have localised the lesion accurately, but I think this combination of symptoms might be accounted for by a lesion involving the pulvinar and anterior corpora quadrigemina.

Loss of the power of convergence, accompanied sometimes, as in Eales's interesting case,⁸ by paralysis of accommodation, is a symptom of lesion of the posterior quadrigeminal bodies. Probably, however, this symptom is occasionally a distant one, for there are three cases published, one of them by Senator,⁹ in which the lesion was in the pons, and did not involve the corpora quadrigemina.

A symptom which is the very opposite of conjugate deviation or paralysis may be best referred to here. It consists in a deviation of one eye downwards and outwards, while its fellow is turned upwards and inwards. This remarkable, and, as yet, wholly inexplicable, symptom has only been seen with lesion of the middle cerebral peduncle, and the lesion may or may not involve the neighbouring cerebellar substance. The last-mentioned symptom, as well as loss of power of the upward, or of the upward and downward, motions of the eyeballs, are almost always direct symptoms.

We have now to consider the localising value of those paralyses of the cranial nerves for the supply of the motor apparatus of the eye other than conjugate paralyses.

As regards the third nerve, we are at once struck with the fact that ptosis, partial or complete, may be present as a focal symptom in cortical lesions—cerebral ptosis, as it is called—without any other third-nerve branch being paralysed. That a separate cortical centre for this branch of the third nerve exists, and that it innervates the muscle of the opposite side, is very probable. The existence of such a centre would not be inconsistent with the view that, as regards the motions of the eyeballs, associated centres alone are present; for although, as a rule, the elevators of the lids are associated in their motions, yet by an effort of the will most people can throw one of them into motion separately, or more than the other. No doubt the power to voluntarily innervate one levator and orbicularis alone varies in different individuals, and Wernicke¹⁰ leans to the opinion that in many persons the levator centres are practically associated centres,

¹ *Lehrbuch der Gehirnkrankheiten* Bd. i, p. 353.

² *Medical Ophthalmoscopy*, 2nd Ed., p. 340, and *Diseases of the Brain*, 1888, p. 174, foot note.

³ *Rev. Klin. Wochenschr.* 1864, No. 13.

⁴ *Trans. Ophth. Soc.*, vol. ii, p. 230.

⁵ *Trans. Ophth. Soc.*, vol. iv, p. 300.

⁶ *Arch. f. Psychiatrie*, Bd. xiv, p. 644.

⁷ *Loc. cit.*, Bd. i, p. 323.

⁸ *De la Deviation Conjugée*, Paris, 1879.

⁹ Page 167.

¹⁰ *Medical Times and Gazette*, 1872, vol. i.

and that this is the reason why cerebral ptosis is rather rare. The position of this centre is still an open question, for the view of Landouzy that it is situated in the posterior part of the inferior parietal lobule has not met with acceptance.

Ptosis, then, has no value as indicating the locality of a lesion in the cortex; but, according to Nothnagel,¹¹ it may be of use in distinguishing a cortical lesion from one situated elsewhere in the brain, for monolateral ptosis, as the only focal symptom, occurs with cortical lesions alone.

It is probable that ptosis, as the result of a cortical lesion, is a distant symptom in not a few of the cases where it is present.

Double ptosis was seen by Steffen¹² as the only focal symptom in a case of tubercular degeneration of the corpora quadrigemina, the neighbouring parts of the brain being absolutely uninvolved in the lesion. It is interesting to note that in this case the motions of the eyeballs were not impeded, nor was there any defect of vision.

It is evident that lesions causing bilateral paralysis of branches of the third nerve which are wont to be innervated together—loss of motion of the eye upwards, of motion of the eye downwards, of convergence, and double ptosis—are to be sought for in the quadrigeminal bodies. Basal lesions do not give rise to similar paralysis.

Ptosis on the side of the lesion has occasionally formed a symptom in disease of the pons, without paralysis of the other branches of the third nerve—except, sometimes, in so far as conjugate deviation is concerned—and without the third nerve being involved in the lesion. An interesting case of this kind has been published by Wernicke.¹³ Both upper lids drooped, the left more so than the right; there was also conjugate lateral deviation of the eyes to the right, the left eye being turned more inwards than the right eye was turned outwards. On an effort to look to the left the left eye did not move at all, while the right eye only reached the middle line. The whole of the left facial nerve was paralysed, and sensation on the right side of the face and head was dull. These were only the focal symptoms, and they continued until the patient's death. At no time was there any paralysis, either of motion or of sensation, in the extremities. The carefully made necropsy showed a tumour in the floor of the fourth ventricle, a little to the left of the middle line. Towards the left it did not extend beyond the lateral boundary of the pons; its posterior edge reached within $\frac{1}{2}$ centimètre of the calamus scriptorius, and its anterior edge was 1 centimètre from the corpora quadrigemina, while the width of the tumour was 2 centimètres. The growth did not involve the middle peduncle of the cerebellum, but it did extend into the inferior cerebellar peduncle. The rest of the brain was normal. The zone of nerve tissue surrounding the tumour was free from all pathological changes, nor were there any traces of pressure in the pons or in the medulla. The upper half of the tumour was found to have involved, on the left side, the nucleus said to be common (Stilling) to the facial and sixth nerves. The nucleus of the small motor root of the fifth nerve on the same side had suffered, and the fibres of the descending root of the right fifth nerve had been interrupted. The lower part of the tumour had destroyed a large part of the facial nucleus, and a great part of the internal and external auditory nuclei, and the upper end of the glossopharyngeal nucleus. The nuclei and fibres of the third nerve were not in any way implicated in the lesion. It is as yet impossible to explain the occurrence of ptosis in this and other such cases—they have all been cases of tumour—unless we regard it as a distant symptom. It is important to note that in each of these cases paralysis of the facial was a constantly associated symptom. Consequently, in this combination, even if it should be a distant symptom, ptosis may have some localising value.

Again, ptosis, by forming a factor of a crossed paralysis, may serve to localise a lesion in the crus cerebri. When the third nerve is paralysed by a lesion in this situation, it is the rule to find it paralysed as a whole, but paralysis of only some of the third-nerve branches may be produced by a lesion of the cerebral peduncle, and the branch to the levator palpebræ seems to be the one most frequently implicated alone. A case of this kind was published by Dr. Rickards,¹⁴ in which there was loss of motion and sensation of the left side of the body, with ptosis of the right eye; later on ptosis in the left eye appeared. The necropsy dis-

covered a hæmorrhage of the size of a hazel nut occupying the upper and inner surface of the crura cerebri, but involving chiefly the right crus.

Another important case of this kind (important especially on account of the accurate necropsy) was published last year by Professor Leube.¹⁵

In order now to complete this subject of ptosis as a focal symptom, I must refer to a rare form of it which has been described by Nothnagel,¹⁶ and which does not depend on a lesion of the third nerve. It may be called sympathetic, or pseudo-ptosis, and is accompanied by other eye symptoms as well as by symptoms of vasomotor paralysis of one side of the body, such as elevation of temperature and redness and œdema of the skin. In these cases, this author says, there is—1. Apparent ptosis on the paralysed side, owing to the contraction of the palpebral aperture, but the lid can be raised. 2. Contraction of the pupil on the same side. 3. A shrinking back of the eyeball into the orbit, so that it seems to have become smaller. 4. An abnormal secretion of thin mucus from the corresponding nostril, of tears from the affected eye, and of saliva from the corresponding side of the mouth. Nothnagel states he has found this train of symptoms in lesions of the corpus striatum.

A common sign of disease of the crus cerebri is what is known as crossed hemiplegia. Paralysis of the third nerve on the side of the lesion, with hemiplegia, hemianæsthesia, often facial, and sometimes hypoglossal, paralysis of the opposite side of the body, is a frequent form of it. The lesion may implicate all the branches of the third nerve or only some of them. I have already given an instance in which only the branch to the levator palpebræ is involved. But the localising value of crossed hemiplegia, as Hughlings Jackson long ago pointed out,¹⁷ depends chiefly on the hemiplegia and paralysis of the cranial nerve coming on simultaneously. If they occur at different times, they may be due to two distinct lesions, neither of which may be in the crus; for the hemiplegia might be due to a lesion in the hemisphere, and the third-nerve paralysis to a basal lesion of earlier or later date. Yet a few cases have been observed where, with a lesion in the cerebral peduncle, the third nerve paralysis preceded the hemiplegia by a considerable interval.

That basal lesions are by far the most frequent cause of paralysis of the third nerve is beyond a doubt; and here it is usual, but not constant, to find it paralysed in all its branches. The diagnosis to be made, when direct symptoms are being considered, is, for the most part, between a lesion in the crus and a lesion at the base. We cannot pretend to be able to make this diagnosis with certainty in all cases. Complete paralysis of every branch of the third nerve without any other paralysis is almost always basal; so also, as Graefe pointed out, are those cases in which, where there is hemiplegia, it is slight as compared with the degree of the third-nerve paralysis; and those cases, too, to which I have already referred, where there is an interval between the onset of the paralysis of the extremities and of the third nerve, are apt to be basal. Of course, there may be such a combination of paralysis of the other cerebral nerves with that of the third nerve as to leave no doubt with reference to the basal position of the lesion. But into all this I need not here enter.

Third-nerve symptoms—in addition to those included under the headings conjugate deviation, or paralysis and ptosis—are sometimes distant symptoms. Tumours of the cerebral hemispheres, more particularly if accompanied by violent general head symptoms, indicating probably high intracranial pressure, are the lesions most apt to produce these distant third-nerve symptoms. As a rule, the slighter the general cerebral symptoms are, the most likely are the third-nerve paralyses to be direct symptoms. This rule, indeed, applies to other, as well as to third-nerve, focal symptoms.

Paralysis of the fourth nerve, when combined with paralysis of other motor eye nerves, is difficult to recognise; and, consequently, in such cases, it supplies but little aid for localisation. Solitary paralysis of this nerve, as a symptom of cerebral focal lesion, is extremely rare. Niden has placed a case on record¹⁸ in which paralysis of one fourth nerve was the only focal symptom to which a tumour of the pineal gland, of the size of a walnut, gave rise. But the isolated fourth-nerve paralysis is more apt to be produced by a basal lesion. In combination with

¹¹ *Topische Diagnostik der Gehirnkrankheiten* p. 454.

¹² *Berl. Klin. Wochenschr.*, 1884, No. 20.

¹³ *Archiv für Psychiatrie und Nervenkrankheiten*, Bd. vii, p. 513.

¹⁴ *JOURNAL*, 1886, vol. i, p. 774.

¹⁵ *Deutsches Archiv für Klinische Medizin*, Bd. xl, No. 2, p. 217.

¹⁶ *Loc. cit.*, p. 327.

¹⁷ *Lancet*, September 6th, 1873.

¹⁸ *Centralblatt für Nervenheilkunde*, 1879, No. 8.

paralysis of the third nerve it speaks for a lesion in the cerebral peduncle, extending back to the valve of Vieussens, and has, I believe, been utilised by Meynert in this sense.

Paralysis of the sixth nerve associated with paralysis of the internal rectus branch of the third nerve has been already considered. When paralysis of the sixth nerve occurs as the only focal sign, it is probably due to disease at the base, or it is a distant symptom. I think I am correct in stating that there is no cranial nerve so liable to provide a distant symptom as the sixth. Gowers refers this liability to the lengthened course these nerves take over the most prominent part of the pons, which renders them readily affected by distant pressure. One or both nerves may in this way be paralysed. I was recently consulted in a case in which there were diffuse cerebral symptoms, which left no doubt as to the presence of an intracranial tumour. The only focal symptoms were complete paralysis of the left external rectus muscle, and very slight loss of motion upwards of the eyeball, and these had been present for only a few days. On one occasion, for a few minutes, there had been a tingling sensation in the left index finger. The shortness of the duration of the sixth-nerve paralysis rendered it, for the time, useless as a localising symptom; and that it was really only a distant symptom seemed not unlikely from the fact that, as I was informed, there had been a few days previously paralysis of the right internal rectus, of which now there was no trace. Wernicke¹⁹ states that sixth-nerve paralysis is most apt to be present as a distant symptom, when the lesion, especially a tumour, is situated in the cerebellum; differing in this way from the third nerve, which, as I have said, is more likely to give distant symptoms with a lesion in the cerebral hemisphere.

Paralysis of the sixth nerve, simultaneous in its onset with hemiplegia of the opposite side of the body, indicates a lesion in the pons, usually a hæmorrhage, on the side corresponding to the paralysed nerve. We know that the fifth and facial, and sometimes the auditory, spinal accessory, and hypoglossal nerves, may all, in varying combinations, form one of the elements in a crossed paralysis from a lesion in this position; but, in the opinion of Nothnagel,²⁰ if special localising value is to be given here to the participation of any one cranial nerve, that nerve is the sixth. The paralysis of this nerve, simultaneously with palsy of the opposite side of the body, while other conditions point to an intracerebral lesion, speaks then almost certainly for pontine disease.

Paralysis of the facial with the sixth is not an uncommon combination caused by a lesion in the pons, which at the same time produces hemiplegia of the opposite side of the body. This combination is a natural one, in view of the close relations of the nuclei of the sixth and seventh nerves. Indeed, Lockhart Clarke, Meynert, and others, as you are aware, are of opinion that there is one nucleus which is common to both nerves, a view not shared in by Gowers and others. The manner in which the root of the facial nerve winds round the sixth-nerve nucleus must also have an important bearing on the occurrence of associated paralyses of these nerves.

Hemiplegia due to a lesion of the cortical motor region, which might happen to be combined with paralysis of the sixth nerve as a distant symptom, offers no difficulty in its diagnosis from hemiplegia with sixth-nerve paralysis in pontine disease; for while the latter is a crossed paralysis, the former is homonymous.

Lagophthalmos, we know, is the eye symptom to which paralysis of the facial nerve gives rise. It is useful for localisation, inasmuch as it assists in differentiating a lesion in the internal capsule, or in the facial motor centre of the cortex, from one implicating the portio dura in the pons, as it is absent or very slight in the former cases, but very often markedly present in the latter. With a lesion in the lower part of the pons we are apt to have lagophthalmos with crossed hemiplegia; but, if the lesion be in the upper part of the pons, the fibres from the opposite side having here joined the motor tract, the hemiplegia and lagophthalmos will be homonymous.

Nystagmus, as a focal symptom, has little localising value. Indeed Nothnagel and Raehlmann agree in the view that it has no such value. It occurs as a distant symptom in many focal, as well as diffuse, cerebral diseases in various situations, but Gowers states that it is "especially common in tumours of the cerebellum."²¹

It will be convenient to refer here briefly to the eye symptoms of localising value afforded by paralysis of the fifth nerve. Similarly as the sixth nerve, crossed paralysis here points to a lesion in the pons. Moreover, when the nucleus or fibres of the nerve in the pons are diseased, neuroparalytic ophthalmia rarely supervenes; while, if the lesion be basal, the corneal affection is the rule.

The condition of the pupils is rarely of much value in regional diagnosis. Bilateral myosis is often seen with hæmorrhage in the pons, and is usually reckoned amongst the more important symptoms of the lesion; but it is by no means a constant symptom of it, the pupils here being frequently of normal size. Although myosis is most commonly the result of pontine lesions, yet it may accompany disease in other parts, as for instance, hæmorrhage in the corpus striatum, which bursts into the latter ventricle, and in meningeal hæmorrhages. Bilateral mydriasis is frequently present in apoplectic coma, without reference to any particular locality. The same holds good as regards monolateral myosis and mydriasis.

To the monolateral mydriasis, which forms one of the symptoms of paralysis of the third nerve, I shall refer only to point out that this monolateral paralytic mydriasis has greatly assisted in localising a lesion in the cerebral peduncle. Rudolf Arndt has recorded a case²² in which, among the early symptoms, the most marked were uncertainty and occasional loss of power of both lower extremities, incontinence of urine and of feces, vomiting, and frequently recurring dilatation of the left pupil, no other branch of the third nerve being at the time affected. The necropsy discovered, as had been foreseen, a tumour in the intrapeduncular space, which pressed more on the left than upon the right peduncle.

Loss of the pupillary reflex to light, apart from cases of paralysis of the third nerve, is a sign of lesion of the anterior quadrigeminal bodies, or of the optic tracts, and may be utilised to distinguish these lesions from others which may cause loss of sight by implicating both visual paths beyond the corpora quadrigemina, or both visual centres; for, in such cases, notwithstanding the amaurosis, the pupil reflex is maintained. Gudden's investigations²³ showed that there are special afferent fibres in the optic nerves and tracts for the pupil reflex, distinct from those for vision.

In cases of hemianopsia, similarly, the pupil reflex serves to establish a diagnosis between a lesion in an optic tract and one further on in the visual path, or in the visual centre, of the same side; for, if the pupil contracts actively to light concentrated on the blind side of the field, the lesion cannot be in the tract; but if it does not react, the lesion must be in the tract. It must be admitted that to decide the presence or absence of this hemiopic pupil is not always a simple matter, owing to the diffusion of the light when thrown into the eye, yet with care it may be obtained, or at least a marked difference in the promptness of reaction of the pupil, according as the light is concentrated more on the seeing or on the blind side of the retina.

And now, as to the second part of my subject—the localising symptoms derivable from the visual apparatus. Of these hemianopsia is one of the most common, as it is one of the most valuable. I may at once dismiss the three forms of hemianopsia which are known as temporal, nasal, and altitudinal, for they are only found with lesions of the chiasma, and provide a means, upon which I need not dilate, of localising those lesions at one or other side, above or below, the chiasma. But complete and absolute homonymous lateral hemianopsia is often a symptom of great localising value. It may be caused by a lesion in the cerebral cortex, or by one situated anywhere in the course of the fibres between the cerebral cortex and the optic chiasma; and, by taking concomitant symptoms into account, we are frequently enabled to say in what part of this course the lesion lies.

By complete hemianopsia we, of course, mean, that the defect in the field extends up to the vertical dividing line, and by absolute hemianopsia, a defect which involves each of the three visual perceptions—colour, light, and form.

And, first, as regards hemianopsia due to a lesion in the cerebral cortex, a lesion of the visual centre. Pathological anatomy leaves no doubt, but that, in man, the visual centre is situated in the occipital lobe rather than in the angular gyrus or elsewhere, and the evidence goes to show that the absolute optical centre chiefly

¹⁹ *Loc. cit.*, Bd. iii, p. 369.

²⁰ *Loc. cit.*, p. 159.

²¹ *Loc. cit.*, p. 193.

²² *Archiv f. Psych.*, iv, 2, p. 432.

²³ *Sitzungsber. d. Münch. Ges. f. Morphol. u. Physiol.*, 1886, p. 168.

occupies the cortex of the cuneus and of the superior occipital convolution, and also, especially in respect of the colour sense, the posterior part of the superior and inferior occipito-temporal convolutions. The most important evidence to this effect is supplied by four cases published respectively by Haab,²¹ Huguenin,²⁵ Féré,²⁶ and Seguin,²⁷ in which homonymous hemianopsia was present for a length of time before death, and in each of which the only cerebral disease consisted in a circumscribed lesion of the cuneus. Also an interesting case published by Bouveret,²⁸ in which sudden blindness of both eyes was caused by arterial thrombosis, resulting in softening of a symmetrical region on the mesial surface of each hemisphere, which included the cuneus on each side, and, it is true, a considerable extent of the occipito-temporal convolutions.

Then Berger, and Nothnagel²⁹ have each observed cases in which the lesion causing hemianopsia was restricted to the superior occipital convolution. Seguin, indeed, would confine the absolute optical centre to the cuneus, but Nothnagel seems to have good grounds for extending it over to the superior occipital convolution as well. Nothnagel points out that in nearly all the cases of extensive lesion of the occipital cortex which gave rise to hemianopsia, the cuneus and the superior occipital convolution were both implicated. He also draws attention to the fact that the middle and inferior occipital convolutions, the lingual and fusiform gyri, may all be disorganised without hemianopsia resulting. He is speaking here of absolute hemianopsia, but he does not discuss hemiachromatopsia, nor the question of separate centres for the three visual perceptions, and Verrey's case of hemiachromatopsia, to which I shall shortly again have to refer, makes it tolerably certain that the centre for the colour sense is in the posterior part of the occipito-temporal convolutions.

Yet cases of hemianopsia are on record in which the disease has attacked one or other or all of these parts, and has left the cuneus and superior occipital convolution unscathed. In some of these cases the lesion, no doubt, extended deep enough to involve the optic fibres on their way from the cuneus and superior occipital convolution; but in others it certainly did not do so. For these exceptional cases Nothnagel thinks that, as there is sometimes a departure from the usual anatomical arrangement of the convolutions of the occipital lobe, so, also, there may, in these cases, be an apparent irregularity in the distribution of its functions.

I may here mention that Reinhard³⁰ inclines to the view that at a point in the superior occipital convolution, close to the intraparietal fissure, there is a half-vision centre for the macula lutea. Should this prove to be correct, a lesion of the cortical centre for vision which included this spot should always give a hemianopsia, of which the dividing line would pass through the fixation point. These observations of Reinhard's require confirmation, for his paper, being the product of investigations made upon persons of unsound mind, is necessarily unreliable in many particulars. It is especially so as regards the field of vision, for perimetric observations could not be made.

Wilbrand, also, believes that the macula lutea is specially represented in the visual centre.

But, so far as our present knowledge goes, it seems most probable that the view held by Gowers regarding the variations in the dividing line in hemianopic fields is the correct one, namely, that the slight differences with which we are familiar in the position of this line, with reference to the fixation point and to the central vertical line of the field, depend upon slight individual differences in the decussation of the central optic fibres, and have no important localising value.

It would appear, then, that we cannot expect to be able to distinguish clinically between an absolute hemianopsia due to a lesion confined to the absolute visual centre, with or without its efferent fibres, from one due to a lesion involving at the same time other parts of the occipital cortex; and this we find to be borne out in practice.

But can we distinguish a complete and absolute hemianopsia, due to a lesion confined to the occipital lobe, from a similar defect in the field, due to a lesion in the optic radiations, internal capsule, pulvinar, or optic tract? We may conclude that the hemianopsia depends upon an occipital lesion, if it be unaccompanied

by hemiplegia, motor aphasia, or paralysis of cerebral nerves, as direct symptoms—as might occur with a lesion in the posterior limb of the internal capsule on the left side (*vide infra*)—; but, be it remembered, that one and all of these are liable to accompany lesions of the occipital lobe as distant symptoms.

Aphasia, too, occasionally accompanies right cortical hemianopsia as a direct symptom. It is not easy to offer a satisfactory explanation of this fact, unless we accept Naunyn's view,³¹ that there is sometimes a centre for speech in the region where the angular gyrus passes over to the occipital lobe, a region, consequently, very close to the centre for vision. It is also close to the centre for visual memory of words; and hence, aphasia, hemianopsia, and word-blindness—of which more later on—may be found in varying combinations in different cases. The form of aphasia, which Naunyn describes as produced by a lesion in this locality, is not altogether sensory, nor altogether motor, but is an undefined or mixed form. It is, moreover, often caused by a lesion in Broca's lobe, or by one in the centres which usually give sensory aphasia. But in about 40 per cent. of the cases, the lesion producing this undefined aphasia is, according to Naunyn, in neither of these localities, and then this part of the angular gyrus—that long-suffering region—is found to be the most common seat of disease.

Cortical hemianopsia may be a distant symptom. Gowers has observed that, at the onset of many attacks of cerebral hemorrhage, hemianopsia is present as a distant symptom of very fleeting character, so fleeting indeed that it does not complicate attempts at localisation. Except under this condition, distant hemianopsia seems to be rare, a fact which enhances the localising value of the symptom.

Cortical hemianopsia may be incomplete, but we do not as yet know that a lesion of the visual centre can be so situated as to produce loss of precisely the upper or lower half of the half field; yet clinicians should bear in mind those interesting experiments made by Schäfer³² on monkeys, which show that in these animals, as Munk had already proved for dogs, there is a correlation between the parts of the retina and of the occipital lobe. Removal of a certain part of each occipital lobe in the monkey gave rise to loss of the lower part of the field only in each eye. In future cases of incomplete cortical hemianopsia in which necropsies can be obtained, it will be important to note with the greatest possible exactness the precise extent and position of the disease.

So much for absolute hemianopsia. But the lesion may be such as to destroy only the colour centre, without reaching those for form and light. Eight cases of hemiachromatopsia are on record.³³ There are also some cases of loss of colour-vision in the whole field of each eye with retention of the form and light-senses.³⁴

Again, the form-sense may be lost in the half field along with the colour-sense, while only the light-sense is retained. Such cases are hardly less rare than the loss of the colour-sense alone. Furthermore, cases of hemianopsia are on record³⁵ in which, in part of the defect, both the colour and form-senses were absent, but the light-sense present, while in the remainder of the defect all three visual perceptions were lost. In some of these cases partial recovery took place by the light and form-senses returning, while the colour-sense still remained wanting; or, it may be, that the functions which returned had merely played the part of distant symptoms.

Until quite lately we had no necropsy in a case of loss of the colour-vision alone. But Dr. Verrey, of Neuchâtel, has been so fortunate as to be able to publish, in the last number of the *Archives d'Ophthalmologie*,³⁶ a case which practically fulfils these conditions. Dr. Verrey is also to be congratulated upon the good sense which led him to submit his valuable preparation to the skilful handling of Dr. Burkhardt. The patient was a lady, 60 years of age, who consulted Dr. Verrey for a difficulty of sight she experienced in continued reading. This was the result of an apoplectic attack, from which she had rapidly recovered, without

²⁴ *Monatsh. f. Augenhtk.*, 1882, p. 149.

²⁵ *Monatsh. f. Augenhtk.*, 1882, p. 143.

²⁶ Quoted by Seguin.

²⁷ *Archives de Neurologie*, 1886, p. 176.

²⁸ *Revue Générale d'Ophthalmologie*, November, 1887, p. 461.

²⁹ *Verhandlungen des VI Congresses für Innere Medicin*, 1887.

³⁰ *Zur Frage der Hirnlocalisation, etc.*, *Arch. f. Psych.*, Bd. xviii, 2, p. 478.

³¹ *Verhandlungen des VI Congresses für Innere Medicin zu Wiesbaden*, 1887.

³² "Experiments on Special Sense Localisation in the Cortex Cerebri of the Monkey," *Brain*, Parts 39 and 40, January, 1888.

³³ 1. Charpentier, *Thèse de Paris*, 1877. 2. Bierum, *Centralbl. f. Augenhtk.*, 1881, p. 471. 3. Samelson, *Centralbl. f. d. Med. Wissensch.*, 1881, Nos. xlvii and l. 4. Swanzy, *Trans. Ophth. Soc.*, vol. iii, p. 185, 1883. 5. Eperon, *Arch. d'Ophthal.*, Juillet, 1884. 6. Wilbrand, *loc. cit.* 7. Landolt, *De la Cécité Verbale*, 1888. 8. Verrey, *Arch. d'Ophthal.*, Juillet, 1888.

³⁴ Wilbrand, *Zur Diagnostik der Gehirnkrankheiten*, 1884, p. 24; Steffan, *Arch. Ophthal.*, xxvii, 2, p. 11.

³⁵ Wilbrand, *loc. cit.*

³⁶ T. viii, No. 4, Juillet-Août, 1888.

any other symptom of which she complained remaining. A perimetric examination showed that there was absolute colour-blindness in the whole of the right side of the field of each eye, with diminution, but no absolute loss, of the light and form-senses in the same part of each field. Dr. Verrey considers that the patient's difficulty in reading was due to cerebral asthenopia, and expressly states that she had no dyslexia, properly so-called, nor any word-blindness. He also does not think that the slight diminution in the light and form-senses in the right side of the field would account for the difficulty in reading which was present. In the course of the following twenty months, during which the patient lived and enjoyed fair general health, Dr. Verrey examined the fields several times, and found them unchanged. Finally the patient succumbed to another apoplectic attack.

The cause of death was found in a fresh hæmorrhage in the right centrum ovale and lateral ventricle; while an old hæmorrhagic cyst, in the lower part of the left occipital lobe, extending into the temporal lobe on the mesial side, explained the hemiachromatopsia. This cyst was situated between the floor of the posterior horn of the left lateral ventricle and the basal surface of the occipital lobe. It had occupied the white substance of the inferior occipital convolution, and had almost completely destroyed the white substance of the posterior extremity of the occipito-temporal convolutions, as well as that of the postero-inferior part of the cuneus. The cyst came almost to the surface of the cuneus and of the occipito-temporal convolutions, having destroyed the deeper layers of their cortex.

There can, then, be no doubt but that the centres for the colour form, and light senses are all present in the occipital lobe and, posterior end of the occipito-temporal convolutions; and it is probable that they are either, as Wilbrand suggests, arranged in layers one over the other in the cortex, or, as others think, they are placed side by side. The latter is the arrangement Dr. Verrey deduces from his case, and he thinks the colour-sense occupies the most inferior part of the occipital lobe, and probably the posterior part of the lingual and fusiform convolutions of the temporal lobe; while higher, and more towards the superior part of the occipital lobe, is situated the cortical centre for the light-sense; and probably between these two is the centre for the form-sense, for the latter was the function which, after the colour-sense, he found most affected. This agrees in the main with the view of Seguin and Nothnagel.

Relative hemianopsia can only occur with lesions of the cortex, hemianopsia from lesions elsewhere must always include all the visual perceptions. Those cases of hemianopsia, also, with some peripheral contraction of the other side of the field, are due to cortical lesions.

Hemianopsia from a lesion in the optic radiations will often be indistinguishable from the same symptom due to a cortical lesion. The defect may be incomplete, as the lesion may implicate only some of the radiating fibres; or it may be complete, if they are all involved. Pronounced distant symptoms, such as hemiplegia, hemianæsthesia, ptosis, and so on, are more apt to be caused by a lesion here than in the cortex.

A lesion in the posterior third of the posterior limb of the internal capsule—the sensory crossway—is likely to produce complete hemianopsia; because the nerve fibres are here collected together in a small space. Hemianæsthesia will be present as an accompanying direct symptom; and also, sometimes, loss of the other special senses on the opposite side from the lesion; and, should the disease extend forwards to the anterior part of the posterior limb, hemiplegia will be added as a direct symptom. Moreover, if the lesion be on the left side, motor aphasia may be present, by reason of the proximity of the path for speech on its way to the cerebral peduncle. I have already spoken of the combination of cortical hemianopsia with aphasia.

There are a few cases on record of hemianopsia caused by a lesion in the pulvinar. The symptoms in such cases strongly simulate those present in many cases of cortical hemianopsia, so that a differential localisation as regards these two positions may be impossible. The hemianopsia will be absolute, and probably complete; but lesions just in this situation seem to be very rare.

In hemianopsia due to a lesion of the optic tract the defect in the field is usually complete. The characteristic sign which enables us to localise a lesion in this position from one elsewhere causing hemianopsia is the hemiopic pupil already described.

Two cases may be mentioned which exemplify the peculiar service this pupil-symptom is sometimes capable of rendering. One of these—interesting, too, for other reasons—is published by

Brandenburg in the last volume of Graefe's *Archiv*.³⁷ In the right eye there was a hemiopic defect of the field to the right side. But the field in the left eye could not be examined, owing to complete leucoma of the cornea. The defect in the right field, therefore, might have been the representative of a bitemporal hemianopsia due to a lesion at the chiasma, in so far as the perimeter could decide. The absence of the hemiopic pupil it was which showed that the lesion was not at the chiasma, and, no doubt, if the field of the left eye could have been examined, a homonymous defect would have been found in it.

The other case, observed by Brieger, is quoted by Wernicke.³⁸ In it there was complete blindness of both eyes, and, as the pupil in a general way reacted to light, a diagnosis of double hemianopsia due to lesions of both occipital lobes was made. At the necropsy only the right occipital lobe was found diseased, while the left occipital lobe was healthy; consequently, the left-sided hemianopsia alone was so far accounted for. But the cause of the right-sided hemianopsia was soon discovered. It consisted in a degeneration of the left optic tract. If, then, during life the pupillary reaction had been more carefully examined, the existence of two lesions, one in the left tract and the other on the right side, beyond or higher than the tract of that side, could have been diagnosed: for the pupillary reflex would have been found wanting or defective if light had been concentrated on the left half of one or other retina.

Lesions of the optic tract are, of course apt to implicate the crus cerebri, but do not necessarily do so. Leber has pointed out that atrophy of the optic nerve is likely to make its appearance, and at an early stage of the case, in lesions of the tract.

Total blindness of both eyes, when it appears as a focal symptom—apart from such cases of double cortical hemianopsia, or of cortical hemianopsia combined with a lesion of the opposite optic tract, as I have already referred to—can only be due to a lesion involving the whole of the chiasma, or of both optic tracts. The great mass of clinical evidence is opposed to the idea that lesions of the corpora quadrigemina produce blindness.

I must now ask you to return with me for a moment to the occipital lobe. Visual aura has sometimes been noted as a symptom of disease of this lobe. Gowers has recorded a case³⁹ in which flashes of light were seen before both eyes, but especially before the left eye, and in which the necropsy discovered a sarcomatous tumour occupying the first and second occipital convolutions, the posterior half of the superior and inferior parietal lobes, and, on the mesial surface, the cuneus and the præcuneus. Other authors, too, have recorded occipital tumours with similar symptoms.

A very remarkable visual defect is that known as mind-blindness, or loss of visual memory—blindness, we may say, of what has long been known as "the mind's eye." Sight in the ordinary sense of the word—the reception of the retinal images by the visual centre—is unimpaired, but the psychological realisation of the retinal images is not effected. The objects are seen, but the sight of them suggests no corresponding idea in the patient's mind.

We are acquainted with several varieties or degrees of this symptom. In a well-marked case the patient may be unable to recognise the streets of the city in which he has been resident for many years, and will feel as if in a strange place; he may not know his own hall door. He may be unable to distinguish his wife from his mother, and his children may appear as strangers to him. He will be unable, when away from them, to recall to his mind's eye the appearance of the places and people that have been familiar to him all his life. Yet such a man will be perfectly capable, so far as his other intellectual faculties are concerned, of transacting business, and of entering into the enjoyments of life.

In a case which I brought before this Society five years ago, mind-blindness of a partial character was one of the symptoms; although at the time I did not sufficiently direct attention to it, for I did not myself understand its significance. The case is recorded in our *Transactions*⁴⁰ as one of hemiachromatopsia; but it was partial mind-blindness which brought the gentleman, aged 77, to me. Five months previously he had had a slight apoplectic attack, and he came complaining that ever since then he could not see people properly, although he saw everything else as plainly as ever. Then he expressed himself differently, and said

³⁷ Bd. xxxiii, 3, p. 93.

³⁸ *Loc. cit.*, Bd. iii, p. 335.

³⁹ *Lancet*, 1879, Vol. 1, p. 363.

⁴⁰ *Loc. cit.*

he did see people indeed, but did not know them, even when near to them. His own wife he would pass in the street, and if she stopped him, he would look straight in her face, she told me, without any expression of recognition in his until she spoke, and he then at once knew who it was. He said that although he saw me at that moment quite plainly, yet if he met me unexpectedly five minutes later, my appearance would be as new to him as though he had never seen me before; but he always identified people when they spoke. He was hypermetropic 1.5 dioptres, and had an acuteness of vision of $\frac{1}{2}$. With the exception of some peripheral cataract, his eyes were organically sound. In the course of the functional examination I discovered the hemiachromatopsia, which afterwards formed the main subject of my communication, although in my paper I did mention his difficulty of recognising people, and stated that I regarded it as a disturbance of a cerebral function. In addition to the difficulty of recognising well-known faces, the patient only complained of a confusion of ideas, when he made an unwonted mental effort. He did not tell me anything about colour-blindness, until he saw that I was going to test his colour-sense, and then he said he had found it most difficult to distinguish colours since his attack.

Mind-blindness is seen in cases of general paralysis, usually in the advanced stages. In one case under Wernicke's care,⁴¹ and which was examined by Schweigger—where the symptoms came on at an early stage, the patient's intelligence being still good—the remarkable circumstance was noted that, with good acuteness of vision, and without any absolute defect in the field, there were, disseminated over a great portion of the field, a number of relative scotomata, within the area of any one of which, although objects could be seen by the patient, yet he could not recognise them—could not tell what they were. The same objects in other parts of the field he was able, not only to see, but also to recognise. The region of the macula lutea did not correspond to one of these scotomata. Wernicke regrets that the colour-vision in the scotomata was not tested, and it would, indeed, have been interesting to know what its condition there may have been.

The position of the cortical centre for visual memory is still a subject of discussion. Hitherto all the necropsies have been made in cases of general paralysis, and for definite knowledge we must wait for the *post-mortem* examination of a case in which the symptom has been caused by cerebral hemorrhage. Nothnagel, Wernicke, Wilbrand, and some other writers assign this function to all, or to most, of that part of the occipital cortex which does not form the centre of vision. Gowers thinks it is either in the anterior part of the occipital lobes, or in the posterior part of the parietal lobes; but the latter, he believes, is the more probable. The curious and interesting fact that, in Charcot's case, in a case recorded by Quaglino, in one by Landolt, and in my own case, a derangement of the colour-sense came on simultaneously with mind-blindness seems strong evidence in favour of a localisation of visual memory very close to the visual centre. My case is, perhaps, of special importance in this connection, as the lesion was evidently of slight extent. Cases, too, are on record in which absolute hemianopsia accompanied mind-blindness. I should, therefore, be inclined, in the absence of a conclusive necropsy, to localise this function very close to the centre for vision, probably in the occipito-temporal lobe.

Gowers thinks that when mind-blindness results from disease of one hemisphere, it is probably transient; but my case, where the lesion was evidently in the right hemisphere, and where, even when I first saw the patient, the symptom had lasted five months, does not support this view.

It would seem that mind-blindness may be a distant symptom, for Wilbrand records a case in which it appeared to have only this significance.

I must here refer to a case reported by Dr. Macewen, of Glasgow, in his remarkable address on "The Surgery of the Brain and Spinal Cord," delivered at the last meeting of the British Medical Association.⁴² A man, who had received an injury about a year previously, suffered from deep melancholy and strong homicidal impulses, which were relieved by paroxysms of pain in the head of indefinite seat. The only localising symptom Dr. Macewen could obtain was mind-blindness, which had been present immediately after the accident, and for about two weeks subsequently. On operation, the angular gyrus was exposed, and it was found that a portion of the internal table of the skull had been detached from the outer, and had exercised pressure on the posterior por-

tion of the supra-marginal convolution, while a corner of it had penetrated and lay imbedded in the anterior portion of the angular gyrus. The bone was removed from the brain and reimplanted in proper position, after which the patient became greatly relieved in his mental state, though still excitable.

Dr. Macewen thinks, that the definite localisation in this case will assist in indicating, in man, what function the anterior portion of the angular gyrus and the posterior portion of the supra-marginal convolution subserve. The brilliant result obtained places the case almost beyond the region of any criticism; and yet I cannot help expressing the view that Dr. Macewen ventured a good deal here in adopting the mind-blindness as a localising symptom. The fact that it was present immediately after the accident for only a fortnight, although in marked degree, would raise a strong suspicion that the mind-blindness may have been a distant symptom, and nothing more. Again, I do not quite agree with Dr. Macewen in thinking his case conclusive as to the function of the injured convolutions. If visual memory be their function, why was it so completely destroyed in the first fortnight during which the lesion existed, and completely restored from that time on, although the lesion remained for a year afterwards? The centre for visual memory may, indeed, reside in the angular gyrus; but I hardly think Dr. Macewen's interesting case can be taken as a proof that it does reside there.

Word-blindness, or alexia—loss of the power of understanding printed or written speech-symbols—is held by many to be nothing more than partial mind-blindness. Gowers, Wernicke, and Wilbrand are of this opinion, and all these authors, as well as Ferrier and Broadbent, localise the lesions which produce the symptom in the angular gyrus of the left hemisphere. Nothnagel,⁴³ while he localises the centre for visual speech-symbols in this same region, dissents from the view that word-blindness is to be included in mind-blindness; and the fact that in some well-marked cases of mind-blindness there was no word-blindness might seem to support this view. But, if the centre for visual memory of every kind, except that for printed and written speech symbols, is situated in the occipital lobe, as Nothnagel himself holds, there is no reason why word-blindness should accompany mind-blindness, unless in those cases in which the lesion extends to the angular gyrus. The fact that word-blindness may exist without mind-blindness is, for a similar reason, no argument in favour of Nothnagel's opinion.

A good many cases of alexia with right hemianopsia have been recorded, and a natural explanation of this combination of symptoms is supplied by the proximity of the centre for vision to the angular gyrus. Indeed, some authors go so far as to state that hemianopsia is present in all cases of word-blindness. It is interesting to note that there is one case, reported by Dr. Borel,⁴⁴ in which, in a right-handed person, word-blindness along with mind-blindness accompanied left hemianopsia. And Landolt⁴⁵ has placed on record a case of right hemiachromatopsia in which there was word-blindness.

I may be permitted to briefly refer to a peculiar variety of alexia which has been seen in two cases, although we cannot at present utilise it for the purposes of localisation. One of these is reported by Brandenburg.⁴⁶ The patient, who was also affected with right homonymous hemianopsia, was unable to read any printed or written letters or words, yet he could read off with ease long numbers reaching to tens of thousands in Arabic characters. In the second case—quoted by Brandenburg—Joly⁴⁷ observed the same power of reading Arabic numbers, while the power of reading words and letters was lost.

I do not know that alexia has been seen as a distant symptom. The remarkable symptom termed dyslexia was first described by Berlin.⁴⁸ He has observed it in six cases, and Nieden⁴⁹ and Bruns⁵⁰ have each recorded one case of it. The patient is unable to read more than a very few words consecutively, either aloud or to himself, owing to a feeling of dislike or disgust which suddenly invades him, and which he cannot overcome. After he has read a few words, which he can understand well, he pushes the book away, or hands it to the surgeon,

⁴³ *Verhandlungen des VI Congresses für Innere Medicin zu Wiesbaden, 1887.*

⁴⁴ Landolt, *loc. cit.*

⁴⁵ *Loc. cit.*

⁴⁶ *Loc. cit.*

⁴⁷ *Amnesie et Cécité verbale, Le Scalpel*, 1183, No. 21, p. 147.

⁴⁸ *Arch. f. Psych.*, Bd. xv, p. 276, and in his monograph, *Eine besondere Art der Wortblindheit, Dysteria*, Wiesbaden, 1887.

⁴⁹ *Arch. f. Augenheilk.*, Bd. xviii, p. 162.

⁵⁰ *Neurolog. Centralbl.*, Nos. 2 and 3, 1888.

⁴¹ *Loc. cit.*, Bd. iii, p. 552.

⁴² *JOURNAL*, August 11th, 1888.

while he draws his head back, and turns it aside. After a brief interval the attempt may be renewed, but with the same result, when a few words have been read. There is no dimness of sight, defect of accommodation, or pain in the eyes or head to account for the symptom. It usually comes on suddenly, and is the first sign of serious cerebral disease, being soon followed by other symptoms, such as headache, giddiness, aphasia, hemianopsia, hemiplegia, and so on. All of these cases have ended fatally. The details of the necropsies, in those cases where they were obtained, leave much to be desired. In every instance the lesion was on the left side of the brain, the patients being all right-handed. The disease seems to have occupied chiefly the inferior parietal lobule, extending sometimes as far forwards as the inferior frontal convolution, and sometimes as far backwards as the angular gyrus. It is evident, however, that we must be content to wait for more definite information with regard to the usual seat of the lesion, before this symptom can be utilised in practice for the purposes of localisation.

In crossed amblyopia—in which the eye on the side away from the cerebral lesion is almost blind, with very contracted field, while the field of the other eye is also contracted, but in a less degree—the lesion has been found in the lower and hinder part of the inferior parietal lobule.

I must mention that so distinguished an author as Nothnagel, writing about a year ago,⁵¹ expressed his doubt as to the occurrence of crossed amblyopia. He does not think the perimetric examinations have been made in these cases with care sufficient to exclude the possibility of their being examples of incomplete hemianopsia. Nothnagel does not offer any experience of his own, and I am slow to think that observers like Ferrier, Gowers, and Sharkey could fall into such error.

In conclusion, with reference to optic neuritis, it is merely necessary to state that it has practically no localising value; nor should I think it is necessary even to do this much but that, in some recent cases of brain surgery, it seems to have been tried to utilise the optic neuritis for localisation. Optic neuritis occurs in most cases of intra-cerebral tumour, irrespective of the seat of the disease.

And now, gentlemen, I have come to the end of this lecture, and have not as yet made any reference to that distinguished man in whose honour I have the honour to address you. But, I do not believe you could wish that we should separate this evening, without your having heard from me some expression of the esteem in which we hold Sir William Bowman, nor will I be guilty of such an omission. And here I might dwell upon Bowman's scientific attainments, upon the good work he did in years gone by—not alone in ophthalmology, but also in physiology—upon all this Society owes to his prestige and to his generosity, upon his clinical knowledge and his operative skill, and upon many another topic which his name suggests. But I prefer to remind you of the high standard of professional life he has shown us, a standard which we of a younger generation must endeavour to maintain. Conscientious in his relations with his patients, honourable in his relations with his professional brethren, careful not to put himself forward in any unrecognised manner, not seeking notoriety, simple, kind, courteous, dignified, William Bowman is presented to our mind's eye as the personification of the best of those qualities which go to make an English gentleman. We are indeed privileged in being permitted during his lifetime to offer some tribute to his distinguished scientific merit; but, while recognising how much we owe to him scientifically, I think, and I believe you will admit, we should not forget how much his every-day professional life has tended to give to the noble speciality he adorns that high tone which belongs to it. Our sincere hope is that Sir William Bowman may long live to enjoy the honour conferred on him by his Queen, the love of those who are dearest to him, and the warm, heart-felt esteem of this Society.

⁵¹ Congress für Innere Medicin, Wiesbaden, 1887.

REGULATION OF THE SALE OF CARBOLIC ACID.—From the report of the meeting of the Council of the Pharmaceutical Society of Ireland, it is seen that the subject of accidental poisoning by carbolic acid has again been under consideration by that body, and that, in pursuance of the view taken with regard to it on a previous occasion, it has been resolved to urge again on the Privy Council the desirability of adding carbolic acid to the poison schedule. The proposal to do so met with the approval of the King and Queen's College of Physicians in Ireland in 1882.

GENERAL SUMMARY OF CONCLUSIONS FROM A SECOND SERIES OF ONE THOUSAND CON- SECUTIVE CASES OF ABDOMINAL SECTION.

Read before the Birmingham and Midland Counties Branch.

By LAWSON TAIT, F.R.C.S., LL.D.

Professor of Gynæcology in Queen's College, Birmingham; late President of the British Gynæcological Society; Honorary Consulting Surgeon to the Brooklyn Hospital for Women; etc.

WHEN I had the privilege some four years ago of communicating to this Society the results of my first series of 1,000 consecutive cases of abdominal section, I said that I did not know whether the general death-rate, 9.3 (the figures ought to have been 9.2, the fractional increase being an error), was high or low, for no such series had up to that time been published, nor have I seen such a series since. I thought the figure must be high, and I expressed my conviction that if I lived to complete another such series the death-rate would be very much lower, and it is with no small satisfaction that I can record a diminution from 9.2 in the first series to 5.3 in the second. I pointed out that as my first series included my early work, which had been so heavily tainted by the mortality of clamp and other blunders arising from the traditions of the elders, the mere freedom from these alone ought to give me far better results. That this prediction has been completely verified is certain, from the mere fact that my mortality in the treatment of ovarian and parovarian cystoma has fallen from 8.1 to 3.3. But that by no means represents the full gain which I can point out, for in this class of cases I had, in my previous series, a large group of uncompleted operations with a mortality of at least 50 per cent., and these have now practically disappeared. They were cases chiefly of sessile ovarian tumours and imbedded cysts of the broad ligament, the removal of which I always now complete, and thus at once diminish my mortality together with that greatest of all opprobria, imperfect work. I am now in a position to say that no cystic tumour of the abdomen exists which cannot be removed, and this is quite as true of solid tumours of a non-malignant character associated with any organs in the abdomen except the liver and spleen, and I lay far less stress upon my diminished death-rate as evidence of the improvement of my work, than upon the fact that in my second series I have only to plead guilty to 6 uncompleted operations in 1,000, whereas in my first series the number was 30, with the same mortality in each—50 per cent. Even in these six I know now, for *post-mortem* examination revealed the fact, that I could have finished the operations in at least two, perhaps in three, if I had screwed my courage up to a still harder strain, and I blame myself accordingly.

I therefore lay it down as an axiom that a man who has a large proportion of unfinished operations is a bad surgeon, worse than one with a high mortality. I may here again assert that I never in any way select my cases or refuse to operate in such as look unfavourable for any statistical results. This diminution of uncompleted operations may, I think, be fairly and entirely put down to increased experience, giving greater readiness of resource, better judgment, and increased manual dexterity; and to some extent the diminution of the death-rate may be credited in the same direction. But I am proud to be able to say that there is certainly another factor which enters largely into this increased success, and that is the fact that the mind of the general profession is now rapidly freeing itself from the shackles of the ancient authorities of abdominal surgery, and is opening itself to the new rules of progress. The most important of these are two: first, that the principles which govern our actions in general surgery must apply to the abdomen; and the second is that no doubt or difficulty of diagnosis should stand in the way of adopting that