that none such has been published. It is most important that the Vienna portrait should be kept in memory, and its features made familiar, in order that careful search should be made for others. No doubt, they illustrate new laws in reference to the disease in question, so also of rhinoscleroma.

In a portfolio which is on the table I have collected from various sources published portraits of diseases which, so far as my knowledge extends, are almost unique. We have amongst them an enormous erectile hypertrophy of the female breast (Mr. Image). Portraits of granuloma fungoides (Dr. Pringle). A most unusual case of urticaria tuberosum, by Mr. Morrant Baker. A case of dermatolysis, in which the man's ear and eyelids travelled half-way down his neck. The condition shown in this latter portrait is so extraordinary, so inconceivable without the sketch—and, I may add, so almost incredible with it—that we can but wish it had occurred within the possibilities of photography.

Here is a case of tinea, possibly favus, in which extensive disease of the skull bones was induced. A portrait by Dr. Anthony Todd Thompson, of a rapidly serpiginous form of nevus. In this other portrait we have a representation—rough, I may confess, as a work of art, but very good for the demonstration of what it is designed to show—in which, after premature ossification of the skull bones, the hydrocephalic brain grew through a hole caused by absorption of the calvaria, and hung over the temples (Fig. 6). I must not trouble you with more, my object simply being to enforce the point that of all extreme rarities portraits should be taken and conspicuously exhibited, in the hope of obtaining parallel observations.

We have in Dr. Duhring's Neoplasm a sort of type case of a new group (Fig. 2). If it be not now unique, it probably was so at the time of publication. The growths in the end killed the patient, and caused gland disease; they were therefore malignant, but they were not to be distinguished from inflammatory hypertrophies, and many underwent complete removal whilst the others were growing. The cases constitute most important links between inflammation and cancer.

It would be great omission if, in connection with the present-day development of such a museum as ours, I should say nothing as to the collection of illustrations of surgical instruments and appliances. The time at my disposal is, however, very short. I must content myself with affirming that such a collection ought certainly not to confine itself to objects of historical interest, but should aim at displaying every novelty of the day. It is only by such public display that the merits of what is good can be easily appreciated and made widely known, whilst that which is not good may be permitted to pass quietly away. At present the surgeon wishing to make himself acquainted with new instruments, etc., must visit an instrument maker's shop, or must content himself with what is shown at the annual museum of the British Medical Association. Neither of these can be considered as fully satisfying, and I cannot admit that the institution in which we are now assembled will have done all that it can for the perfecting of the art of surgery until it has within its walls a collection, easily arranged for inspection and comparison, well labelled, described, and catalogued, of all instruments and appliances, concerning the merits of which the practitioner may desire information.

It remains only, Mr. President and gentlemen, that I should now say a few words by way of summary and conclusion. The design of the Bradshaw Lecture is, I believe, to keep in memory the name of a surgeon who was proud of his profession and had faith in its aims. When you, Sir, did me the honour to convey to me your wish that I should deliver it on the present occasion, I had no difficulty in selecting a subject, than which I believe there is none other of equal importance at present. I can only hope that you have not thought it inappropriate. Apart, however, from the general interest which attaches to the development of modern museums, there were other and more temporary reasons which impelled me to take up their advocacy. It is well known that our Council contemplates large additions to our museum accommodation. The additions to our library are already complete, and have, I cannot but believe, given great and general satisfaction. When, however, it got abroad that one or more new wings were to be added to the museum, the announcement was not received by our critics with unmixed approval. It has been said that we have abundance of bottles and bones already; I have even heard it hinted that the age of museums is passed. Now, Sir, the Council of this College has, as you well know, in nowise commissioned me to defend its acts, nor do those acts

stand in the slightest need of apology from anyone. It occurred to me that I might, however, usefully employ the present occasion in the endeavour to enlist the sympathy and co-operation of the whole profession in the plans referred to. With this design I have tried to show that our museum, rich as it is, magnificent as in many departments are the accumulations which it already possesses, still stands in need of further developments. These developments are necessary in all directions, but, as being of paramount importance, and as having been comparatively neglected, I have urged the formation of clinical departments, which should be designed to assist in the diagnostic discrimination of disease.

I have in the advocacy of this object produced, I fear with some diffuseness, evidence that such a museum would be of the utmost advantage both for the education of students and the advancement of knowledge. I am sanguine that some of the facts adduced must have carried conviction to all doubters, should such be present, that it is most desirable that we should as speedily as possible form a collection as complete as possible of all that the modeller, the artist, and the photographer can do in the representation of disease as seen in the living subject. I am almost ashamed to attempt to meet the only objection which I can conceive to remain in the minds of any, that, namely, on the score of expense. If, however, this must be met, let it be urged that in the lowest and most mercantile calculations, such expenditure would pay. The victims of cancer, for instance, and of such diseases as malignant pustule, are often men in the prime of life, the fathers of families; and their death prematurely, quite apart from motives of humanity, may be easily shown to be a heavy pecuniary loss to the community. These diseases are curable only by early operation, and early operation is possible only as the outcome of early and confident diagnosis. I assert that a few good models would make such power of diagnosis a common possession of the whole profession to an extent very far greater than is at present the case. I name these two diseases as examples, but there are hundreds of others. Let us throw aside lukewarmness, and join heartily in the work. Let us stint no requisite outlay. I cannot believe that our critics wish that we should hoard our wealth; there could be no surer step downwards as regards both the repute and the usefulness of this noble College. Others there are, I doubt not, ready to emulate the liberality of Sir Erasmus Wilson if they see that the College knows how to make wealth useful, and has the moral courage to employ it. Surely it was to the high honour both of Aldrovandus and of John Hunter that they died poor. Although both enjoyed large incomes, yet both saw so clearly how money might be made to subserve science and both had such faith in science, that they spent in its service all they had. If I were asked to denote by any one fact the difference in the intensity of character of the brothers Hunter, who in most respects were much alike, I would say that William died wealthy and John almost a bankrupt. It cannot be wished that an institution like ours should show less of public spirit than has been displayed by individuals who were surrounded by family claims. If such cowardice should ever come over us, as I am convinced it never will, we shall indeed have lost sight of the example of the man whose name our museum bears, and shall have failed grievously in the duties of our trust.

THE MORTON LECTURE ON CANCER AND CANCEROUS DISEASES.

*Delivered at the Royal College of Surgeons of England on
Thursday, November 29th, 1888.*

BY SIR SPENCER WELLS, BART., F.R.C.S.,
Surgeon to the Queen's Household.

[Concluded from page 1265.]

LET me mention another matter for your consideration. We may be consulted in cases where the period for early and hopeful operation has passed by, or where, months or years after operation, recurrence has taken place—it may be that recurring growths have also been removed, once or several times—and then comes a time when a patient, exhausted by loss of blood or by profuse offensive discharge approaches the end, and further operative treatment is abandoned. Are we to be content with relieving pain, prolonging sleep, and the use of deodorising or

disinfecting appliances, or can we do more? Can we encourage the forlorn hope of the patient that even yet growth may be stopped; that some benign alteration may take place in its structure—something like the “spontaneous involution” of Virchow—some retrograde fatty change in the cells or elementary components of the tumour, while extension or growth of new cancer cells is stopped? If this could be attained before the whole body were infected it would amount to curing cancer. From time to time such cases of cure are reported; but few, if any, stand the test of accurate investigation. Virchow, however, cautions us against too great scepticism in such therapeutical investigations, and says that the hopeless condition of the patient justifies the trial of remedies of whose mode of action we have no clear idea. Whether we can go still further, and secure for healthy tissues immunity against infection from an existing cancer—a resistance to invasion—remains a subject for future inquiry. All we can yet say is, that under certain conditions the nutrition of cancer cells may be so far arrested as to lead to their complete disintegration. This is often observed in small fragments, or within a limited circumference, and Nussbaum has recently succeeded in the attempt to effect such changes in larger growths by cutting off their supply of blood. He makes deep furrows with Paquelin's cautery through the integuments and subcutaneous fat quite down to the adjacent fascia or muscles, thus cuts off all peripheral supply of blood, and so lessens the abnormal vascularity of many malignant tumours. Soft fungoid masses which bleed readily soon become firm, and the whole tumour more solid, while the general health improves and life is prolonged. And we also hear from Munich of aims at a somewhat similar object by injecting solutions of ozone into the substance of cancerous growths and the tissues surrounding them. Not only is the proliferation of the epithelial cells said to be arrested, and the formation of cicatricial tissue obtained, but the surrounding textures are protected against invasion. So long as the disease is local it must be curable. As Virchow says in concluding his last paper, “If cancer in its beginning, and often very long afterwards, is a local disease, it must be possible during this period locally to cure it.”

When a superficial cancer on a limb has returned after destruction by cauterisation or caustics, or after removal by the knife and grafting of healthy skin, and the lymphatic glands nearer the body remain free from infection, the propriety of amputating the limb must become the subject for consultation. It will sometimes be the painful duty of the surgeon to urge upon a reluctant patient to sacrifice a limb in the hope of saving life; and if this advice is followed before infection of the glands has taken place, the result has often proved the soundness of the advice; while too great delay, or want of earnestness in urging submission to so serious an alternative as the loss of a limb must always be, may lead to protracted suffering and inevitable death.

I have yet another subject for your consideration; one which the general public enters into with a good deal of warm feeling and an amount of ignorance which calls rather for instruction than blame. I mean the removal of cancerous growths by caustics instead of by the knife. On this point let me read to you what our great master, John Hunter, said just a hundred years ago¹ as to cures for cancer.

“No cure has yet been found: for what I call a cure is an alteration of the disposition and the effect of that disposition, and not the destruction of the cancerous parts. But as we have no such medicine, we are often obliged to remove cancerous parts, which extirpation, however, will often cure as well as we could do by changing the disposition and action. Arsenic seems to have some power of this kind, and its effects might be increased, by being used internally and externally; but its use is very dangerous, and I am afraid insufficient for the disease. This is a remedy which enters into the empirical nostrums which are in vogue for curing cancer, and among which Plunkett's holds the highest rank. But this is no new discovery, for Senertus, who lived the Lord knows how long ago, mentions a Roderiguez and Flusius, who obtained considerable fame and fortune by such a composition. I was desired to meet Mr. Plunkett, to decide on the propriety of using his medicine in a particular case. I have no objection to meet anybody. It was the young one; the old one is dead, and might have died himself of a cancer for aught I know. I asked him what he intended to do with his medicine. He said, ‘To cure the patient.’ ‘Let me know what you mean by that. Do you mean to alter the diseased state of the parts, or do you mean by your medicine to remove the parts diseased?’ ‘I mean

to destroy them,’ he replied. ‘Well, then, that is nothing more than I or any other surgeon can do with less pain to the patient.’ Poor Woollett, the engraver, died under one of these cancer cures; he was under my care when this person took him in hand. He had been a life-guardsmen, I think, and had got a never-failing receipt. I continued to call on Woollett as a friend, and received great accounts of the good effects, upon hearing which I said, if the man would give me leave to watch regularly the appearance of the cancer and see myself the good effects, and should be satisfied of its curing only that cancer (mind, not by destroying it), I would exert all my power to make him the richest man in the kingdom. But he would have nothing to do with me, and tortured poor Woollett for some time, till at last I heard the sound testicle was gone, and at length he died.”

How this story of John Hunter reminds us of much that goes on every year in this great city. More than thirty years ago, in 1857, I published a lecture on the cancer cures of that day—Pattison and Fell, Landolfi and the Black Doctor, Beveridge, and the Rev. Hugh Reed. I believe the exposure of Mr. Reed's practice did something to prevent other cancer cures from doing much harm in London since his disappearance. But there are still irregular practitioners heard of who follow the line of Plunkett, Pattison, and Fell, who profess not only to destroy a cancerous tumour, but also what they call its roots, and thus prevent a return; or, as Fell professed, to destroy the “tendency existing in many cases in the constitution for the reproduction of cancerous cells,” and to “destroy the cancerous diathesis.” It is curious to compare this claim with John Hunter's speculations on the use of arsenic to “destroy the cancerous disposition,” and Dr. Creighton's remarks on arsenic among other alterative or “habit breaking” drugs, metals, or metalloidal substances, having elective affinities for particular tissues, and his views of cancer as an acquired habit of the tissues, “a habit that might be broken if we only knew how.” The inhalation of oxygen and the use of oxygenated water as a beverage have been supported upon somewhat similar grounds, and did no harm beyond disappointing the patient. But, as a fashion, its day was short. The use of methodical compression, which often did good, has led to the more modern massage, which has already done much harm.

On looking over my old lecture, I find this sentence: “I should not be at all surprised to hear that the next great empiric who appears in London will profess to cure cancer by galvanism.” I have been surprised that this advent has been so long delayed, although there have been proofs of late that female galvanic doctors are at work, and others calling themselves electro-homoeopaths, doing some harm, but not deserving the title of great empirics. I heard only last week of certain cancer cures whose headquarters are at Brussels, but who have correspondents in London and Southampton, whose practice leads to results quite as bad as any of those of Pattison and Fell. I must not pursue this rather tempting subject further than to repeat my assertion that “we have no reason to fear a comparison between what we can do by fair and open means, and what really can be done, or ever has been done, by any cancer curer or any secret remedy.”

I wished to say something more as to the use of caustics rather than the knife, scissors, or cautery, but I have only time to add that potassa fusa has been largely used in this way, and that I have heard of, but never seen, very good results following; and I have seen several cases treated by the late Dr. Wynn Williams with bromine. But not one ended satisfactorily, although temporary good was done. On the table is a very remarkable specimen of the entire uterus, which sloughed away after the use of chloride of zinc. Dr. Marion Sims carefully scraped away with his curette all the diseased structure till he came upon hard uterine tissue. Then he applied persulphate of iron upon cotton-wool, left this for twenty-four hours, removed the styptic plug, applied another plug of wool charged with a strong solution of chloride of zinc, and left this for four days. After removing this plug antiseptic vaginal injections were used for several days, and then I found this mass quite loose in the vagina. It consists of the entire uterus, and when fresh it was quite evident that some of the peritoneal coat of the fundus came away with it. The patient suffered a great deal, and died about two months after the application of the caustic; sooner, I believe, than if she had been left alone.

More than ten years ago, in June, 1878, lecturing in this theatre, I gave a short account of Freund's method of entirely removing a cancerous uterus through an incision in the abdominal wall, and referred to Blundell's four cases of removal through the vagina.

¹ Palmer's edition of his works, vol. i, page 625.

I then said I thought it would be very rarely that an entire uterus affected by cancer would be removed in this way, and I need now only add that the results of the practice have been so disappointing, that it may be said to be practically abandoned. I have only twice removed the entire uterus by laparotomy, and the specimens in both cases are on the table before you. One is of some historical interest as having been the first instance in this country of excision of the entire gravid uterus. The patient was six months pregnant, and suffered from epithelioma of the cervix uteri. You see the entire uterus and the ovaries just as they were removed, and the opening through which the foetus was extracted. The patient recovered from the operation, and was in fairly good health for several months; but the disease recurred in the left iliac fossa, and death took place thirteen months after operation.

The other case was much more satisfactory. A patient, at the full term of pregnancy, had her pelvis so blocked up by a uterine tumour that delivery was impossible. You see the uterus, which I removed with both ovaries, and the cavity from which a living child was drawn out. Recovery was as rapid as after a natural labour, and I received a photograph of mother and child a year after the operation, both being quite well. I need say no more of these cases, as they have both been published, except that they have proved that, in case of need, the entire uterus containing a foetus or a viable child, with considerable tumours, and with both ovaries may be successfully removed through the abdominal wall. But excision of the uterus by laparotomy must be an exceptional proceeding, as removal through the vagina is a much more successful operation. It did not attract serious attention in England until after a discussion at the Obstetrical Society in March, 1885, where it may be said to have been generally condemned. In the report of that discussion I find the following remarks of my own:—

"It is to be hoped that at least as good results may be obtained here as in Germany, and that condemnation of the principle of the operation will not be the verdict of the Obstetrical Society. Have we sufficient facts before us to justify such a verdict? Admitting that abdominal extirpations have resulted in a mortality of 72 per cent., and vaginal extirpations of 28 per cent., is it to be supposed that improved methods will not lead to diminished mortality? Admitting that recurrence of the disease within a few months, or, at most, in from two to five years, has hitherto been the rule, may not a more accurate diagnosis, earlier operations, and improving methods lead to better results—not only to a lower death-rate, but to a retarded recurrence and sometimes to complete recovery?"

It will hardly be disputed now that the mode of operating has been improved, and that mortality has diminished; and I think I cannot employ the few minutes still allotted to me better than in rapidly sketching what I believe to be the best mode of operating, and in bringing before you the results already attained by experienced operators.

But before speaking of the removal of the entire uterus, I must allude to the removal of the diseased part only, by the operations known as infra-vaginal and supra-vaginal amputations. These diagrams show what is meant by these alternative proceedings:—

Supra-vaginal amputation of cervix. Infra-vaginal amputation of cervix.
 i o internal os. The dotted lines indicate the line of incision.

Of course it must be difficult to draw a very definite line, and say exactly where one or other of the two operations begins or ends, but these are near enough for all practical purposes, and if they are compared with this other diagram after Freund, showing what is done when the entire uterus is removed, both Fallopian tubes and both ovaries being left behind, a very good idea may be formed as to the comparative gravity of the different procedures.

Line of Ligatures in Freund's operation.

In many cases of total extirpation it may be advisable to remove both Fallopian tubes as well as the uterus, and in some cases one or both ovaries should also be removed. This will, I need hardly say, depend upon the healthy or diseased condition of the different parts.

I am sure we are not yet in a position to compare numerically the risks of partial or total excisions, either as regards the immediate danger of death from the operation, or as to the liability to recurrence at longer or shorter intervals. This must be done at some future time, when we have a larger number of cases accurately observed and faithfully recorded. At present I can only offer a general statement that in my own practice, when the disease is strictly limited to the parts near the os, I prefer infra-vaginal amputation, and I do it by the galvanic cautery *écraseur*, using the wire at a low red heat, and tightening the loop very slowly. I have removed nearly the whole of the cervix in this way without losing one drop of blood. A white, dry eschar is left, and cicatrisation follows without fever or much pain. When disease has extended rather higher, amputation must be done with knife or scissors, but it is well to cauterise the raw or bleeding surfaces by copper or gas cautery, or Paquelin's, to serve the double purpose of checking bleeding and destroying any infective cells that may possibly have invaded the tissues above the line of amputation.

After what I have said it cannot be surprising that in any case where cancerous disease has extended much higher than the os, and there is good reason to believe from the mobility of the organ that the surrounding tissues are still free from invasion, I advise total excision as the best practice; and increasing experience is materially simplifying the operation. I cannot pretend now to describe the different steps in detail, and I need only allude to the importance of preliminary disinfection of the vagina, removal of any dead or softening cauliflower excrescences, and shaving off all hair from pubes and vulva. This may be done the day before operation.

During the operation the patient is fixed in the lithotomy position by anklets and wristbands, or by a thigh crutch; and a *douche* and elastic tube are so arranged that a stream of warm, slightly carbolicised water may continually irrigate the vagina. The body is well covered, as well as the arms, legs, and feet, to prevent chilling. If the uterus is fairly mobile, no speculum (at most one or two retractors) is required. Some operators draw down the uterus by a simple or double hook, or by hooked forceps; but, if the cervix is firm, any friable part having been cleared away, the safest plan is to pass a strong wire through the cervix. This is quite safe, and not so likely to be cut as silk. Supposing that any intention of turning the fundus either forward or backward is, as it ought to be, abandoned, the cervix is pulled by the wire towards the vulvar orifice, the labia held aside, and the sides of the vagina separated if necessary. The operation is completed by different stages.

1. Division and loosening, and pushing up to the extent of half

to three-quarters of an inch of the vaginal mucous membrane behind and in front of the cervix.

2. Pushing upwards or separation of the bladder and ureters from the neck and anterior surface of the body of the uterus.

3. Opening Douglas's pouch, and introduction of a sponge to keep up the small intestines.

4. Securing the uterine arteries on both sides by ligature or by pressure forceps.

5. Division of the broad ligaments and other attachments, and removal of the uterus.

6. Cleansing vagina, removal of sponges, arranging ligature-threads or forceps, and loose plugging with iodoform gauze.

It is unnecessary to such an audience as this to enter into further detail, but I will refer briefly to two matters of great importance—the safety of the bladder and ureters, and the substitution of forcipressure for the ligature.

Sometimes the separation of the bladder from the uterus is quite easy, sometimes very difficult, and I am sure great care is always required to avoid the ureters. In several recorded cases one, and in more than one case both, ureters have been tied or divided. I need hardly remind you that one mode of death in uterine cancer is uræmic coma, caused by closing up of one or both ureters; and I show you here the parts removed after the death of a patient from uræmic coma. She was operated upon by a very experienced American surgeon for vesico-vaginal fistula; yet one ureter was included in the sutures which closed the fistula, and the other was left opening into the vagina behind the united edges of the fistula. In some recorded cases the bladder has been wounded, and in others the catheter or sound passed through the urethra, and used as a guide to the separating finger or to pull the bladder forward, has been seen through the thin muscular coat and mucous membrane of the bladder.

In one of these drawings the mode of passing one of the ligatures used to secure the uterine arteries is shown. A threaded blunt needle is guided by one finger passed into Douglas's pouch. One or more may afterwards be passed in the same way to secure the broad ligament on either side, or the same object may be gained by the use of pressure-forceps, as shown in the next drawing. These are not removed, but are left on for twenty-four, thirty-six, or forty-eight hours, until all danger of bleeding has ceased. Leopold and other German operators prefer the ligature.

Péan, Richelot, and other Frenchmen trust to the forceps. I believe I was the first to suggest their use, and one of our Fellows, Mr. Jennings, the first in this country to carry out the suggestion with success. Some warm controversy has arisen in Paris lately on this matter, and Dr. Pozzi, in a small work on vaginal hysterectomy for cancer just published, puts the historical question as follows. After quoting what I wrote in 1882,* recommending the

“securing any bleeding vessel as it is divided by pressure-forceps, not using any ligatures, but leaving the forceps hanging out of the vagina for two or three days until all danger of hæmorrhage has ceased,” Dr. Pozzi refers to Mr. Jennings's successful case in October, 1885, before M. Richelot's first case in April, 1886, but after M. Péan's first in June, 1885, which was not published until April, 1886, and concludes that, although M. Richelot has “vulgarised” forcipressure in this operation, Mr. Jennings's case had been published nine months before M. Richelot made his communication to the Academy of Medicine.

But it matters very little who first proposed or first practised a particular method. The chief thing is to decide whether it should be followed or not, and my own belief is that it must be left to the choice of the surgeon in each case. He should be prepared to secure bleeding vessels by ligature or the forceps, as he finds he can do it more easily. Forceps with a variety of curves may prove more convenient than the straight instrument shown in the drawing. It may be that ligatures will do all that is required, and the best way of applying them that of Leopold, as shown in the drawings, each ligature passing twice through the peritoneum of Douglas's pouch and the vagina, and being tied so as to lie parallel to the line of incision—not simply bringing serous and mucous edges together, but compressing in separate portions the whole of the vaginal wall. I think the quickest way to do this would be to have a long ligature threaded at each end on a small needle, which would be passed by a needle-holder from the serous outward through the mucous membrane.

When these ligatures are tied they form a chain of compression all round the edges, and may be used afterwards by loosely tying some or all of the long ends together to close the opening from the vagina into the peritoneal cavity as far as may be desirable. By crossing their handles pressure-forceps would have the same effect. My own feeling would be rather to trust as a rule to forceps, and only exceptionally to ligatures. Forcipressure is simpler, more secure, takes much less time, and saves much trouble and suffering to the patient after operation. When an operator who has prepared himself by sufficient practice on the dead body trusts to forcipressure, and has prepared everything before he begins, I feel sure that fifteen or twenty minutes would give ample time to

* *Ovarian and Uterine Tumours*, p. 525.

complete the operation if the ovaries are not disturbed. If healthy, I think they had better be left alone, and so may the Fallopian tubes. But if the disease has extended high up the uterine cavity it would be safer to remove the tubes; and this would add very little to the duration of the operation.

I shall not attempt to give the results of this operation statistically, but it is important that we should know what a few of the most experienced operators have been able to accomplish. Leopold wrote to me on the 11th of this month, stating that up to that date he had performed total extirpation of the uterus by the vagina in 83 cases with only 5 deaths, a mortality of 6 per cent. Péan, writing on the 13th, says that from 1882 to the end of 1887 he had done 22 operations, 15 successful, but with 7 deaths; but that this year he has had 16 cases, all successful and no deaths. Professor Olshausen, of Berlin, appears to have the largest experience of this operation, for he writes to me that between May, 1887, and November 5th, 1888, he had 99 cases with 13 deaths. In 1881-82, he had 22 cases with 8 deaths; from January, 1883, to April, 1887, 31 cases with 5 deaths = 53 cases in Halle with 13 deaths, to be compared with 99 in Berlin, also 13 deaths. All these were cases of total extirpation by the vagina for cancer. He has had 1 case where seven years have elapsed without recurrence, and 3 of five years and upwards. Martin, of Berlin, who had previously published 66 cases where cancer led to the operation, with 11 deaths, or 16 per cent. mortality, writes on the 16th of this month:—"Since the Washington Congress in 1887, I have performed the vaginal total operation in 22 instances" (he does not say in how many for cancer). "All recovered but one, who died of hæmorrhage. Of supravaginal amputation I performed altogether 6 with one death." It would be easy to collect a very large number of cases, such as Schroeder, 59, with 5 per cent. mortality; Fritsch, 60, 7 per cent. fatal. But these do not bring us later than 1886, and they all tell the same story of greater success following larger experience.

As to the comparative mortality of infra- and supravaginal amputation and total excision, the results vary so much in successive years and in the practice of different operators, that I have been unable to collect materials for a trustworthy contrast either as to the immediate fatality or as to the frequency of recurrence of disease, but it certainly appears that recurrence is earlier after partial than after complete operations. In some series the remarkable fact comes out that supravaginal amputation stands the test of recurrence better than complete excision. This would be inexplicable if it were not for the probability that the disease had been generally much further advanced in the cases where total excision was performed than in the others.

Our American brethren have been before us in adopting the operation of total excision of the uterus, and, as elsewhere, their results have improved with time and experience.

The cases hitherto performed in this country have not been enough to serve for any comparison of results with those abroad, but there have been many successful cases published since that of Mr. Jennings, five by Dr. Purcell, and others by Dr. Duncan, Mr. Reeves, and Dr. Sinclair, of Manchester; and they all teach the lesson that if the operation is to be successful it must be done early, before surrounding structures are involved in the disease.

There are other parts of this subject on which I should like to say much more. I wished to allude to cancerous disease of the ovary, and to say how difficult it is to decide whether an ovarian tumour is innocent or malignant, even after it has been removed and carefully examined. Now I can do no more than support the rule to operate early as the best way to avoid recurrence. I wished also to remind you how the mortality after amputation of the breast and after removal of other parts affected by cancerous disease has been so much lessened by modern sanitary science that all the old arguments against operating at all in cases of cancer require revision. I wished to say something about inadequate operations, and about operations which are unnecessarily severe—and much more than I did say against operating in hopeless cases, and in studying how to "soothe where we cannot save." Of cases which especially interest us, where cancer has affected the lips or tongue, the œsophagus, the larynx, the pylorus, the spleen or kidneys, the intestines, the bladder, and the female genital organs, I have not time to say another word.

And now, in conclusion, I will only venture to express the belief that if some who have heard or who may read this lecture are convinced that cancer is increasing, and will try to find out the cause of the increase, and why cancerous diseases are so prevalent and so fatal; and if they will also try to learn how to

prevent or to cure them, no one can forecast the result. No one could have imagined that a forgotten book of Marco Polo would rouse the spirit of research in Columbus, and lead to the discovery of a new continent; and no one can foretell what the work of one man may accomplish. It may be trifling, but

Think nought a trifle, though it small appear,
Small sands the mountain, moments make the year,
And trifles life.

Each one may bring his offering, however small, to the altar of one of the temples of Esculapius; or, if his own search for knowledge fail, he may assist in preventing the failure of others, and in guiding them along the path to full success. And, year after year, the Morton Lecture may exert some influence for good upon all mankind in all coming generations.

[Some paragraphs of the lecture were not delivered for want of time.]

OBSERVATIONS

ON

TWO CASES OF RAYNAUD'S DISEASE (SYMMETRICAL GANGRENE).

WITH CHRONOLITHOGRAPH.

Read in the Section of Medicine at the Annual Meeting of the British Medical Association, held in Glasgow, August, 1888.

By J. O. AFFLECK, M.D., F.R.C.P.E.,

Physician to the Royal Infirmary, Edinburgh; Lecturer on Practice of Physics and on Clinical Medicine in the Edinburgh Medical School.

ALTHOUGH numerous instances of the remarkable train of symptoms constituting the disease which bears the name of its eminent discoverer, M. Maurice Raynaud, have of late years been recorded, and although the subject has received considerable attention and discussion, it must be admitted that there remain many points both in its etiology and pathology which are still unsettled, and respecting which further information is much to be desired.

The main phenomena of this affection, when typically manifested, are, as is well known, the sudden occurrence, after it may be various sensory disturbances, of marked ischæmia, affecting in general portions of the extremities, particularly the points of the fingers and toes, but sometimes other parts, such as the lobes of the ears, tip of nose, etc. This displays itself in two or three stages. At first the affected parts present the appearance of so-called "dead fingers," being pallid and cold like the fingers of a corpse. Pain, often of intense character, is present; but along with this there may be numbness and other perverted sensations. The affected textures, too, are often abnormally hard and resistant. These symptoms, constituting the *syncope locale* of M. Raynaud, are held to depend upon arrest of blood-supply from spasm of the arteries. They may soon pass off, and the circulation being restored, the parts regain their normal condition and appearance. But in many cases this stage continues, with, it may be, intermissions over a series of days, and further vascular changes emerge in the form of extreme lividity, together with swelling of the affected localities, due to their engorgement with venous blood (*asphyrie locale*), which is held to depend either upon backward flow into the capillaries, or upon an actual spasm of the veins themselves arresting the return blood. Painful burning feelings and other sensory disturbances are present during this stage. The whole symptoms, however, may still pass off, or they may increase, and, with or without a further state of erythematous change, advance to gangrene. This gangrene is usually of the dry form, and superficial in extent, involving the upper layers of the skin of the points of the fingers or toes, which separate, leaving an ulcerated surface, which cicatrises with some amount of loss of substance, or, it may be, of one or more phalanges. Occasionally, but rarely, the gangrene is of much greater extent. The disease usually affects young adults, and it appears to be liable to recur, particularly on the return of any exciting cause, of which exposure to cold is one of the most potent. It may vary greatly in intensity