than any tests we can apply. I have been entirely satisfied, after a very considerable experience, with my results. The operation, under the illumination afforded by my head lantern, containing an electric lamp, is easy and safe. The drumhead is detached all around by a small tenotome, an opening for the introduction of which is previously made with a trowel-pointed knife, which may also be used to detach the fastenings of the hammer and to divide the tensor tympani muscle. The tissues in front of the incudo-stapedial joint are then pushed aside with a blunt silver probe, and any oozing blood wiped away with cotton, when the joint may be divided with the angular knife made for that purpose. The separation usually occurs with a distinct snap. The long arm of the incus, which is usually in view, is then seized with forceps, and is very easily brought away. The malleus is best secured with the foreign-body forceps (known by my name), but often it can be brought down with Toynbee's aural dressing forceps. Care must be taken else the tip of the manubrium may be caught in the bottom of the drum, and broken. Usually there is but little bleeding, though the ether narcosis under which both operations are done tends to increase the vascularity about the head. Hot water is useful in arresting bleeding, poured in on cotton-wool. It is best to place these patients on restricted diet for some weeks before the operation, and to keep them recumbent afterwards for a day at least. In some cases there is no subsequent suppuration or reproduction of the drumhead, but in a certain number, the percentage of which I cannot at the moment recall, there is more or less discharge and more or less rapid reproduction. When this occurs it is best to not interfere, but in a few months remove the membrane, which is not a painful proceeding, using cocaine if required. In all cases where there is a tendency to gradual reproduction, the edges of the advancing tissue may be turned back, so to speak, by painting the presenting membrane with collodion containing either salicylic or tannic acid.

Electric head-lantern.

There has been a reluctance on the part of some of the older otologists to enter into a somewhat new field of aural surgery, but the speaker believed we should come to it before long. In otorrhœa we may prevent the extension of purulent processes to the brain, for exacerbations can scarcely no longer occur. In sclerosing aural catarrh we may in all cases, with but few exceptions, expect to improve hearing, and in all it may confidently be promised, from the speaker's experience at least, that the further progress of deafness will be arrested.

The operation requires, after complete narcosis, from ten to twenty minutes for its completion, though in otorrhœa, owing to the greater liability to bleeding, it may sometimes require a longer time.

An alarming epidemic of measles is reported to be raging at Alton, Hants. All the public schools are closed, and over 300 cases are reported. Several deaths are said to be taking place daily.

CACTUS GRANDIFLORUS IN SOME FORMS OF HEART DISEASE.

By A. ORLANDO JONES, M.D.
Harrogate.

THE difficulty of finding a remedy for some forms of heart disease, and the fact, which is more or less recognised, that digitalis, strophanthus, and convallaria, although very valuable, are not always reliable in the varied conditions of the heart with which we meet, appeared to be a sufficient reason for bringing under notice a drug which I had found useful in many cases, namely, *cactus grandiflorus*. I do not presume that it is a panacea for all forms of disease to which that organ is liable; I only claim for it that it is likely to prove to be a useful adjunct to our resources in some conditions of the heart.

By the cases to which I propose very briefly to refer, I may be able to show that this remedy may fill a gap where other remedies appear to be lacking, for instance, the beneficial action of digitalis is often most marked where the heart has been overstimulated, as in cases of delirium tremens; and, on the other hand, its action has been not infrequently disappointing when dealing with a feeble heart, especially if that feebleness is excessive and of long duration. But the able and valuable expositions of the virtues of digitalis by Dr. Lauder Brunton and others will be so fresh in most memories that for the purpose of a slight comparison with *cactus grandiflorus* I need only briefly refer to the fact that the action of digitalis is divided into three stages.

The first is that of stimulation of the vagus. The second action is the sudden depression of the vasomotor apparatus of the renal arteries. The third stage is the depression of the vagus, exhaustion of the ganglia, weakening of the heart, and, as Dr. Mitchell Bruce very wisely puts it, the circulation begins to fail.

But with *cactus grandiflorus*, the remedy to which I wish to draw attention, the action seems to be the very opposite, that is, the final stage of the action of the drug is, that it strengthens the heart, and consequently the circulation improves, therefore, the final result of the exhibition of this drug is the reverse of that seen in digitalis.

Thus where digitalis appears to be a drug that is most applicable to sthenic or over-stimulated conditions of the heart, *cactus grandiflorus* appears to be applicable to asthenic conditions of the heart.

This *cactus grandiflorus* is a native of Jamaica and Vera Cruz. A good description of this species is given in Green's *Botanical Dictionary*. It belongs, of course, to the order Cactaceæ, the flowers being very large, the calyx measuring about a foot in diameter. It is possible that other plants of this order may be equally valuable as possessing properties that will control the heart's action, for Robert Bentley, in his valuable work on botany, states that the fruit of many species is useful in febrile diseases.

The first case to which I would draw attention was a youth of about 13 years of age, of strumous diathesis, weedy habit, very feeble, and the heart very weak. Under the steady exhibition of *cactus* the heart improved, and the patient grew to be a fairly healthy young man.

The next case I will mention was a man, aged about 60, who came to me saying that he could not continue his work owing to the suffering produced by exertion. I found him to be suffering from mitral disease, accompanied with loud mitral murmur, and gave him *cactus* and ammonia. He so far improved that he was able to resume work, and continued it for some years till he left the neighbourhood.

A third case, a more important one, came under my care about seven years ago, at the age of about 40, apparently in a far-gone and hopeless condition of enfeebled heart, liver disease, and general dropsy, extending from head to foot. The legs were very much distended with fluid; they were hard and turgid and nearly bursting. The patient had been tapped several times previous to coming to me. The heart sounds were barely audible; the pulse could scarcely be found at the wrist.

The first time of tapping, after coming under my care, the operation was performed very successfully by Mr. Nunneley, of Leeds, when twenty-six pints were drawn off from the abdomen.

During the last six years the patient has been tapped about fifty times, the quantity gradually diminishing. Mr. Nunneley,

who has always operated since I have had charge of the case, states that he is not aware of a similar case being on record.

The dropsy is now confined chiefly to the abdomen, the legs, feet, hands, and arms having become almost normal, though the thighs are distended at times. All the organs have improved in their action, but the improvement in the heart's action is the most marked. The exhibition of cactus has been continued throughout, but other drugs have been given freely as occasion has required, in order to meet the disordered condition of other organs; for instance, for a lengthened period constipation has been a very persistent symptom, refusing to yield alike to pil. hydrarg. and mist. sennæ co., belladonna or cascara, but latterly even this symptom has materially lessened.

A fourth case was that of a young woman, about 20 years of age, who suffered from a very severe attack of rheumatic fever, when, towards the end of the fever, a loud mitral murmur developed. The exhibition of cactus and alkalies resulted in recovery so far that she has been able to resume her work.

Other cases might be brought forward, but these will perhaps suffice to suggest that cactus grandiflorus offers an interesting and valuable field for research, especially in the treatment of asthenic conditions of the heart.

NEW AURAL INJECTOR AND EVACUATOR, WITH A FLEXIBLE EUSTACHIAN CATHETER.¹

By JOHN WARD COUSINS, M.D.LOND., F.R.C.S.,

Senior Surgeon to the Royal Portsmouth Hospital, and the Portsmouth and South Hants Eye and Ear Infirmary.

SINCE the publication of my paper in the JOURNAL² on Tinnitus Aurium, and its Treatment by a New Method of Alternate Injection and Evacuation of Air, I have considerably modified my apparatus, and have now adapted it for self-application. It consists of one hand ball fitted with a recoil spring, which increases the force and rapidity of its expansion (Fig. 1). The injecting

Fig. 1.

and evacuating tubes both terminate in a vulcanite joint, to which a nasal piece of soft rubber can be attached. The nasal pieces are made in different sizes, and several of them are supplied with each instrument. When medicated air is to be used, the volatile substance must be dropped upon a pledget of cotton wool placed in the vulcanite box, which is fixed on the injecting tube. In this way I use chloroform, ether, creasote, pinol, and many other remedies. For the successful application of the treatment it is essential that the patient should be instructed to practise injection and evacuation of the tympanum in the following manner:

The nasal orifices must first of all be carefully wiped dry, and the nose pieces adjusted, so as to securely and completely plug the nostrils. The nasal tubes are now held in position by the forefinger and thumb of the left hand, and the hand ball is elevated and compressed by the right, and then allowed to ex-

pand by its own elasticity. During the steady performance of these movements the patient should take a full inspiration, and while holding the breath and pressing the lips together he should make an expiratory effort, just as if he were in the act of blowing through a tube. With many patients the entrance of air into the tympanic cavity is rendered easy by the act of deglutition, for the muscular movements of the pharynx act directly upon the open ends of the Eustachian tubes. As a general rule, my patients require very little instruction, and soon become very skilful with the instrument. The alternate method presents many advantages over other forms of tubal inflation. It promotes the drainage of the tympanum and the discharge of pent-up fluid into the pharynx. It also proves of material assistance in cases of collapsed and adherent membrane. Its daily application is especially useful in the treatment of tinnitus aurium caused by abnormal tension, and also in other forms of chronic middle ear disease.

The apparatus for alternate injection and evacuation with the Eustachian catheter is represented in Fig. 2. The tubes are

Fig. 2.

joined together, and near their junction a vulcanite terminal is fixed for the insertion of the catheter. The flexible catheter is a flattened tube of soft metal, and it can be readily bent to any shape by protecting the bore with the stilette (Fig. 2). For the successful application of the Eustachian catheter, it is often necessary to adapt the instrument to the curvatures of the nose and pharynx. Sometimes the nasal passages are by conformation irregular, or they become tortuous and obstructed by morbid changes in the bone or lining membrane; in other cases the position of the tubal orifices and mucous folds of the pharynx require a special flexion of the beak. For some patients I keep a catheter always bent and ready. The instrument must be carefully washed and deodorised after each operation.

In performing the operation of alternate injection and evacuation of the tympanic cavity, the hand ball must always be attached to the catheter after its introduction. The surgeon should elevate and compress the hand ball with the right hand, and hold the catheter and close the nostrils with the left; at the same time, he must direct the patient to practise the movements already described. During the expansion of the hand ball, the catheter should be drawn outwards for half an inch, for by this manoeuvre the evacuating power will be increased. Special care must be taken to prevent twisting or kinking of the india-rubber tubes. Sometimes the point of the catheter gets lodged in the folds of the pharynx, so that the injection of air is obstructed. When this occurs, the instrument should be rotated a little, and then replaced over the Eustachian orifice. If from any cause considerable difficulty is experienced in passing the flexible metallic catheter through the nasal meatus, I substitute for it a fine vulcanite instrument, which can be bent to any curve by gentle heat.

In conclusion, I desire to mention that all the instruments described in this paper have been made for me by Messrs. Maw, Son, and Thompson, of London.

THE ACTION OF POTABLE WATERS ON LEAD

By SURGEON GILBERT KIRKER, M.D., R.N.

[Communicated by THE DIRECTOR-GENERAL OF THE MEDICAL DEPARTMENT OF THE NAVY.]

WHILE engaged in some investigations in connection with the bacteriology of drinking water last summer, I read Mr. Power's memorandum on Lead Poisoning by Certain Public Water Supplies, etc., which is contained in the report of the medical officer of the Local Government Board for 1887. My interest was particularly aroused by Mr. Power's suggestion that the seasonal variations in the action of water on lead, which had been observed by Dr. Sinclair White at Sheffield, might be owing to low organic life;

¹ Exhibited in the Section of Otolaryngology at the Annual Meeting of the British Medical Association, held in Leeds, August, 1889.

² See JOURNALS of February 26th and March 26th, 1887.