cases. Teething can only be indirectly concerned, and then as part of the developmental changes of early childhood, for attacks occur when the first dentition is over. Hot weather is not a direct cause, for though young children suffer much from various nervous diseases in hot climates, they are not specially liable to this one. The only hereditary tendency traceable in my cases is towards the arthritic; one or two of them may be neurotic, only one case, the first, developed a subsequent tuberculosis; all were absolutely free from other latent hereditary taint.

Thus two forms of spinal paralysis occur in infancy, the one, more accidental and infinitely rarer than the other, requires rest, sedatives, and delay in the use of mechanical means; the other demands mechanical aid from the earliest period to guard against associated deformities that would otherwise follow, and in this respect differs from diphtheritic paralysis, which may be expected to amend if immediate dangers be guarded against; in both local stimulation is required, and the employment of tonics, including the use of strychnine, is advantageous. Excepting the rarer instances where diphtheria leads to lesion of the nerve centre, diphtheritic paralysis is associated with a local degeneration of the small fibres only of nerve and muscle, leaving no permanent disability or deformity of the bodily frame and limbs; but in the anterior poliomyelitis of childhood an entire muscle or group of muscles is cut off from its centre of innervation, and degenerative changes that are irremediable follow, inducing collateral effects, often of the most serious consequence, many of which could be remedied or averted by timely care and treatment.

THE CAUSES OF LARYNGISMUS IN YOUNG CHILDREN, WITH SPECIAL REFERENCE TO ITS PRODUCTION BY ELONGATION OF THE UVULA.¹

By ALFRED MANTLE, M.D.,
Durham.

THE condition of laryngismus in the young is alarming both to the friends of the patient and also to the medical man. I am not now, however, going to speak of that form of laryngeal spasm in which stridor is a marked feature, and which is in my experience by no means common, but of another form, not so much characterised by stridulous breathing as by a distinct catch in the breath, bringing respiration temporarily to a standstill. I cannot do better than give an example of the class of cases of which I speak.

An infant which may be a few weeks or months old is observed suddenly to awake out of its sleep and struggle for breath. The facial muscles may show some slight convulsive disturbance, and in some the face and lips are livid, while in others they are very pale. The mother tells you she thought her child was suffocating, but that eventually it recovered its breath, sometimes, it is said, with a more or less crowing inspiration, but frequently, in the simplest form of the disease, merely a disturbance of the natural rhythm of respiration is noticeable. These attacks are said to occur many times during the day and night, and to come on without any disturbance whatever, but the excitement of laughter or crying or the effort of sucking sometimes provokes the seizures. They are common when the child first wakes in the morning.

Laryngismus is met with in the newly-born infant, but is most common during the first half year, and must be the disease which Bouchut, in his memoir on Croup, speaks of when he says, "croup is observed in newborn children and in children at the breast." Thus far, we have a spasm confined to the laryngeal muscles only; but experience teaches that, however slight this may be, if it continue day after day and week after week, other muscles will sooner or later become involved. The flexors and adductors of the hands and feet first show signs of irritation, producing carpo-pedal contractions, or that condition now known as tetany; but should the laryngismal attacks become more urgent, a condition of general eclampsia will soon follow. To put it in Dr. Cheadle's words, "Laryngismus, tetany, and general convulsions are the positive, comparative, and superlative of the convulsive state in children."² Of course it is to be understood that Dr.

Cheadle here speaks of convulsions produced by laryngeal obstruction.

What are the causes of this form of laryngeal spasm? There is no doubt that rachitis predisposes to such attacks, and it has been said by some that rickets is always present; but we know this not to be the case. The frequency of a rickety condition, however, led some to believe that the attacks were produced by pressure on the head when the child lay on its back, through the cranial bones being in a condition of defective ossification.

A much more likely cause, as Dr. Hughlings Jackson has pointed out, may exist in the weakened and deformed chest wall of rickety children interfering with the proper and complete aëration of the blood, for we believe that increased vascosity tends to continual excitement and irritation of the respiratory centres.

Anæmia, on the other hand, may be said to favour the condition, for we sometimes see these attacks during the exhaustion consequent on diarrhoea and vomiting, and when the body is ill-nourished and emaciated by defective diet in hand-fed children.

Another cause, described by Dr. Goodhart in his book on the *Diseases of Children*, is found in the epiglottis, for he states that Dr. Lees found in one of these cases the epiglottis excessively recurved in its vertical axis, as if it had been bent in half down the middle, and that thus the ary-epiglottic folds were brought almost into apposition, and a mere chink was left between them. Dr. Goodhart says that more or less of this recurvation is a common thing in infancy and early childhood, and may explain some of the cases of laryngismus which would otherwise be swept into the net of convulsive laryngismus on account of the coexistence of a very moderate rachitis. "These symptoms," says Dr. Goodhart, "gradually pass off as the child gets older, and are little relieved by treatment." Such a cause is, no doubt, very difficult to diagnose.

Then, again, we have a similar train of symptoms produced reflexly by enlarged bronchial glands irritating branches of the vagi. *Post mortem* this has been proved to be the case, but it only accounts for a very few cases of laryngismus, whilst enlargement of the thymus gland, which has also been assigned as a cause, will account for still fewer.

I wish to draw your attention to yet another cause which experience has taught me occasionally exists. Almost the first case of laryngismus I had under observation was a very anxious one. Three or four years ago I was asked to see a child 8 weeks old, decidedly rickety, and from the description given by the mother, I concluded that the child was suffering from laryngismus.

I was fortunate enough, a few days afterwards, to witness one of the attacks complained of. The child, while being nursed on its mother's knee, and I being about to examine it, suddenly stopped breathing; the eyes were fixed and directed upwards, the conjunctivæ were insensible to touch; there was lividity of the lips and face. A struggle for breath took place, followed by a shrill inspiratory sound, the attack then suddenly ending with the child crying. These attacks would come on many times, both by day and by night, the mother stating that she had always to raise the child to prevent it suffocating.

I treated the case as one of laryngismus due to rachitis, but with no good result.

On making an examination of the throat on one occasion I was much struck with the thickened and elongated condition of the uvula. The tonsils, as would be expected, were not more than visible at that age, but the uvula and palatal folds were much congested and thickened, with a viscid secretion about the parts. After applying astringents locally and administering bromides internally without any benefit, I urged upon the parents the necessity of excising the uvula, but could not obtain their consent. They consulted another medical man, but the child's condition grew worse, the attacks were more frequent and urgent, carpo-pedal contractions and general convulsions were produced, and a certain amount of exophthalmia was noticeable. I was now asked to operate. I removed the uvula, and, with the exception of one slight attack on the same evening, no further spasm took place, and the child is now a strong healthy boy.

Here was a case where a child suffered these attacks for four months, the cause evidently having been irritation by an elongated uvula in a rickety child, producing spasm of the larynx, and eventually general convulsions.

Dr. Gay, late clinical assistant at the Great Ormond Street Hospital for Children, related to me a similar case in which a very elongated uvula was noticed in a case of laryngismus. The case was sent to the surgeon to ask the advisability of removing it,

¹ Read in the Section of Diseases of Children at the Annual Meeting of the British Medical Association, held in Leeds.

² "Pathology and Treatment of Laryngismus, Tetany, and Convulsions," *Lancet*, vol. i. page 919, 1887.

The operation was not done; tetany and general convulsions eventually developed, and the child died unrelieved. I have noticed a remark by Dr. Sidney Ringer, in his *Handbook of Therapeutics*, which is the only reference I can find to the palate as a cause of laryngismus. He says:² "It may be useful here to advert to a condition not uncommon in infants. An infant in poor health frequently wakes up at night from 'a catch in the breath.' From some unexplained reason, it cannot for a time get its breath, and wakes up with a loud snore. This condition is altogether different from that of laryngismus stridulus, and the fault appears to lie in the soft palate, not in the larynx; moreover, it is not due to enlarged tonsils, as this 'catch in the breath' occurs in children of tender age, long before the morbid condition of the tonsils takes place."

I must say that I have not since seen any case with such an amount of elongation and thickening of the uvula, although, having examined a considerable number of cases, I have nearly always found the soft palate considerably congested, and I would strongly urge everyone to examine the fauces in all cases of laryngismus. It is just possible that the treatment which I have adopted in every case since, that of cold sponging, so strongly advocated by Dr. Ringer, has been the means of preventing such an aggravated condition developing, and I can strongly endorse what Dr. Ringer says of its good effects. An important part of the treatment, however, is diet and fresh air. Most of these children are hand-fed and breathing bad air; they get not only too much starchy food, but also a deficiency of albuminates and fats.

The hygienic surroundings are usually bad. It is a condition much more common in the early part of the year than at any other time (Dr. Gee says from January to the end of June), and more common among boys than girls. Several cases may occur in a family, the same causes operating in each case. The majority of such cases get well if the right treatment is adopted in time, but I would strongly advise an examination of the throat as a matter of routine in all cases of laryngismus, for occasionally the cause will be found there, and if so may easily be removed.

NOTE ON A POSSIBLE MEANS OF ARRESTING THE PROGRESS OF MYXEDEMA, CACHEXIA STRUMIPRIVA, AND ALLIED DISEASES.

By VICTOR HORSLEY, B.S., F.R.S., F.R.C.S.,

Surgeon to the National Hospital for the Paralysed and Epileptic, Assistant
Surgeon to University College Hospital, Professor of Pathology in
University College, London, etc.

It is believed, probably, by most practitioners and pathologists of the present day, that the diseases known as myxœdema, cretinism, cachexia strumipriva are the result of want of function of the thyroid gland. This position, at any rate, is that which received most support from the committee appointed by the Clinical Society under the chairmanship of Dr. Ord to inquire into the subject, and it has been endorsed by everyone who has paid personal attention to the point with the exception of Professor H. Munk and Dr. Drobnik. For the purposes of this brief note I shall assume that the view of the majority is correct.

Quite recently a very valuable paper has been written on the prominent early symptom which follows total ablation of the thyroid, namely, tetany, by Dr. Anton von Eiselsberg, Assistant in Professor Billroth's Clinic. In this monograph (entitled *Ueber Tetanie im Anschluss an Kropf-operationen*, Wien, 1890), Dr. Von Eiselsberg, after a masterly examination of this dangerous phenomenon, gives the records of an elaborate and careful series of experiments upon cats, from which he confirms the opinion referred to above in the fullest manner, and adds many exact results to the literature of the subject. I particularly wish, however, to draw attention to the cases in which he repeated and extended Professor Schiff's fundamentally important observation on the effects of transplantation of the gland into the peritoneal cavity, etc.

In 1886, I referred in this JOURNAL¹ to Professor Schiff's pioneering investigations. That observer found that in dogs "thyroidectomy loses its danger, and an essential amount of its effect, if one previously introduces and fixes in the abdominal

cavity other thyroid glands from an animal of the same species."²

Dr. von Eiselsberg, pp. 23, 24, 25, describes a number of similar experiments in the following manner: In nine animals extirpation of one (the left) thyroid gland was performed, the lobe kept warm in normal saline solution, and quickly transplanted either into a fold of mesentery or into the subperitoneal tissue. At dates varying up to three weeks the other half was removed from the neck. Of the nine animals eight died of typical symptoms of loss of the thyroid. *Post-mortem* examination showed in these eight fatal cases that the transplanted gland had not healed *in situ* or was degenerated. The ninth animal, however, survived, and increased in weight. It was killed, and a *post-mortem* examination revealed the fact that the gland transplanted into the peritoneal cavity had organised, was vascularised, and in no way degenerated. In four more cases the transplanted gland was placed between the fascia and the peritoneum. Of these, in three cases the transplantation failed, owing to the necrosis of the graft, and of course the animals succumbed. In the fourth case, however, the animal was perfectly well (and had grown) three months and a half after the extirpation. It was killed, and, on *post-mortem* examination, the gland was found in functional activity in the abdominal wall, highly vascularised, etc.

These results show conclusively that the transplanted gland, if it can be got to live after the grafting, will, in its new situation, provide for the needs of the body as well as if it were in the neck, and consequently saves the life of the animal. It seems to me that these observations of Professor Schiff, and of Dr. von Eiselsberg, are of especial value, as they suggest to my mind that possibly the diseases mentioned before, namely, myxœdema, etc., may be treated with success by transplanting thyroid tissue into the patient. In the first place these diseases are hopelessly incurable with present remedies, and consequently anything possibly remedial not endangering life is justifiable. The transplantation of a thyroid gland from one of the lower animals into the peritoneal cavity, or into the subcutaneous tissue might be followed by the successful growth of the grafted gland, and so bring about the arrest of the disease process by reason of the restoration of the lost function. Performed under strict aseptic conditions the operation would be without risk or practically inconvenience.

A far more difficult question is that involved in Professor Schiff's dictum, and which experiment will solve, namely, What kind of gland would be best for the purpose? Doubtless one from an anthropoid ape; but the infrequency of opportunity and very considerable expense would afford some obstacles (not, however, insurmountable by any means) to this proposal.

I have elsewhere shown that the thyroid gland of the sheep very closely resembles in anatomical characteristics that of man, and further that the sheep, of all animals as yet experimentally investigated in the present question, resembles man most in the duration and character of its symptoms after thyroidectomy. I would propose, therefore, when opportunity offers, to try transplanting a portion of the thyroid gland from the sheep. By a portion, I mean that possibly the grafting of one lobe or even half, would be more successful than an attempt to transfer the whole gland; and moreover, if the operation were successful, no doubt the imbedded piece would hypertrophy to meet the requirements of compensation.

A point of practical difficulty would arise when one attempted to gauge the success of the operation, although doubtless the development of the grafted gland, or the reverse, its atrophy, might to some extent be estimated by palpation. The real plan, however, would obviously be that of watching the symptoms very closely, and if no improvement resulted at the end of two or three months, the operation should be repeated.

From experimental evidence, I would venture to suggest that the progress of the anæmia would be the best indication of relief. I may perhaps be allowed to take this opportunity of correcting ideas which have been attributed to me concerning the relation of the thyroid gland to so-called blood formation, and for which I have been in consequence adversely criticised.

Attention undoubtedly in the past has been too much given to the origin of the corpuscular elements of the blood as easier of observation than the source of the components of the plasma; and, indeed, the expression hæmopoiesis, or blood formation, has provisionally been limited to speaking of the corpuscles.

In 1885, I pointed out that marked anæmia followed loss of the thyroid gland, and that this anæmia included a fall in the number

² *Handbook of Therapeutics*, 11th edition, p. 70.

¹ Brown Lectures, 1885.

² Transl., p. 440, *Revue Médicale de la Suisse Romande*, 1884.